

# Chapter-13

## Nervous System-III

Ms. Neha Goel

Associate Professor  
Rajiv Gandhi Institute of Pharmacy  
Faculty of Pharmaceutical Science & Technology  
AKS University Satna, (M.P.).

---

---

---

### Abstract

The nervous system is essential for regulating bodily functions and cognitive processes, but it can be affected by complex disorders such as schizophrenia and Alzheimer's disease. Schizophrenia is a chronic psychiatric disorder characterized by distortions in thinking, perception, emotions, and behavior. The pathophysiology involves a combination of genetic, neurochemical, and structural brain abnormalities, particularly in dopaminergic pathways. Epidemiologically, schizophrenia affects about 1% of the global population, typically emerging in late adolescence or early adulthood. Symptoms include hallucinations, delusions, disorganized speech, and impaired cognitive function. Diagnosis is based on clinical evaluation and the presence of characteristic symptoms over time. Treatment involves antipsychotic medications, psychotherapy, and social support. Complications can include disability, social isolation, and increased risk of substance abuse and suicide. Prevention strategies focus on early intervention and reducing stressors that may trigger or exacerbate symptoms.

Alzheimer's disease is a progressive neurodegenerative disorder primarily affecting the elderly, characterized by memory loss, cognitive decline, and behavioral changes. The pathophysiology involves the accumulation of amyloid-beta plaques and tau tangles in the brain, leading to neuronal death and brain atrophy. Epidemiologically, Alzheimer's disease is the most common cause of dementia, affecting millions worldwide and increasing in prevalence with age. Symptoms start with mild memory lapses and progress to severe cognitive and functional impairments. Diagnosis is made through clinical assessment, cognitive testing, and imaging studies. There is no cure, but treatments such as cholinesterase inhibitors and memantine can help manage symptoms. Complications include complete dependence on caregivers, infections, and increased mortality. Prevention focuses on lifestyle modifications like regular exercise, a healthy diet, mental stimulation, and management of cardiovascular risk factors to potentially reduce the risk of developing Alzheimer's disease.

### I. SCHIZOPHRENIA

Schizophrenia is a chronic, severe mental disorder that affects how a person thinks, feels, and behaves. Individuals with schizophrenia may seem like they have lost touch with reality,

which can be distressing for them and for those around them. Schizophrenia can also have significant impacts on the gastrointestinal (GI) system, often due to the side effects of medications and the lifestyle associated with managing a chronic mental illness.

## Pathophysiology

### Nervous System

- 1. Neurotransmitter Dysregulation:** Abnormalities in dopamine pathways are central to schizophrenia, particularly hyperactivity in the mesolimbic pathway and hypoactivity in the prefrontal cortex.
- 2. Neurodevelopmental Hypothesis:** Schizophrenia may arise from disruptions in brain development during prenatal and early life stages.
- 3. Genetic Factors:** A significant genetic component exists, with multiple genes implicated in increasing susceptibility to schizophrenia.
- 4. Structural Brain Abnormalities:** Reduced gray matter volume in the brain, enlarged ventricles, and abnormalities in hippocampal and prefrontal areas.

### Gastrointestinal System

- a. Gut-Brain Axis:** Increasing evidence suggests that gut microbiota dysbiosis may contribute to the pathophysiology of schizophrenia.
- b. Inflammation:** Systemic inflammation and elevated levels of inflammatory markers have been noted in schizophrenia, potentially influencing both brain and gut health.
- c. Medication Effects:** Antipsychotic medications can lead to various GI issues, including weight gain, metabolic syndrome, and altered gut motility.

### Epidemiology

- a. Prevalence:** Schizophrenia affects approximately 1% of the global population.
- b. Onset:** Typically manifests in late adolescence to early adulthood, with males often presenting earlier than females.
- c. Risk Factors:** Family history, prenatal exposure to infections or malnutrition, urban upbringing, and cannabis use during adolescence.

## Symptoms and Complications

### Nervous System

- a. Positive Symptoms:** Hallucinations, delusions, thought disorders, movement disorders.
- b. Negative Symptoms:** Reduced expression of emotions, lack of pleasure, difficulty beginning and sustaining activities, reduced speaking.
- c. Cognitive Symptoms:** Impaired executive function, attention, working memory.

### Gastrointestinal System

- a. GI Symptoms:** Constipation, diarrhea, bloating, abdominal pain, nausea.
- b. Metabolic Issues:** Weight gain, increased risk of diabetes and cardiovascular diseases due to antipsychotic medications.
- c. Nutritional Deficiencies:** Poor diet and reduced appetite, leading to potential vitamin and mineral deficiencies.

### Diagnosis

- a. Clinical Assessment:** Detailed patient history and symptom evaluation.

- b. **Psychiatric Evaluation:** Structured interviews and use of standardized diagnostic tools like the DSM-5 or ICD-10 criteria.
- c. **Neuroimaging:** MRI or CT scans to rule out other conditions and to observe brain structural abnormalities.
- d. **Laboratory Tests:** Blood tests to exclude other medical causes of symptoms and to monitor the side effects of medications.

## Treatment

### Nervous System

- a. **Pharmacotherapy:** Antipsychotics (first-generation and second-generation), mood stabilizers, antidepressants.
- b. **Psychotherapy:** Cognitive-behavioral therapy (CBT), supportive therapy, family therapy.
- c. **Psychosocial Interventions:** Social skills training, vocational rehabilitation, community support programs.
- d. **Lifestyle Changes:** Regular physical activity, healthy diet, and adequate sleep.

### Gastrointestinal System

- a. **Medications:** Probiotics, laxatives, antidiarrheals, and medications to manage metabolic side effects.
- b. **Dietary Management:** Balanced diet rich in fiber, regular meal schedules, and avoiding foods that trigger GI symptoms.
- c. **Behavioral Therapy:** Stress management techniques, education on healthy eating habits.

### Complications

- a. **Chronic Psychiatric Symptoms:** Persistent and severe symptoms can lead to significant disability.
- b. **Physical Health Issues:** Increased risk of cardiovascular diseases, diabetes, and other metabolic disorders.
- c. **Substance Abuse:** Higher likelihood of alcohol and drug misuse, often as a form of self-medication.
- d. **Social and Occupational Impairment:** Difficulty maintaining relationships, education, and employment.
- e. **Suicide Risk:** Increased risk of suicide compared to the general population.

### Prevention

- a. **Early Intervention:** Identifying and treating symptoms early can improve long-term outcomes.
- b. **Genetic Counseling:** For individuals with a family history of schizophrenia.
- c. **Healthy Lifestyle:** Regular exercise, balanced diet, and stress management techniques.
- d. **Avoiding Substance Use:** Particularly cannabis and other recreational drugs during adolescence.
- e. **Regular Medical Check-ups:** Monitoring physical health and managing medication side effects.
- f. **Education and Awareness:** Promoting mental health awareness and reducing stigma.

## II. ALZHEIMER'S DISEASE

Alzheimer's disease (AD) is a progressive neurodegenerative disorder that primarily affects memory, thinking skills, and behavior. It is the most common cause of dementia, accounting for 60-70% of cases.

### Introduction

Alzheimer's disease gradually destroys brain cells, leading to cognitive decline and memory loss. It impacts daily functioning and eventually interferes with the ability to carry out simple tasks.

### Pathophysiology

The exact cause of Alzheimer's disease is not fully understood, but it involves several key pathological changes in the brain:

- 1. Amyloid Plaques:** Abnormal deposits of beta-amyloid protein form plaques between neurons.
- 2. Neurofibrillary Tangles:** Twisted fibers of tau protein accumulate inside neurons, disrupting cell transport systems.
- 3. Neuronal Loss:** Progressive death of nerve cells (neurons) leads to brain shrinkage (atrophy) over time, especially in the hippocampus and cortex.
- 4. Neurotransmitter Disruption:** Decreased levels of neurotransmitters, including acetylcholine, which is crucial for memory and learning.
- 5. Inflammation and Oxidative Stress:** Chronic inflammation and oxidative damage contribute to neuronal dysfunction and death.

### Epidemiology

- a. Prevalence:** Alzheimer's disease affects millions of people worldwide, with numbers expected to rise as the population ages.
- b. Age Distribution:** Most common in older adults, with risk increasing significantly after the age of 65.
- c. Gender:** Women are more likely to develop Alzheimer's disease compared to men, partly due to their longer lifespan.

### Symptoms and Complications

#### a. Early Symptoms

- Mild memory loss, especially of recent events.
- Difficulty performing familiar tasks.
- Problems with language (aphasia).
- Disorientation to time and place.
- Poor judgment.
- Changes in mood or behavior.

#### b. Advanced Symptoms:

- Severe memory loss, including forgetting names of family members or familiar objects.
- Inability to recognize faces or places.
- Difficulty speaking, swallowing, and walking.

- Behavioral changes, including agitation, aggression, and hallucinations.
  - Dependency on others for daily care.
- c. Complications**
- **Progressive Cognitive Decline:** Loss of ability to function independently.
  - **Medical Complications:** Increased risk of infections, falls, and fractures.
  - **Behavioral and Psychological Symptoms:** Agitation, anxiety, depression, and sleep disturbances.
  - **End-Stage Complications:** Complete loss of ability to communicate, respond to surroundings, and control movement.

### Diagnosis

- a. Clinical Evaluation:** Based on medical history, symptoms, and cognitive tests to assess memory, language, and problem-solving abilities.
- b. Neuropsychological Testing:** Detailed assessment of cognitive function, including memory, attention, and executive function.
- c. Brain Imaging:** MRI or CT scans to detect changes in brain structure and rule out other causes of dementia.
- d. Biomarkers:** Tests for beta-amyloid and tau proteins in cerebrospinal fluid or imaging techniques (e.g., PET scans) to identify characteristic changes in the brain.

### Treatment

- a. Medications**
  - **Cholinesterase Inhibitors:** Donepezil, rivastigmine, galantamine to improve symptoms by increasing acetylcholine levels.
  - **Memantine:** NMDA receptor antagonist that regulates glutamate activity in the brain, improving symptoms and slowing progression.
- b. Symptomatic Treatment**
  - **Behavioral Interventions:** Cognitive stimulation therapy, reality orientation therapy to improve quality of life and reduce behavioral symptoms.
  - **Psychological Support:** Counseling for patients and caregivers to cope with the emotional and practical challenges of the disease.
- c. Experimental Therapies**
  - **Immunotherapy:** Targeting beta-amyloid or tau proteins to reduce their accumulation in the brain.
  - **Gene Therapy and Stem Cell Therapy:** Investigational approaches to repair or regenerate damaged neurons.

### Complications and Management

- a. Care Planning:** Long-term care planning to address safety, medical needs, and financial considerations.
- b. Support Services:** Accessing community resources, support groups, and respite care for caregivers.
- c. Advance Directives:** Documenting preferences for medical care and end-of-life decisions.

## Prevention

### a. Healthy Lifestyle Choices

- **Physical Activity:** Regular exercise to improve cardiovascular health and reduce risk of cognitive decline.
- **Healthy Diet:** Mediterranean-style diet rich in fruits, vegetables, whole grains, and lean proteins.
- **Mental Stimulation:** Engaging in intellectually stimulating activities, such as reading, puzzles, and social interactions.
- **Management of Chronic Conditions:** Controlling hypertension, diabetes, and other vascular risk factors that may contribute to dementia risk.
- **Social Engagement:** Maintaining social connections and participating in social activities.

### b. Research:

Participation in clinical trials to explore new treatments and preventive strategies.

## Alzheimer's Disease and the Gastrointestinal System:

While Alzheimer's disease primarily affects the brain, it can indirectly impact the gastrointestinal (GI) system through several mechanisms:

### GI Symptoms and Complications

- Dysphagia:** Difficulty swallowing, leading to aspiration pneumonia and malnutrition.
- Weight Loss:** Due to decreased appetite, forgetfulness of meals, or difficulty in chewing and swallowing.
- Constipation:** Common in Alzheimer's patients due to decreased physical activity, medications, and changes in routine.
- Incontinence:** Loss of bowel or bladder control, which can lead to embarrassment and social withdrawal.

### Managing GI Symptoms

- Dietary Modifications:** Soft or pureed foods, smaller and more frequent meals, and adequate hydration.
- Behavioral Interventions:** Assistance with eating, reminders to use the restroom, and establishing a regular toileting schedule.
- Medications:** Laxatives or stool softeners for constipation, anticholinergic medications for urinary incontinence.

### Preventive Measures for GI Issues

- Routine Monitoring:** Regular assessment of swallowing function, bowel habits, and nutritional status.
- Hydration and Nutrition:** Ensuring adequate fluid intake and a balanced diet rich in fiber and nutrients.
- Physical Activity:** Encouraging mobility and regular exercise to promote GI motility.