

Chapter-7

Haematological Disease-I

Mr. Abu Tahir

Assistant Professor
Rajiv Gandhi Institute of Pharmacy
Faculty of Pharmaceutical Science & Technology
AKS University Satna, (M.P.)

Abstract

Hematological diseases encompass a wide range of disorders affecting blood components, including red and white blood cells, platelets, and the coagulation system. Iron deficiency anemia (IDA) and megaloblastic anemia are two common types of hematological disorders. Iron deficiency anemia occurs when there is insufficient iron to produce hemoglobin, leading to reduced oxygen-carrying capacity of the blood. The pathophysiology involves depleted iron stores due to inadequate dietary intake, malabsorption, or chronic blood loss. Epidemiologically, IDA is the most common nutritional deficiency worldwide, particularly affecting women and children. Symptoms include fatigue, pallor, shortness of breath, and dizziness. Diagnosis is made through blood tests showing low hemoglobin, hematocrit, and serum ferritin levels. Treatment involves iron supplementation and addressing the underlying cause. Complications of untreated IDA can include heart problems and developmental delays in children. Prevention focuses on adequate dietary iron intake and early intervention in at-risk populations. Megaloblastic anemia, on the other hand, is caused by deficiencies in vitamin B12 or folic acid, leading to impaired DNA synthesis and the production of abnormally large red blood cells. The pathophysiology involves insufficient intake, malabsorption, or increased requirements of these vitamins. Epidemiologically, it is prevalent in populations with poor nutrition, the elderly, and individuals with malabsorption syndromes. Symptoms include fatigue, weakness, pallor, and neurological deficits such as numbness and cognitive changes in vitamin B12 deficiency. Diagnosis is confirmed through blood tests showing megaloblastic changes in red blood cells and low levels of vitamin B12 or folic acid. Treatment includes supplementation with the deficient vitamin and managing the underlying cause. Complications can include severe neurological damage in untreated vitamin B12 deficiency and increased risk of cardiovascular diseases. Prevention involves ensuring adequate dietary intake of vitamin B12 and folic acid, particularly in vulnerable groups such as vegetarians, pregnant women, and the elderly.

I. INTRODUCTION

Hematological Diseases

Hematological diseases affect the blood, bone marrow, and lymphatic systems. They can impact the production, function, and quality of blood cells and can be broadly classified into several categories:

1. **Anemias:** Conditions characterized by a deficiency in the number or quality of red blood cells (RBCs) or hemoglobin, leading to reduced oxygen delivery to tissues. Examples include:
 - a. **Iron Deficiency Anemia:** Caused by a lack of iron, which is crucial for hemoglobin production.
 - b. **Vitamin B12 and Folate Deficiency Anemia:** Resulting from insufficient vitamin B12 or folate, essential for RBC production.
 - c. **Aplastic Anemia:** A condition where the bone marrow fails to produce adequate amounts of blood cells.
2. **Leukemias:** Cancers of the blood and bone marrow characterized by the overproduction of abnormal white blood cells (WBCs). Types include:
 - a. **Acute Lymphoblastic Leukemia (ALL):** A rapid-growing leukemia affecting lymphoblasts.
 - b. **Acute Myeloid Leukemia (AML):** Affects myeloid cells and progresses quickly.
 - c. **Chronic Lymphocytic Leukemia (CLL):** A slow-growing leukemia affecting B lymphocytes.
 - d. **Chronic Myeloid Leukemia (CML):** Affects myeloid cells and is characterized by the presence of the Philadelphia chromosome.
3. **Lymphomas:** Cancers that originate in the lymphatic system. Major types are:
 - a. **Hodgkin Lymphoma:** Characterized by the presence of Reed-Sternberg cells.
 - b. **Non-Hodgkin Lymphoma:** A diverse group of blood cancers that includes various subtypes, such as diffuse large B-cell lymphoma (DLBCL) and follicular lymphoma.
4. **Myeloma:** Cancer of plasma cells in the bone marrow, leading to the production of abnormal antibodies. **Multiple Myeloma** is the most common form, characterized by bone pain, anemia, and kidney dysfunction.
5. **Bleeding Disorders:** Conditions that impair blood clotting, leading to excessive bleeding or bruising. Examples include:
 - a. **Hemophilia:** A genetic disorder resulting in insufficient clotting factors.
 - b. **Von Willebrand Disease:** A bleeding disorder caused by a deficiency or dysfunction of von Willebrand factor.
6. **Polycythemia:** Conditions characterized by an overproduction of red blood cells. **Polycythemia Vera** is a primary form associated with JAK2 mutation.

Endocrine System

The endocrine system is responsible for producing and regulating hormones that control various physiological processes. Key components include:

1. **Hypothalamus:** Regulates the pituitary gland and controls the release of hormones that influence growth, metabolism, and stress responses.
2. **Pituitary Gland:** Known as the "master gland," it secretes hormones that regulate other endocrine glands. It has two main lobes:
 - a. **Anterior Pituitary:** Produces hormones like growth hormone (GH), adrenocorticotropic hormone (ACTH), thyroid-stimulating hormone (TSH), luteinizing hormone (LH), and follicle-stimulating hormone (FSH).
 - b. **Posterior Pituitary:** Releases hormones such as oxytocin and antidiuretic hormone (ADH) produced by the hypothalamus.
3. **Thyroid Gland:** Produces thyroid hormones (thyroxine T4 and triiodothyronine T3) that regulate metabolism, energy levels, and growth.
4. **Parathyroid Glands:** Secrete parathyroid hormone (PTH), which regulates calcium and phosphate levels in the blood and bones.
5. **Adrenal Glands:** Located on top of the kidneys, they produce hormones such as cortisol (involved in stress response), aldosterone (regulates blood pressure), and adrenal androgens.
6. **Pancreas:** Functions as both an endocrine and exocrine gland. The endocrine component produces insulin and glucagon, which regulate blood sugar levels.
7. **Gonads**
 - a. **Ovaries:** Produce estrogen and progesterone, which regulate reproductive functions and secondary sexual characteristics.
 - b. **Testes:** Produce testosterone, which regulates sperm production and secondary sexual characteristics.
8. **Pineal Gland:** Produces melatonin, which regulates sleep-wake cycles.

Interactions between Hematological Diseases and the Endocrine System

1. **Endocrine Disorders and Blood Cell Production:** Conditions like hypothyroidism or hyperthyroidism can affect red and white blood cell production. For example, hypothyroidism can lead to anemia, while hyperthyroidism might cause increased red cell turnover.
2. **Adrenal Disorders:** Diseases such as Addison's disease (adrenal insufficiency) or Cushing's syndrome (excess cortisol) can impact various blood parameters, including blood pressure and glucose levels.
3. **Diabetes and Hematological Health:** Chronic diabetes can lead to various hematological complications, including an increased risk of infections due to impaired immune function.

II. HEMATOLOGICAL DISEASES

Iron deficiency

Introduction

Iron deficiency is one of the most common nutritional deficiencies worldwide, impacting various aspects of hematological and endocrine health. Iron is crucial for producing hemoglobin, which transports oxygen in the blood. Iron deficiency can lead to anemia and various systemic issues, impacting overall health and quality of life.

Pathophysiology

Iron deficiency impairs hemoglobin production due to insufficient iron availability. Hemoglobin is essential for oxygen transport in the blood, and low iron levels reduce its synthesis, leading to decreased oxygen delivery to tissues.

The pathophysiology of iron deficiency anemia (IDA) involves:

1. **Reduced Hemoglobin Production:** Low iron levels limit the synthesis of hemoglobin, leading to smaller and fewer red blood cells (RBCs).
2. **Impaired Oxygen Transport:** Anemia reduces the blood's oxygen-carrying capacity, affecting cellular metabolism and function.
3. **Increased Erythropoiesis:** The body tries to compensate by increasing red blood cell production, which may not be effective without adequate iron.

Epidemiology

Iron deficiency is prevalent globally and can affect any age group. The incidence varies by region and demographic factors:

1. **Infants and Children:** Rapid growth and increased iron requirements make them vulnerable. Iron deficiency is common in developing countries due to poor diet and inadequate breastfeeding.
2. **Pregnant Women:** Increased iron needs due to fetal development and blood volume expansion.
3. **Women of Reproductive Age:** Menstrual blood loss can contribute to iron deficiency.
4. **Elderly:** Chronic diseases, poor dietary intake, and gastrointestinal bleeding can lead to iron deficiency.
5. **Athletes:** Increased iron loss through sweat and high demand can result in deficiency.

Symptoms and Complications

Symptoms of Iron Deficiency

1. **Fatigue and Weakness:** Due to reduced oxygen delivery to tissues.
2. **Paleness:** Due to low hemoglobin levels.
3. **Shortness of Breath and Dizziness:** Especially during physical activity.
4. **Cold Hands and Feet:** Due to poor circulation.
5. **Brittle Nails and Hair Loss:** Reflecting poor cellular function.
6. **Pica:** Craving for non-nutritive substances like ice, dirt, or starch.

Complications

1. **Severe Anemia:** Can lead to cardiovascular complications, including heart failure.
2. **Pregnancy Complications:** Increased risk of preterm birth, low birth weight, and postpartum depression.
3. **Impaired Cognitive and Developmental Function:** In children, iron deficiency can affect cognitive development and academic performance.

Diagnosis

Diagnostic Tests

1. **Complete Blood Count (CBC):** To evaluate red blood cell count, hemoglobin levels, and hematocrit.

2. **Serum Ferritin:** Measures the amount of stored iron in the body. Low levels indicate iron deficiency.
3. **Serum Iron:** Measures the amount of circulating iron. Low levels can indicate deficiency.
4. **Total Iron-Binding Capacity (TIBC):** Elevated in iron deficiency; measures the blood's ability to bind iron.
5. **Transferrin Saturation:** Low percentage indicates iron deficiency.

Additional Tests

1. **Peripheral Blood Smear:** To observe red blood cell morphology (e.g., microcytic, hypochromic cells).
2. **Bone Marrow Aspiration:** In some cases, to evaluate iron stores if the diagnosis is unclear.

Treatment

Iron Supplementation

1. **Oral Iron Supplements:** Ferrous sulfate, ferrous gluconate, or ferrous fumarate are commonly used. Usually prescribed at a dose of 100-200 mg of elemental iron per day.
2. **Intravenous Iron:** For patients who cannot tolerate oral iron or have severe deficiency. Examples include iron sucrose, ferric gluconate, and iron dextran.

Dietary Changes

1. **Iron-Rich Foods:** Incorporate sources like red meat, poultry, fish, lentils, beans, and fortified cereals.
2. **Vitamin C:** Enhances iron absorption. Include citrus fruits, tomatoes, and peppers in the diet.

Addressing Underlying Causes

1. **Gastrointestinal Disorders:** Treat conditions like peptic ulcers or celiac disease that may contribute to iron deficiency.
2. **Menstrual Issues:** Address heavy menstrual bleeding with appropriate treatments.

Prevention

General Recommendations

1. **Balanced Diet:** Ensure adequate intake of iron-rich foods, particularly in high-risk populations.
2. **Iron Supplementation:** For at-risk groups such as pregnant women, infants, and individuals with chronic conditions.
3. **Regular Screening:** For populations at risk (e.g., pregnant women, children, and the elderly) to detect iron deficiency early.
4. **Education and Awareness:** Promote dietary diversity and awareness of iron-rich foods and proper supplement use.

Specific Interventions

1. **Pregnant Women:** Routine iron supplementation during pregnancy.
2. **Infants and Children:** Supplementation and dietary guidance to prevent deficiency.

Effective management of iron deficiency involves a combination of appropriate treatment, addressing underlying causes, and preventive measures to ensure optimal health outcomes.

Megaloblastic anemia (Vit B₁₂ and folic acid):

Introduction

Megaloblastic anemia is a type of anemia characterized by the presence of abnormally large and immature red blood cells (megaloblasts) in the bone marrow and peripheral blood. It is primarily caused by deficiencies in vitamin B12 (cobalamin) or folic acid (vitamin B9), both of which are essential for proper DNA synthesis and red blood cell maturation.

Pathophysiology

1. Vitamin B12 Deficiency

- a. **Impaired DNA Synthesis:** Vitamin B12 is essential for DNA synthesis and cell division. Its deficiency leads to impaired DNA replication, resulting in the production of large, immature red blood cells.
- b. **Neurological Effects:** Vitamin B12 is also crucial for maintaining myelin sheaths in nerves. Deficiency can lead to neurological symptoms due to demyelination.

2. Folic Acid Deficiency

- a. **Impaired Cell Division:** Folic acid is vital for DNA synthesis and cell division. Deficiency leads to the production of large, dysfunctional red blood cells.
- b. **Effect on Rapidly Dividing Cells:** Folic acid deficiency affects rapidly dividing cells, including those in the bone marrow, leading to megaloblastic anemia.

In both deficiencies, the bone marrow produces large, immature red blood cells that are released into the bloodstream, where they cannot function effectively.

Epidemiology

1. Vitamin B12 Deficiency

- a. More common in older adults due to decreased absorption in the gastrointestinal tract.
- b. Higher prevalence in individuals with gastrointestinal disorders (e.g., Crohn's disease, pernicious anemia) or those undergoing gastric surgery.
- c. Vegetarians and vegans are at higher risk due to the absence of vitamin B12 in plant-based diets.

2. Folic Acid Deficiency

- a. Common in pregnant women due to increased demand.
- b. Higher incidence in individuals with poor dietary intake, alcoholism, or malabsorption issues (e.g., celiac disease).
- c. Certain medications (e.g., methotrexate, antiepileptics) can interfere with folic acid metabolism.

Symptoms and Complications

Symptoms

1. **General Anemia Symptoms:** Fatigue, weakness, pallor, and shortness of breath.
2. **Vitamin B12 Deficiency Symptoms**

- a. **Neurological Symptoms:** Numbness or tingling in extremities, difficulty walking, and memory loss.
- b. **Glossitis:** Inflammation of the tongue.
- c. **Mood Changes:** Depression or irritability.

3. Folic Acid Deficiency Symptoms

- a. **Glossitis and Mouth Ulcers:** Inflammation and sores in the mouth.
- b. **Diarrhea:** Gastrointestinal discomfort.
- c. **Pregnancy Complications:** Neural tube defects in the fetus if deficiency occurs during pregnancy.

Complications

1. **Neurological Damage:** Irreversible neurological damage if vitamin B12 deficiency is not treated promptly.
2. **Pregnancy Complications:** Increased risk of birth defects, including neural tube defects in the baby.
3. **Impaired Immune Function:** Both deficiencies can impact immune system function, leading to increased susceptibility to infections.

Diagnosis

Diagnostic Tests:

1. **Complete Blood Count (CBC):** Shows macrocytic anemia with high mean corpuscular volume (MCV).
2. **Peripheral Blood Smear:** Reveals megaloblasts and hypersegmented neutrophils.
3. **Serum Vitamin B12 Levels:** Low levels indicate deficiency.
4. **Serum Folate Levels:** Low levels indicate deficiency.
5. **Methylmalonic Acid (MMA):** Elevated in vitamin B12 deficiency.
6. **Homocysteine Levels:** Elevated in both vitamin B12 and folic acid deficiencies, but more specific to vitamin B12 deficiency.

Additional Tests

1. **Intrinsic Factor Antibodies:** To diagnose pernicious anemia (a common cause of vitamin B12 deficiency).
2. **Bone Marrow Biopsy:** In some cases, to assess for megaloblasts and rule out other causes of anemia.

Treatment

Vitamin B12 Deficiency

1. **Oral Supplements:** Typically prescribed at 1,000-2,000 µg daily if absorption is adequate.
2. **Intramuscular Injections:** 1,000 µg of vitamin B12 administered intramuscularly weekly or monthly, depending on the severity and cause of the deficiency.

Folic Acid Deficiency

1. **Oral Supplements:** Typically prescribed at 1 mg daily.
2. **Dietary Changes:** Increase intake of folate-rich foods such as leafy greens, legumes, and fortified cereals.

Addressing Underlying Causes

1. **Vitamin B12 Malabsorption:** Treat underlying conditions like pernicious anemia or gastrointestinal issues.
2. **Folic Acid Absorption Issues:** Address any underlying causes, such as malabsorption syndromes or drug interactions.

Complications

1. **Neurological Damage:** Especially in vitamin B12 deficiency, if treatment is delayed.
2. **Persistent Anemia:** If the underlying cause is not addressed, anemia may persist or recur.
3. **Pregnancy Complications:** Inadequate folic acid intake during pregnancy can lead to serious fetal developmental issues.

Prevention

General Recommendations

1. **Balanced Diet:** Ensure adequate intake of vitamin B12 and folic acid through a varied diet.
2. **Supplements:** For individuals at risk, such as the elderly, pregnant women, and those with malabsorption issues.

Specific Interventions

1. **Pregnant Women:** Routine supplementation of folic acid to prevent neural tube defects.
2. **Vegetarians and Vegans:** Regular vitamin B12 supplementation or fortified foods.
3. **Elderly Individuals:** Regular screening and supplementation if needed.