

PIONEERING CROP IMPROVEMENT THROUGH EXPLORATION OF PROTEOMICS FOR NEXT GENERATION AGRICULTURE

Abstract

The field of agriculture is facing significant challenges due to a growing global population, climate change, and diminishing natural resources. To ensure sustainable food production, there is an urgent need for innovative approaches to improve crop yields, nutritional content, and stress tolerance. In recent years, proteomics has emerged as a powerful tool for understanding the complex network of proteins involved in plant growth, development, and response to environmental stimuli. This chapter discusses various proteomic techniques, including gel-based and gel-free approaches, mass spectrometry-based methods, and protein-protein interaction studies, highlighting their strengths and limitations. Researchers have gained insights into the mechanisms underlying nutrient uptake, disease suppression, and enhanced growth by studying the plant proteome. These findings can be exploited to enhance the nutritional value, flavor, and shelf life of crops, addressing the increasing demands of consumers for healthier and more sustainable food options. While proteomics has made significant strides in crop improvement, challenges still exist, such as the complexity of protein extraction, the dynamic nature of the proteome, and the need for robust data analysis pipelines. Overcoming these hurdles will require interdisciplinary collaborations and advancements in technology and bioinformatics. This chapter aims to provide an overview of the potential applications of proteomics in next-generation agriculture and its role in crop improvement.

Keywords: Pioneering Crop, proteomic techniques, agriculture

Authors

Dr. Arpita Mahobia
Biotechnology Centre
Department of Agricultural Botany
Dr. Panjabrao Deshmukh Krishi
Vidyapeeth
Akola, Maharashtra, India.
35arpita89@gmail.com

Sagar L. Zanjali
Biotechnology Centre
Department of Agricultural Botany
Dr. Panjabrao Deshmukh Krishi
Vidyapeeth
Akola, Maharashtra, India.

Dr. Sanjay B. Sakhare
Biotechnology Centre
Department of Agricultural Botany
Dr. Panjabrao Deshmukh Krishi
Vidyapeeth
Akola, Maharashtra, India.

Dr. Rameshwar B. Ghorade
Biotechnology Centre
Department of Agricultural Botany
Dr. Panjabrao Deshmukh Krishi
Vidyapeeth
Akola, Maharashtra, India.

Dr. Shyamsunder S. Mane
Department of Plant Pathology
Dr. Panjabrao Deshmukh Krishi
Vidyapeeth
Akola, Maharashtra, India.

I. INTRODUCTION: PROTEOMICS AND CROP IMPROVEMENT

According to an estimate by World Hunger Fact and Statistics, there are approximately 783 million people on the globe who live in a state of hunger (www.actionagainsthunger.org). Furthermore, the global human population reached 8.0 billion in mid-November 2022, adding 2 billion people since 1998. The world's population is anticipated to increase by nearly 2 billion persons in the next 30 years, from the current 8 billion to 9.7 billion in 2050 and could crown at nearly 10.4 billion in the mid-2080s. By combining various factors and bolstering the plant breeding tools for crop advances, we can greatly enhance food production and supply in an endeavor to erase that unsightly patch of hunger from the lovely face of humanity [1]. A key challenge for plant breeders working on crop improvement projects is the small gene pool of domesticated crop species. A critical step in improving essential crop features is the identification of possibly beneficial genes from the animal and plant kingdoms. These genes are typically discovered through study in molecular biology, including genomics and proteomics. These recently found genes may benefit human society when they are introduced into a desired crop species and used in breeding operations. Additionally, climate change poses significant challenges, including increased occurrence of drought, salinity, heat, and pest infestations, which further threaten agricultural productivity. Addressing these challenges requires innovative approaches that go beyond traditional breeding methods. However, with the advent of molecular biology and genomics, significant progress has been made in understanding the genetic basis of plant traits. Nevertheless, the information provided by genomics alone is insufficient to decipher the complex regulatory networks and molecular mechanisms underlying plant growth, development, and response to environmental stresses

Over the recent decade, the use of proteomics for crop plant analysis has grown dramatically. Although proteomic techniques are widely utilized in plant laboratories around the world and are effective research tools, there is still much opportunity for advancement. The fraction of the plant proteome that can be detected with current methodologies, in particular, is significantly smaller than that of other "Omics" techniques, and so does not provide a complete picture of the cellular proteins. Proteomics, as a discipline, has revolutionized our understanding of complex biological systems by focusing on the comprehensive study of proteins, their structure, function, and interactions within cells and organisms. With the ever-growing global population, climate change, and limited availability of arable land and resources, there is a pressing need to develop sustainable and efficient agricultural practices. Crop improvement, aimed at enhancing crop productivity, quality, and stress tolerance, is essential to meet the increasing demands for food, feed, and bioenergy. By employing proteomic techniques, researchers aim to identify key regulatory proteins, elucidate signaling pathways, and discover novel biomarkers that can contribute to crop improvement and sustainable agriculture. Traditionally, crop improvement strategies relied on classical breeding techniques, which often took several years to develop new varieties with desired traits.

This is where proteomics steps in. Proteomics complements genomics by providing insights into the dynamic and functional aspects of the plant proteome, which comprises the entire set of proteins expressed in a specific tissue, developmental stage, or under different environmental conditions. By studying the proteome, proteomics enables researchers to identify and quantify proteins, understand their interactions and modifications, and unravel

their roles in various cellular processes. In the context of crop improvement, proteomics offers several distinct advantages. Firstly, it provides a direct measurement of the functional molecules responsible for plant phenotypes. Proteins are the primary effectors of biochemical reactions and cellular processes, making them crucial targets for crop enhancement. Secondly, proteomics can identify proteins that are specifically induced or suppressed under stress conditions, facilitating the discovery of stress-responsive biomarkers. These biomarkers can be utilized for early stress detection, monitoring crop health, and guiding breeding programs for stress tolerance. Furthermore, proteomics aids in unraveling the complexities of post-translational modifications (PTMs), which play crucial roles in protein function and regulation. PTMs, such as phosphorylation, acetylation, and glycosylation, can modulate protein activity, stability, and subcellular localization [2]. Understanding these modifications is essential for deciphering plant signaling pathways and regulatory networks.

In recent years, significant progress has been made in proteomic techniques, including advancements in protein separation, identification, and quantification methodologies. Mass spectrometry-based approaches, such as liquid chromatography-mass spectrometry (LC-MS/MS), have revolutionized the field by enabling high-throughput and comprehensive analysis of complex protein mixtures. Additionally, advancements in bioinformatics tools and data analysis algorithms have improved the interpretation and integration of large-scale proteomic datasets. The integration of proteomics with other omics technologies, such as genomics, transcriptomics, and metabolomics, is also crucial for a systems biology approach to crop improvement. Integrative analysis of multi-omics data allows for a comprehensive understanding of the molecular interactions and regulatory networks that govern plant growth and stress responses.

In conclusion, proteomics offers a powerful approach for studying the plant proteome and its implications in crop improvement. By providing insights into protein expression, function, and interactions, proteomics can guide the development of genetically improved crop varieties with enhanced yield, quality, and stress tolerance. The integration of proteomics with other omics technologies and the development of standardized protocols and bioinformatics tools will further advance our understanding of plant biology and contribute to the sustainable future of agriculture.

II. ROLE OF PROTEOMICS IN AGRICULTURE

Proteomics has emerged as a powerful tool in the field of agriculture, offering valuable insights into the complex molecular processes underlying plant growth, development, and response to environmental stresses. By comprehensively studying the proteome, which represents the entire complement of proteins in an organism, proteomics enables researchers to identify key regulatory proteins, elucidate signaling pathways, and discover biomarkers associated with stress tolerance and nutritional traits. This knowledge is crucial for developing innovative strategies to enhance crop productivity, quality, and sustainability.

- 1. Understanding Plant Stress Responses:** Proteomics plays a vital role in unraveling the molecular mechanisms by which plants respond to abiotic and biotic stresses, such as drought, salinity, heat, cold, pathogens, and pests. Proteomic studies have identified stress-responsive proteins, including molecular chaperones, antioxidant enzymes, and

defense-related proteins, providing insights into the adaptive strategies employed by plants to withstand adverse conditions [3]. For example, a study on wheat under drought stress revealed differentially expressed proteins involved in stress signaling, osmotic regulation, and reactive oxygen species scavenging, contributing to a better understanding of drought tolerance mechanisms [4].

- 2. Unraveling Plant Development and Physiology:** Proteomics facilitates the investigation of protein expression patterns during various stages of plant development, differentiation of tissues, and response to hormonal signals. It helps in identifying key regulatory proteins involved in growth, flowering, fruit development, and senescence [4]. For instance, proteomic analysis of tomato fruit ripening identified proteins associated with ethylene signaling, cell wall remodeling, and pigment synthesis, shedding light on the molecular events underlying fruit ripening [5].
- 3. Enhancing Nutritional Traits:** Proteomic approaches contribute to improving the nutritional quality of crops by identifying proteins associated with essential nutrients and anti-nutritional factors. By characterizing the proteome of biofortified crops, researchers can identify proteins involved in nutrient accumulation and bioavailability, facilitating the development of crops with enhanced nutritional content. For example, proteomics has been employed to investigate iron and zinc accumulation in staple crops like rice and wheat, providing insights into the molecular basis of nutrient uptake, transport, and storage [6, 7]
- 4. Uncovering Protein-Protein Interactions and Signaling Networks:** Proteomics enables the exploration of protein-protein interactions and signaling networks in plants, uncovering the complex regulatory mechanisms governing various physiological processes. Techniques such as yeast two-hybrid and affinity purification coupled with mass spectrometry have been employed to identify protein interaction partners and map signaling pathways [7]. These studies have provided valuable insights into protein complexes involved in plant hormone signaling, stress responses, and developmental processes.
- 5. Integrating Proteomics with Other Omics Technologies:** The integration of proteomics with other omics technologies, such as genomics, transcriptomics, and metabolomics, allows for a systems biology approach to understand plant biology comprehensively. By combining multi-omics datasets, researchers can unravel the intricate interactions between genes, transcripts, proteins, and metabolites, providing a holistic view of plant molecular networks [8, 9]. This integrative approach aids in identifying novel targets for crop improvement and understanding the functional relevance of different molecular components.




Figure 1: Role of Proteomics in agriculture

III. PROTEOMIC TECHNIQUES FOR CROP ANALYSIS AND IMPROVEMENT

Proteomic techniques have revolutionized the field of crop science, providing valuable insights into the complex molecular processes underlying plant growth, development, and responses to environmental stresses. By comprehensively studying the proteome - the entire set of proteins expressed by an organism or tissue - researchers can unravel the functional roles of proteins, identify key regulatory factors, and discover potential targets for crop improvement. In this article, we explore some of the proteomic techniques used in crop analysis and improvement. Several distinct types of proteomics approaches have been developed, each with its specific focus and applications. Here are some commonly employed types of proteomics:

- 1. Expression Proteomics:** Expression proteomics aims to identify and quantify changes in protein expression levels under different conditions or in response to specific stimuli. This type of proteomics involves techniques such as two-dimensional gel electrophoresis (2-DE) and shotgun proteomics, which utilize mass spectrometry-based methods for protein identification and quantification. Expression proteomics provides insights into differential protein expression patterns, enabling the identification of proteins associated with specific biological processes or disease states [10].
- 2. Structural Proteomics:** Structural proteomics focuses on determining the three-dimensional structures of proteins, which are crucial for understanding their functions and interactions. Structural proteomics employs techniques such as X-ray crystallography, nuclear magnetic resonance (NMR) spectroscopy, and cryo-electron microscopy (cryo-EM). These methods allow researchers to visualize the atomic-level details of protein structures, aiding in the identification of protein binding sites, enzymatic mechanisms, and drug target interactions [10].

- 3. Functional Proteomics:** Functional proteomics aims to elucidate the functional roles of proteins within a cellular or organism context. It involves studying protein-protein interactions, protein modifications, and protein localization. Techniques commonly employed in functional proteomics include yeast two-hybrid assays, co-immunoprecipitation, protein microarrays, and mass spectrometry-based analysis of post-translational modifications (PTMs) such as phosphorylation, acetylation, and glycosylation. Functional proteomics provides insights into protein networks, cellular signaling pathways, and regulatory mechanisms [10].
- 4. Interactomics:** Interactomics focuses on mapping protein-protein interactions within a biological system. This type of proteomics utilizes various techniques, including affinity purification coupled with mass spectrometry (AP-MS), co-immunoprecipitation, and protein fragment complementation assays. Interactomics enables the identification of protein complexes, signaling cascades, and dynamic interactions between proteins, shedding light on cellular processes and pathways [11]. Affinity purification coupled with mass spectrometry (AP-MS) is commonly used to identify interacting partners of a specific protein. This technique involves the isolation of a protein of interest and the identification of associated proteins using mass spectrometry. Other methods, such as yeast two-hybrid and co-immunoprecipitation, also enable the identification and characterization of protein-protein interactions. Understanding PPIs in crops provides insights into signaling pathways, protein complex formation, and regulatory networks.
- 5. Metaproteomics:** Metaproteomics involves the study of the collective proteome of a microbial community present in an environmental sample. This approach combines techniques from proteomics and metagenomics to analyze complex microbial ecosystems. Metaproteomics allows for the identification and characterization of microbial species, their functional activities, and the interactions between community members. It provides valuable insights into microbial ecology, biogeochemical cycles, and host-microbe interactions [12].




Figure 2: Proteomics branches

IV. PROTEOMICS ANALYSIS TECHNIQUES FOR CROP IMPROVEMENT

Protein expression analysis broadly includes following areas *viz.*, (a) Conventional Techniques, (b) Advanced Techniques, (c) Quantitative Techniques and (d) Bioinformatics Analysis.

1. Conventional Techniques: Conventional methods for protein purification includes mainly chromatography-based techniques, *viz.*, Ion Exchange Chromatography (IEC), Affinity Chromatography, Size Exclusion Chromatography (SEC) etc. The Western blotting technique is used to detect protein N terminal amino acids. Such methodologies are frequently confined to studying a few distinct proteins and are unable to establish the level of protein expression.

- **Ion Exchange Chromatography (IEC):** The IEC approach is used to separate various proteins present in a mixture based on the ionic charges of the relevant proteins' surfaces. Proteins are polymers of zwitter ionic amino acid sequences; some amino acids have an additional negative charge (anionic), while others are cationic. A protein's total charge is reflected by its physiological pH in equilibrium condition. IEC first differentiates the protein based on its charge (anionic and cationic), followed by a comparative charge intensity.
- **Size Exclusion Chromatography (SEC):** Proteins are passed through a stationary phase of porous beads with various pore diameters in SEC. Proteins are classified here according to their molecular sizes. SEC can differentiate proteins in a variety of physiological conditions such as the presence of ions, detergents, co-factors, or at different temperatures. It can separate proteins with low molecular weight and is a significant approach for the purification of non-covalent multimetric protein complexes under biological [13].
- **Affinity chromatography:** A stationary phase in affinity chromatography is a matrix to which antibodies are linked. Proteins operate as antigens, and proteins are separated depending on antigen-antibody interactions. Identification of microbial enzymes responsible for pathogenicity is one of its significant applications[13]. Grice used metal chelate affinity chromatography to isolate homo and hetero dimer components of HIV-I reverse transcriptase enzyme [14].
- **Western blotting technique:** The Western blotting technique detects low abundance proteins using electrophoresis. Proteins are transferred to a nitrocellulose membrane that is positively charged. Proteins with negative charges bind to the membrane. Primary antibodies bind to the target protein, leaving uncharged regions. Milk protein is utilized as a mask to prevent non-specific binding. The system has been cultured with X-phos. Following the elimination of phosphate from X-phos, a secondary antibody with alkaline phosphatase tag is linked to the tail of the primary antibody. As a result, blue precipitate is formed, indicating that the specific protein has been isolated. Western blotting techniques are used to detect antigens from various microorganisms in order to diagnose various infectious illnesses. This approach was used to identify ten key rice proteins. In rice, EF1-₂ and HSPs were substantially

expressed proteins [15]. Kollerova et al [16] extracted PPV virus capsid proteins from *Nicotianabenthamiana* samples.

2. Advanced Techniques: Protein microarrays or chips are used to analyze protein expression. Protein microarrays, on the other hand, are insufficient for investigating the involvement of the entire genome. On the other hand, the Edman degradation method was developed for identifying the amino-acid sequences of a specific protein molecule.

- **Protein Microarrays:** Protein Microarrays are utilized for high throughput sample amount protein identification. It is classified into three types: analytical microarray techniques, functional proteins, and reverse phase microarray. Proteins are detected in Antibody microarray via direct marking after antibody capture. This method is commonly used to calculate protein expression levels and binding affinities. Analytical and experiment-based methodologies have been developed to identify cellular signaling pathways and characterize plant kinases utilizing protein microarrays [17].

➤ **Functional Protein Microarray** A functional protein microarray is constructed using purified protein, allowing the analysis of various types of interactions such as DNA with protein, RNA with protein, protein-protein, ligand and protein, lipid with protein, enzyme-substrate interaction, and so on. It has the potential to change the roles of thousands of proteins. The microarray technique was first used to investigate the degree of substrate specificity of yeast enzyme kinases. The relationship of protein-protein interactions in *Arabidopsis thaliana* was examined, and CML and CaM proteins were found [18].

➤ **Reverse Phase Protein Microarray** In reverse phase protein microarray analysis, cell lysates from various cell stages are put on nitrocellulose slides and tested with target protein antibodies. Antibodies are detected using a combination of fluorescent, chemiluminescent, and colorimetric chemicals. For protein quantitation, reference peptides are written on slides [19]. These microarrays calculate abnormal proteins that are indicators of a certain ailment.

- **SDS-PAGE (Sodium Dodecyl Sulfate-Polyacrylamide Gel Electrophoresis)** SDS-PAGE is used to separate proteins based on their molecular sizes, which are determined using an approximation of the associated molecular weights [21]. An electric field is used to move proteins through a liquid with a pH different from their individual isoelectric point. Because of the m/z ratio of their mass and ionic charge, various proteins in the mixture move at different rates. Sodium dodecyl sulfate denatures the proteins, causing them to separate completely according to molecular weight. This method looked at outer membrane proteins from *Escherichia coli* that were not capable of producing K1 antigen [22]. The storage proteins of chickpea (*Cicerarietinum*) leaves and seeds were profiled under drought stress and control circumstances. Using this method, storage proteins found in the seeds of *Brassica* species are identified in order to study the molecular divergence of diverse genotypes.

- **Two-Dimensional Gel Electrophoresis (2-DE)** is a traditional proteomic technique for protein separation and visualization based on isoelectric point (pI) and molecular

weight. Proteins are separated on a gel by their charge in the first dimension (isoelectric focusing) and size in the second dimension (sodium dodecyl sulfate-polyacrylamide gel electrophoresis) in 2-DE [23]. This method detects differentially expressed proteins comparing samples, revealing changes in protein abundance linked with diverse circumstances or treatments. It can distinguish between 5,000 different proteins at the same time. In the first step, proteins are passed through the gel matrix and sorted based on their charges. Proteins are differentiated in the second dimension based on their masses. Cell membrane proteins from *Listeria monocytogenes* and *Listeria innocua* were examined using 2D-PAGE to visualize isolated proteins, and 30 different types of proteins were isolated [24]. This technique is used to compare the virulence factors and exotoxins produced by two enterotoxigenic *Staphylococcus aureus* strains derived from food [25].

- **Mass Spectrometry (MS)-Based Proteomics** is an effective method for identifying and quantifying proteins. It entails ionizing proteins or peptides and separating them according to their mass-to-charge ratio. Mass spectrometry, when combined with different separation techniques such as liquid chromatography (LC) or capillary electrophoresis (CE), allows for the detection, characterization, and quantification of proteins in complicated mixtures. Shotgun proteomics and targeted proteomics, for example, provide crucial insights into protein expression, post-translational changes, and protein-protein interactions in crops [26].
 - **Edman Sequencing** Edman sequencing was developed by Pehr Edman (1950) to discover N terminal amino acid residues. It is critical in the creation and quality assessment of proteins used in the manufacture of biomedicines. To discover the roles of rice leaf sheath proteins, Mass spectrometry and Edman sequencing approaches were used [27].
- 3. Quantitative Techniques:** Stable Isotope Labeling with Amino Acids in Cell Culture (SILAC), Isotope-Coded Affinity Tag (ICAT), and isobaric Tag for Relative and Absolute Quantitation (iTRAQ) are techniques used in quantitative proteomics research. ICAT is used to precisely quantify and identify the sequence compositions of individual proteins in a mixture by employing labelled peptides, isotopic linker sequences, and reactive groups. To mark metabolites in whole proteomes in a cell system, SILAC is employed in quantitative proteomics studies. iTRAQ is a multiplex technology used for protein tagging and quantification. It tags proteins with isobaric chemicals (8-plex and 4-plex) for relative and absolute quantification [28]. The method involves labeling protein groups at the N-terminus and side chains with amine, which is then separated by liquid chromatography and evaluated by MS. Finding gene control is critical to understanding the disorder's mechanism.
- 4. Bioinformatics Analysis:** Bioinformatics is required for proteomics research. With the introduction of high-throughput next -generation sequencing (NGS) technology, a large amount of sequence data has become publicly available. Bioinformatics advances have made it easier to examine this data with great sensitivity and precision in order to identify, quantify, and assess protein expression levels. The advancement of bioinformatics technology has aided drug design research. Molecular docking is an important step before

this drug designing [30]. It is carried out to find the best shape of ligand molecules at protein binding sites so that total energy is minimized. There are mainly two types of molecular docking, viz., rigid docking and flexible docking. Several bioinformatics tools and databases are developed to facilitate different types of proteomics analysis are given below in (Table 1).

Table 1:- Software of Bioinformatics for Proteomics study

S.No.	Function	Software
1	Protein primary sequence database	SwissProt, UniProt
2	Protein primary structure database	PDB
3	Protein secondary structure database	SCOP, CATH
4	Similarity search	BLASTP
5	Pattern hit initiated BLAST	PHI-BLAST
6	Position specific iterated BLAST	PSI-BLAST
7	Protein domain server	InterproScan
8	Motif based alignment server	MEME
9	Homology modelling tool, BLAST search, T-Coffee alignment, and MODELLER construction	Biskit
10	Homology modelling tool, Automated webserver	ESyPred3D
11	Homology modelling tool, Downloadable program	FoldX
12	Homology modelling tool, Standalone program mainly in Fortran and Python	MODELLER
13	Homology modelling tool, Standalone program mainly in Fortran and Perl	CONFOLD
14	Homology modelling tool, Proprietary platform, supported on Windows, Linux and Mac	MOE (Molecular Operating Environment)
15	Homology modelling tool, Webserver	ROBETTA
16	Homology modelling tool, Protein tertiary structure predictions	BHAGEERATH-H
17	Homology modelling tool, Automated webserver (based on ProModII)	SWISS-MODEL
18	Homology modelling tool, Graphical interface or text mode (clusters)	Yasara
19	Homology modelling tool, Automated webserver	AWSEM-Suite
20	Threading, On-line server for protein modeling	I-TASSER
21	Homology modelling tool, Threading, Automated webserver and some downloadable programs	IntFOLD
22	Homology modelling tool, Threading, Webserver with job manager, automatically updated fold library, genome searching and other facilities	Phyre and Phyre2
23	Homology modelling tool, Threading, Automated webserver and Downloadable program	RaptorX
24	Homology modelling tool, Threading, Interactive webserver with help facility	HHpred

25	Threading	3D-PSSM
26	Molecular docking tool, Open source	Autodock, FlexAID, rDOCK SEED
27	Molecular docking tool, Freeware	DOCK, LeDOCK
28	Molecular docking tool, Commercial	Glide, Molecular Operating Environment (MOE)

V. PROTEOMIC TECHNIQUES OFFER NEW TOOLS FOR PLANT BIOTECHNOLOGY

Proteomics, a potent tool for deciphering the intricate molecular processes within plants, is increasingly being used by researchers in their hunt for sustainable and high-yielding crop types. The capacity to discover essential regulatory proteins and interpret their involvement in various physiological processes is one of the fundamental benefits of studying plant proteomes. Researchers can identify and quantify proteins present in different tissues, organs, and developmental stages of plants using techniques such as mass spectrometry-based proteomics and protein separation methods [31]. This allows them to detect variations in protein expression patterns under varied growth conditions or stress settings, providing important insights into plant adaptation strategies. Knowledge of important proteins that play critical functions in the normal growth and development of a plant is critical for propelling agricultural plant biotechnology innovation [32]. These proteins regulate physiological and biochemical pathways to maintain cellular homeostasis in a particular environment. Continuous identification of genes and proteins responsible for stress tolerance and disease resistance is required to boost agricultural output.

Advances in MS-based proteomics platforms have been dubbed "New Genomics" since MS has become a key tool for investigating PTMs to proteins and protein interactions. Proteomics has a distinct benefit over other "Omics" techniques in that it may identify post-translational modifications (PTMs), which is required to establish the functional impact of protein modification on agricultural plant productivity. Scientists can identify protein interaction partners and map out intricate signaling cascades using methods such as yeast two-hybrid tests and affinity purification coupled with mass spectrometry [33]. Plant proteomics is also important in understanding protein post-translational modifications (PTMs) such as phosphorylation, glycosylation, and acetylation. Protein activity, stability, and subcellular localization are all influenced by PTMs, which are important regulatory mechanisms. Researchers may identify and quantify PTMs using modern proteomic techniques, providing insights into the dynamic changes that occur in protein function under diverse situations [34]. Proteomic investigations, for example, have contributed in the identification of stress-responsive proteins linked to drought, salinity, heat, and disease resistance, suggesting prospective targets for genetic improvement and breeding programs (Kazan, 2015). Proteomics is opening up new methods for crop improvement studies via the signaling, regulatory, and metabolic networks underlying plant phenotypes. Similarly, proteomics has been useful in determining the molecular basis of nutritional features, which has led to the production of crops with increased nutritional content, such as biofortified

cultivars. Jacoby et al. presented a series of studies in which they attempted to map the proteomes of several plant tissues from rice and Arabidopsis. 2-DE was used to separate the proteins from the various tissues [36]. In keeping with this consensus, various plant proteomics studies focusing on specific subcellular proteomes or protein complexes such as the plasma membrane, roots, mitochondria, and chloroplasts have recently been reported. Furthermore, a few intriguing studies on the symbiotic relationship between legume roots and nitrogen-fixing bacteria have emerged.

1. Proteome Analysis of a Biotic Stress Response in Crop Plants

- **Leaf Photosynthesis and Senescence Proteomics to Improve Crop Productivity:** The primary source of plant biomass impacting prospective crop output is leaf photosynthesis. Urban, Milan Oldřich investigated the response of leaf proteome to long-term drought (28 days) was studied in cultivars (cvs): Californium (C), Cadeli (D), Navajo (N), and Viking (V). Analysis of twenty-four 2-D DIGE gels revealed 134 spots quantitatively changed at least 2-fold; from these, 79 proteins were significantly identified by MALDI-TOF/TOF [37]. According to Roberts et al., chloroplast carries up to 75% of leaf nitrogen in the form of Rubisco enzyme constituents in the stroma and a complex of photosystem II in the thylakoid membrane. Advances in organelle proteomic studies combined with large-scale genomic approaches, as well as the identification of enzymes with proteolytic activity, have focused on the complexity of chloroplast proteolytic machinery around leaf senescence and investigated various categories of senescence-associated proteases with distinct physiological roles based on their expression profiles as senescence progresses [38]. Subcellular proteomic analyses of senescence, the process of photosynthesis and stress-responding processes in rice leaves have discovered glycolytic enzymes involved in sucrose production, which are of particular importance in terms of crop output [39].
- **Proteomics of Root-to-Leaf Signaling Pathways During Stress in Xylem and Phloem:** Increasing crop yields also requires the leaves to get an adequate amount of nutrients from the network of roots via the xylem vessels. Several kinds of xylem sap containing proteins that are involved in cell wall development and repair process [40], senescence of leaves [41], abiotic stress responses [42], abiotic stress defense mechanisms [43], and intercellular and intracellular communication [44]. Additional research on the protein and metabolism product composition of xylem sap and apoplast in plant provides further investigation of expression profiles and signals roles of corresponding proteomes, revealing more root contributions to pathogenic and symbiotic microbe interactions, as well as root-to-shoot communication [45]. Other proteomic studies focused mostly on the study of phloem sap exudates from agriculturally significant plants, such as oilseed rape [46], discovering hundreds of physiologically relevant proteins and ribo nucleoprotein complexes. The presence of proteins involved in redox control, defense and stress responses, calcium regulation, RNA metabolism, and G-protein signaling was increased in the phloem sap proteomes. Proteomics investigations gave some key insights into the working of the sieve tube system.

- **Root Proteomics to Increase Legume Productivity of Symbiotic Systems:** The symbiotic relationship between N-fixing bacteria and legumes results in the creation of root nodules, which is extremely significant in agriculture. A number of proteomic studies have sought to study the mutual interactions of symbiotic or pathogenic bacteria and host plant roots in the rhizosphere under a variety of biotic and abiotic soil conditions [47]. Proteomic analysis indicated diverse plant and bacteroid responses to drought stress. Two different groups reported protein expression investigations in nitrogen fixing root nodules of soybean and white sweet clover, respectively [48]. 2DE was also utilized to detect differentially expressed proteins during the bacterium *Sinorhizobium meliloti* strain 1021's symbiotic association with white sweet clover. When compared to the root, over 250 proteins were activated or up-regulated in the nodule, while over 350 proteins were down-regulated in the bacteroid form of the rhizobia. Bacteroid cells demonstrated a decrease in the expression of numerous proteins involved in nitrogen acquisition, indicating that the bacteroid was nitrogen proficient. Both results demonstrate the power of proteomics in plant symbiotic interactions [49].
2. **Recognizing Proteome Signatures For Crop Stress Tolerance:** Drought, heat, salinity, and infections are major environmental factors that threaten agricultural productivity and food security. Understanding the molecular pathways that underpin crop stress tolerance is critical for designing resilient cultivars. Stressful situations frequently result in delayed seed germination, reduced plant growth, and lower crop output. Komatsu and Hossain emphasized the importance of organ-specific proteome analysis for identifying proteins that often accumulate in organs under a variety of biotic stressors [32].

Jacoby et al. described how they used the newly developed proteomic technological advances of multiplexed selective-reaction monitoring MS to improve these approaches and provide a clear method to rank the relative significance of the growing cohort of stress-responsive proteins [36]. In addition to crops, proteomic approaches have been used to analyze various plants that serve as model systems in plant science as well as some agriculturally significant fruits under biotic and abiotic challenges [50]. Takahashi et al. investigated how plants respond to freezing stress, which causes serious problems in agricultural management, and discovered that the plasma membrane plays important roles in signal perception and cellular homeostasis, suggesting that plasma membrane proteins are the most significant variables in determining plant environmental stress tolerance [51]. Salt stress reduces agricultural output and growth significantly; yet, several crop varieties exhibit great resilience to the harmful effects of salt. Many salt-responsive proteins have been found in key crops, and they are expected to boost resistance to salt stress [52]. Hossain and Komatsu discussed current proteomic contributions to the knowledge of heavy metal stress responses in plants, notably the application of redox proteomic techniques for researching heavy metal-induced protein oxidation [32].

- **Differential Protein Expression:** Proteomic studies allow for the identification of proteins that are differentially expressed in response to stress. A proteomic investigation on drought-stressed rice, for example, discovered that numerous proteins involved in stress signaling, osmotic control, and reactive oxygen species scavenging

were elevated, underlining their role in drought tolerance [53]. These differentially expressed proteins act as proteomic fingerprints of crop stress resistance.

- **Stress-Specific Protein Markers:** Proteomic analysis enable the identification of stress-specific protein markers that can be used to measure crop stress tolerance. Researchers can find proteins consistently related with stress tolerance by comparing the proteomes of various crop varieties with varying stress tolerance. A study on salt-tolerant and salt-sensitive wheat types, for example, found particular proteins associated to ion transport, energy metabolism, and antioxidant defense as potential salt tolerance markers. These protein markers can be used to screen for and breed stress-tolerant crop varieties [4].
 - **Post-Translational Modifications:** Proteomic studies allow for the identification of stress-induced PTMs in crop proteins. A study on drought-stressed maize, for example, discovered phosphorylation events in proteins involved in photosynthesis, hormone signaling, and stress response pathways, shedding light on drought adaptation processes [54].
 - **Protein-Protein Interactions:** Proteomic techniques help identify stress-responsive interactions between proteins (PPIs) in crops. Identifying regulatory mechanisms and pathways of signaling involved in stress adaptation can be revealed by mapping PPI networks linked with stress tolerance. A study on salt-stressed soybean, for example, discovered distinct PPIs among proteins involved in ion homeostasis, reactive oxygen species detoxification, and hormone signaling, shedding light on the molecular basis of salt tolerance[11]. Understanding stress-responsive PPIs can help guide crop breeding for stress tolerance.
 - **Multi-Omics Integration:**Integrating proteomic data with other omics methods, including as genomes, transcriptomics, and metabolomics, allows for a more comprehensive knowledge of crop stress tolerance. A work combining transcriptomics and proteomics in drought-stressed maize, for example, identified novel regulatory pathways involved in drought adaptation [40].
3. **Post-Translational Modification:** Proteomics offers the great advantage of being able to detect changes in PTMs as well as changes in expression levels. Protein PTM analysis, such as phosphorylation and glycosylation, is critical for understanding issues like as activity, stability, and turnover. 2-DE and MS are particularly beneficial for the examination of expressed proteins with PTM, such as alkylation, glycosylation, and phosphorylation, which are thought to be the most significant regulatory proteins in a biological cell. Protein ubiquitination is an important regulatory process that regulates protein quantity, location, and activity. Several large-scale plant protein ubiquitination studies have been published [54]. In Arabidopsis, for example, affinity purification with an anti-ubiquity antibody and subsequent MS/MS analysis were used to identify ubiquitinated proteins.
 4. **Analyses of Food Quality, Safety and Nutritional Values:** The field of proteomics has been utilized to investigate differences in the nutritional qualities of food crops by

analyzing their proteomes. Heat stress, according to Mitsui et al., boosted the production of invertases in tomato fruits, increasing their sucrose content and creating sweeter tomatoes [54]. Proteomics studied why heat treatment for peach fruits improves peach fruit quality and shelf-life, and the cause was differentially expressed proteins involved in fruit growth and ripening [39]. Using an allergic patient's serum, a combination of 2-DE and IgE reactive proteins was used to evaluate the allergenicity of dietary proteins [50]. proteome examination of rice leaf, root, and seed revealed the presence of several allergenic proteins in the seeds, implying the utility of proteome analysis of foods for allergen detection [55].

- **Nutrient-Related Protein Identification:** Proteomic techniques allow for the identification and characterization of proteins. By comparing the proteomes of different crop types, researchers can find proteins linked with better nutritional properties. In wheat, for example, researchers have discovered proteins involved in the production of important amino acids and minerals, providing insights into nutritional quality improvement. Similarly, proteomic investigations in maize have discovered proteins involved in vitamin and antioxidant production [56]. The discovery of these nutrient-related proteins lays the groundwork for targeted breeding and genetic engineering approaches to improve nutritional value. Proteomics offers the great advantage of being able to detect changes in PTMs as well as changes in expression levels.
- **Nutrient Metabolism, Uptake and Transport :** Enzymes play an important role in crop nutrient metabolism. Proteomic studies have identified enzymes involved in food metabolism and accumulation. Proteomic investigations on tomato fruits revealed enzymes involved in carotenoid production, which is essential for vitamin A accumulation [56]. Researchers can improve crop nutritional quality by targeting these enzymes. Researchers have identified proteins linked with nutrient absorption and transporters in rice by studying the proteome of roots or conducting subcellular proteomics [45].
- **Nutrient Reserves and Storage Proteins:** Storage proteins act as a reservoir for vital nutrients in crops. Proteomic studies have revealed storage proteins linked to nutritional reserves. Proteomic investigations in legumes such as soybean and chickpea have discovered storage proteins rich in essential amino acids [57, 58].
- **Analyzing Metabolic Pathways:** Researchers can identify important proteins and enzymes that regulate metabolic pathways using the proteome under varied dietary conditions or through genetic alterations. Understanding these pathways makes it easier to optimize nutrient metabolism in crops. Through proteomics, some areas of research have shed information on metabolic pathways.
- **Post-Translational Modifications:** Proteomic analyses allow for the identification of PTMs associated with nutrient-related proteins. PTMs impacting nutrient-related proteins in crops, for example, have been found as phosphorylation, glycosylation, and acetylation. Understanding the function of PTMs can help us understand nutrition control and metabolism [3].
- **Photosynthesis-related proteins for nutrient assimilation:** Photosynthesis-related proteins have been linked to nutritional enhancement in studies. Carbon fixation and

chlorophyll synthesis proteins were shown to be linked with grain zinc and iron levels in wheat [59]. Understanding how these proteins are regulated can lead to techniques for increasing nutritional content in crops. Scientists discovered genotype-specific changes in protein profiles in rice that are connected with amino acid composition and nutritional quality.

- **Impact of Genetic and Environmental Factors:** Genetic and environmental factors have a well-known impact on nutritional quality. Researchers can uncover proteins that alter nutrient accumulation and composition by comparing the proteomes of various genotypes or under different environmental conditions. Proteomic studies on rice have revealed genotype-specific changes in protein profiles linked to amino acid content and nutritional quality as a result of environmental variables [60].
 - **Genetic Diversity:** Proteomic investigations of nutrient-related proteins in different crop varieties have revealed genetic diversity. Genetic differences in proteins linked with amino acid metabolism and grain nutritional quality in maize have been uncovered by researchers [60]. These genetic variances can help direct breeding operations to improve crop nutritional content.
 - **Environmental Stress:** Drought, heat, and nutrient deficit can all have an impact on crop nutritional quality. Wheat proteomic investigations indicated alterations in gluten proteins during heat stress, influencing the nutritional quality of wheat grains[61]. This can aid in reducing the harmful impacts of environmental stress on nutritional content.

5. **Role of Proteomics in Personalized Nutrition:** Because of individual differences in dietary requirements and responses, proteomic investigations have the potential to contribute to customized nutrition techniques. Researchers can gain a full understanding of the relationship between genetics, nutrition, and health outcomes by merging proteomic data with other omics technologies such as genomics and metabolomics. Proteomic profiles can be used to measure nutritional status and metabolic health.

Proteomic investigations, for example, have revealed protein biomarkers linked to dietary treatments and illness risk [62]. Integrating proteomics with personalized nutrition techniques has the potential to improve dietary recommendations and individual health outcomes. By combining proteomic and genomic data, it is possible to identify candidate genes associated with nutrient-related proteins. Researchers can link genetic variants to protein expression and function by comparing proteomic and genomic data. This combination makes marker-assisted selection strategies for breeding nutrient-rich crops more feasible [63]. Integrating proteomic and transcriptome data provides insights into the transcriptional regulation of nutrient-related proteins. Researchers can uncover critical transcription factors and regulatory processes involved in food metabolism by connecting changes in protein abundance with transcript levels. This integration provides a more comprehensive understanding of the molecular mechanisms governing nutrient increase.

6. **Harnessing Proteomic Data for Yield Enhancement in Crops:** Crop yield increase is critical to meeting the world's growing food demand. Researchers can improve crop output by combining proteomic data to find essential regulatory proteins, reveal signaling

pathways, and discover biomarkers associated with yield-related features. This chapter looks at how to use proteomic data to improve yield, such as identifying yield-associated proteins, understanding yield-limiting variables, and integrating proteomics with other omics technologies.

- **Identification of Yield-Associated Proteins:** Proteomic analysis allow for the identification of proteins linked with high-yielding characteristics in crops. Researchers can identify proteins that play a critical role in yield enhancement by comparing the proteomes of high-yielding and low-yielding cultivars or under different yield situations. Different proteomic techniques, such as two-dimensional gel electrophoresis (2-DE) and mass spectrometry (MS), make it easier to identify and quantify differentially expressed proteins.
 - **Photosynthesis-Related Proteins for yield:** Photosynthesis-related proteins have been linked to high yields in studies. Maurya et al. discovered that numerous proteins involved in photosynthetic electron transport and carbon fixation were positively linked with grain production in rice. These proteins could be used as targets for yield-boosting techniques [64].
 - **Protein Synthesis and Metabolism:** Proteins involved in protein synthesis and metabolism have been connected directly to crop yield enhancement. Ribosomal proteins and enzymes involved in protein synthesis and metabolic pathways have been identified as possible yield-associated proteins in wheat and maize studies [65].
 - **Integration of Proteomics with Other Omics Technologies:** Integrating proteomics with other omics technologies such as genomes, transcriptomics, and metabolomics allows for a more complete understanding of crop yield enhancement. Researchers can decipher complex biological networks and find essential regulators by merging multi-omics datasets. Integrative techniques offer a systems-level view of yield enhancement processes and allow for the identification of possible crop improvement targets. This integration of proteomics and genomics allows for the identification of genetic variables that influence crop productivity as well as the use of marker-assisted selection in breeding programs [66]. Researchers can link genetic variants to protein expression and function by comparing proteomic and genomic data. Researchers can identify critical transcription factors and regulatory pathways involved in yield enhancement by correlating changes in protein abundance with transcript levels [67]. Integrating proteome and metabolomic data enables a thorough understanding of metabolic pathways and their link to yield attributes. Key enzymes and metabolic intermediates connected with yield-related processes can be identified by researchers.
7. **Exploring Potential For Disease Resistance In Crops:** By studying the whole set of proteins expressed in plants, researchers can uncover critical proteins and processes involved in plant defense responses, allowing the development of new disease-resistant crop varieties. One technique in disease resistance proteomics research is to compare the proteomes of resistant and susceptible crop varieties or lines after pathogen infection. This allows for the identification of proteins that differ in expression or undergo post-translational modifications (PTMs) in response to the pathogen. 68. Santamaría-Hernando Set al. investigated the response of tomato plants to infection by the bacterial disease *Pseudomonas syringae* using proteomics analysis. Pathogenesis-related proteins,

antioxidant enzymes, and proteins involved in signal transduction pathways were identified by the researchers as proteins connected with defense responses [68].

Protein-protein interaction (PPI) analysis, in addition to differential protein expression analysis, has been used in proteomics research for disease resistance. Mapping protein interactions in defensive signaling can uncover regulatory networks and critical players in plant immunity. Zhenget al., for example, used PPI analysis to investigate the immunological response of rice plants to the blast fungus *Magnaportheoryzae*[69]. The researchers discovered links between proteins important in defense signaling pathways, shedding light on the mechanisms behind rice resistance to the virus.

Furthermore, proteomics techniques have been useful in researching post-translational modifications (PTMs) linked to plant disease resistance. PTMs such as phosphorylation, ubiquitination, and glycosylation can influence plant immunological responses by modulating protein activity and stability. Chivasaet al. (2009), for example, studied the phosphorylation processes that occur in Arabidopsis plants when infected with the oomycete disease *Hyaloperonosporaarabidopsidis*[70]. The researchers discovered phosphorylated proteins during infection, including defense-related proteins and proteins implicated in signal transduction cascades.

Proteomics research has also helped to identify candidate proteins and indicators related with disease resistance in crops. Researchers can identify proteins that confer resistance and use them as breeding or genetic engineering targets by studying the proteome of resistant varieties or wild relatives of crops. Castillejo et al., for example, studied the proteome of *Medicago truncatula*, a legume plant with natural resistance to the parasitic weed *Orobanchecrenata*[71]. The researchers discovered proteins involved in defense responses and parasite detection that could be used to generate crop varieties resistant to parasitic weeds. Researchers can uncover critical proteins and processes involved in plant defensive responses by examining the proteome, protein interactions, and post-translational modifications. This knowledge can be used to create superior crop types with increased disease resistance, hence promoting sustainable and resilient agriculture.

VI. APPLICATION OF ARTIFICIAL INTELLIGENCE IN PROTEOMICS

While significant advancement has been achieved in proteomics, the analysis of proteomics remains time and cost-intensive. With the passage of time, the use of digital tools such as Artificial Intelligence (AI) expands and opens up significant prospects in proteomics to address the issues confronting scientific researchers. Machine learning (ML) is quickly becoming a very popular tool in the field of proteomics due to the increasing availability of publicly available proteomics data. Machine learning (ML) improvements, particularly deep learning (DL), are catalyzing a paradigm shift in science study. State-of-the-art machine learning architectures like as Convolutional Neural Networks (CNNs) and Recurrent Neural Networks (RNNs) have been used to analyse huge and complex proteomic data. The fundamental difference between DL and traditional machine learning techniques such as Support Vector Machine (SVM) and Random Forests (RF) is that DL can learn features and patterns from data automatically. As a result, DL is particularly well suited to scientific domains with large and complex datasets. Recent advances in proteomics have sparked

interest in DL approaches with applications in MS/MS spectrum prediction, PTMs de novo peptide sequencing, protein structure, and structure and function prediction.

Several DL models have been employed and developed for reliable peptide and protein identification from MS data, with the goal of significantly lowering instrument time and costs (Table 2). DL applications in MS-based proteomics include i) improving peptide recognition sensitivity in database scanning, ii) serving as a peptide identification quality appraisal metric, iii) developing spectrum libraries for Data Independent Acquisition (DIA) data analysis, and iv) encouraging targeted proteomics experiments [72].

Table 2: DL models and its description

S.No.	Tools	Description	References
1.	MSpectraAI (Mass Spectra Artificial Intelligence)	Uses deep neural networks (DNNs) to analyze MS data	[73]
2.	DeepIso	Combines CNN and RNN categorizes peptide characteristics using the LCMS/MS plot	[74]
3.	MusiteDeep	Visualizes protein PTM sites based on protein sequences	[75]
4.	DeepUbi	DL frameworks that predict ubiquitination sites	[76]
5.	DeepSuccinylSite	DL frameworks that predict succinylation sites	[77]
6.	DeepACLSTM	DL models for secondary structure prediction, protein function perfection, protein categorization, and protein subcellular localization	[78]
7.	SDN2GO		[79]
8.	UDSMProt		[80]
9.	PSLCNN		[81]
10.	AlphaFold		[82]
11.	C-I-TASSER	Widely employed in conjunction with higher resolution physical modeling.	[83]

VII. CLOUD COMPUTING AND BIG DATA ANALYTICS IN PROTEOMICS

The employment of next generation sequencing and MS technology to research experimental models and crop plants is undoubtedly attributable to the exponential development in the quantity of plant data sets over the last decade. Due to the exorbitant cost of software and high-end servers as well as the lengthy processing times needed for research studies, such development has also created unbelievable difficulties for individual laboratories to store, maintain, and interpret data. There are new challenges in dealing with massive amounts of proteomics data and information. Calculating big data sets takes an impractically long time [84].MassIVE (member of the Proteome Xchange collaboration) , ProteoSAFe, Galaxy-P, and Chorus [85]Exchange Consortium), ProteoSAFe, Galaxy-P, and Chorus [85] are some of the most recent significant proteomic systems to enable data repository, data processing, and are available at Anaconda Cloud for providing a flexible and scalable service for groups and organizations of all sizes.

Cloud hosting allows users to rent these resources from a cloud service provider rather than owning and housing computing, data storage, and other facilities locally, and merely pay for the time spent. A number of cloud service companies today offer a wide range of offerings, with Amazon Web Services (AWS) being the main service that has been used for a wide range of programming activities in bioinformatics. Cloud computing substantially speeds up data analysis and allows researchers to do more complicated analyses. MS-PyCloud is an open source, cloud computing-based pipeline for protein inference, protein PTM evaluation, and MS/MS database scan for spectrum assignment for LC- MS/ MS dataprocessing [86].

QCloud is a cloud-based quality control solution for MS-based proteomics frameworks [87]. The technology eliminates the barriers that often impede the adoption of quality control equipment and provides proteomics laboratories with a system that is ready to use. Firmiana is a proteomic cloud platform that allows researchers to upload raw data from mass spectrometry (MS) files, execute online proteome recognition and quantification, run bioinformatics tests, extract data, and present findings without programming skills [88]. Using the Institute of Systems Biology's first trans-proteomic pipeline (TPP) on the Amazon cloud service two years ago, over 1100 raw data were analyzed in 9 hours.

ProteoCloud (<http://proteocloud.googlecode.com>), a free and open pipeline of applications that allows end-users to easily identify very large MS/MS spectra sets using five distinct methods via cloud computing technologies [89].

VIII. CHALLENGES AND FUTURE PERSPECTIVES

Proteomics has evolved as a potent crop enhancement technique, giving important insights into the molecular mechanisms driving plant characteristics and responses. However, various hurdles must be overcome before the full potential of proteomics in crop enhancement may be realized. Furthermore, there are interesting future prospects that can improve the application of proteomics in this industry. The complexity of the plant proteome is one of the main obstacles in proteomics-based crop development. Plants have a huge and diversified proteome, with proteins ranging in abundance and dynamic range. Because of this intricacy, complete proteome analysis is difficult because extremely abundant proteins might overwhelm low-abundance regulatory proteins. Furthermore, the presence of proteins with high hydrophobicity or extreme molecular weights can make detection and analysis difficult. Overcoming these technical obstacles and establishing better proteomics procedures will be critical for gaining a more complete understanding of the plant proteome. Another challenge is determining the function of discovered proteins. While proteomics can uncover proteins that are differently expressed or changed in relation to specific attributes, identifying their functional roles and how they contribute to phenotypic alterations necessitates more research and validation. To establish the causal association between identified proteins and desired features, functional investigations such as gene deletion or overexpression are required. Integrating proteomics data with other omics methods, like as genomics, transcriptomics, and metabolomics, can also help with functional protein identification and provide a more comprehensive understanding of plant biology.

Furthermore, high-quality reference proteome datasets for many crop species are critical for proteomics research. For reliable protein identification and annotation, extensive

and well-curated databases containing protein sequences, post-translational modifications, and protein-protein interaction information are required. Continuing efforts to establish and upgrade reference proteome databases will allow for more precise and comprehensive proteomics research in crop improvement. Despite these obstacles, proteomics has strong future prospects in crop improvement. The advancement of high-throughput and sensitive proteomics technology is one such viewpoint. Technological advances in mass spectrometry instrumentation, sample preparation methods, and data analysis algorithms are increasing the depth and accuracy of proteomics analysis.

This will allow for more thorough and quantitative characterization of the plant proteome, making it easier to identify critical proteins and pathways linked to crucial agronomic features. Another future possibility is to combine proteomics with other omics methods and phenotypic data. Integrating proteomics data with genomes, transcriptomics, metabolomics, and phenotypic data can lead to a more comprehensive understanding of plant biology and trait regulation. This integration can assist identify candidate genes and proteins for crop improvement, as well as untangle complicated regulatory networks that govern essential features. Furthermore, the introduction of focused proteomics methods, such as selected reaction monitoring (SRM) and parallel reaction monitoring (PRM), holds promise for more precise and quantitative protein analysis.

Targeted proteomics enables the targeted quantification of proteins of interest in large-scale investigations, allowing for the monitoring of protein expression and changes. While proteomics holds enormous promise for crop development, there are several obstacles to overcome, such as the complexity of the plant proteome and the functional characterization of discovered proteins. However, with the advancement of proteomics technologies, integration with other omics approaches, and the creation of comprehensive reference proteome databases, proteomics will continue to play an important role in unraveling the molecular mechanisms underlying crop traits and enabling targeted crop improvement.

IX. CONCLUSION

Proteomics has emerged as a strong crop improvement technique, revealing important insights into the molecular mechanisms underpinning plant characteristics, stress responses, and disease resistance. Researchers can uncover critical proteins and pathways linked with significant agronomic features, stress tolerance, and defense systems by performing a complete investigation of the plant proteome. This understanding opens up new opportunities for focused agricultural improvement, enabling for the production of crops with increased yield, resilience, and resource efficiency. While challenges in proteomics-based crop improvement exist, such as the complexity of the plant proteome and the functional characterization of identified proteins, advances in proteomics technologies, integration with other omics approaches, and the development of comprehensive reference proteome databases offer promising solutions. These advances enable more extensive and quantitative characterization of the plant proteome, facilitate the discovery of candidate proteins and regulatory networks, and contribute to a systems-level knowledge of plant biology.

Plant proteomics studies have improved significantly in recent years due to major advancements in proteomics methodologies and bioinformatics tools. This paradigm shift has improved our ability to recognize plant-pathogen interactions, PTM, disease resistance, and

stress responses. Deep learning methods have enormous promise in delivering major insights into regulatory systems such as response to abiotic stressors, given ongoing developments in plant proteomics and the availability of high quality proteomics data. Crop proteomics has advanced rapidly thanks to novel biotechnology approaches and high throughput omics technologies, which will aid in enhancing crop quality and meeting food production targets by 2050.

The future of proteomics in crop development is bright. Technological advancements in high-throughput and sensitive proteomics, integration with other omics techniques and phenotypic data, and the emergence of focused proteomics approaches enable more precise and quantitative protein analysis. These advances will make it easier to identify essential proteins and pathways linked to desired features, laying the groundwork for the generation of superior crop types through breeding or genetic engineering. Crop improvement driven by proteomics has the potential to greatly contribute to sustainable agriculture by increasing crop output, stress tolerance, and disease resistance.

Targeted proteomics enables the targeted quantification of proteins of interest in large-scale investigations, allowing for the monitoring of protein expression and changes. While proteomics holds enormous promise for crop development, there are several obstacles to overcome, such as the complexity of the plant proteome and the functional characterization of discovered proteins. However, with the advancement of proteomics technologies, integration with other omics approaches, and the creation of comprehensive reference proteome databases, proteomics will continue to play an important role in unraveling the molecular mechanisms underlying crop traits and enabling targeted crop improvement.

REFERENCES

- [1] Beddington, John R., Mohammed Asaduzzaman, Megan E. Clark, Adrian FernándezBremauntz, Marion D. Guillou, Molly M. Jahn, Erda Lin et al. "The role for scientists in tackling food insecurity and climate change." *Agriculture & Food Security* 1, no. 1 (2012): 1-9.
- [2] Hernández-Elvira, Mariana, and Per Sunnerhagen. "Post-transcriptional regulation during stress." *FEMS Yeast Research* 22, no. 1 (2022): foac025.
- [3] Hashiguchi A, Komatsu S. Impact of post-translational modifications of crop proteins under abiotic stress. *Proteomes*. 2016 Dec 21;4(4):42.
- [4] Ashraf, M. "Stress-induced changes in wheat grain composition and quality." *Critical reviews in food science and nutrition* 54, no. 12 (2014): 1576-1583.
- [5] Bashir, Humayra, M. Irfan Qureshi, SowbiyaMuneer, Javed Ahmad, and LelloZolla. "Proteomic approaches to map thylakoid proteins and study differential protein expression under various abiotic stresses." In *Proceedings of International Conference of Biology, Biochemistry and Biotechnology (ICBBB)*, World Academy of Science, Engineering & Technology, Rome, pp. 28-30. 2010.
- [6] Ahmad, Parvaiz, Arafat AH Abdel Latef, SaiemaRasool, Nudrat A. Akram, Muhammad Ashraf, and SalihGucel. "Role of proteomics in crop stress tolerance." *Frontiers in plant science* 7 (2016): 1336.
- [7] Figeys, Daniel. "Proteomics: the basic overview." *Industrial Proteomics: Applications for Biotechnology and Pharmaceuticals* 45 (2005): 1.
- [8] Lorefice, Lorena, MaristellaPitzalis, Federica Murgia, Giuseppe Fenu, Luigi Atzori, and Eleonora Cocco. "Omics approaches to understanding the efficacy and safety of disease-modifying treatments in multiple sclerosis." *Frontiers in Genetics* 14 (2023): 1076421.
- [9] Sharma, Jyoti, LavanyaBalakrishnan, Sandeep Kaushik, and Manoj Kumar Kashyap. "multi-omics approaches to study signaling pathways." *Frontiers in Bioengineering and Biotechnology* 8 (2020): 829.
- [10] Graves, Paul R., and Timothy AJ Haystead. "Molecular biologist's guide to proteomics." *Microbiology and molecular biology reviews* 66, no. 1 (2002): 39-63.

- [11] Bludau, Isabell, and Ruedi Aebersold. "Proteomic and interactomic insights into the molecular basis of cell functional diversity." *Nature Reviews Molecular Cell Biology* 21, no. 6 (2020): 327-340.
- [12] Van Den Bossche, Tim, Magnus Ø. Arntzen, Dörte Becher, Dirk Benndorf, Vincent GH Eijnsink, Céline Henry, Pratik D. Jagtap et al. "The Metaproteomics Initiative: a coordinated approach for propelling the functional characterization of microbiomes." *Microbiome* 9, no. 1 (2021): 1-4.
- [13] Shi, Yang, Rong Xiang, Csaba Horváth, and James A. Wilkins. "The role of liquid chromatography in proteomics." *Journal of Chromatography A* 1053, no. 1-2 (2004): 27-36.
- [14] Le GRICE, Stuart FJ, and Fiona GRÜNINGER-LEITCH. "Rapid purification of homodimer and heterodimer HIV-1 reverse transcriptase by metal chelate affinity chromatography." *European journal of biochemistry* 187, no. 2 (1990): 307-314.
- [15] Li, Xiaoming, Hui Bai, Xianyun Wang, Liyun Li, Yinghao Cao, Jian Wei, Yumeng Liu et al. "Identification and validation of rice reference proteins for western blotting." *Journal of experimental botany* 62, no. 14 (2011): 4763-4772.
- [16] Kollerová, E., M. Glasa, and Z. W. Šubr. "Western blotting analysis of the Plum pox virus capsid protein." *Journal of Plant Pathology* (2008): S19-S22.
- [17] Wellhausen, Robert, and Harald Seitz. "Facing current quantification challenges in protein microarrays." *Journal of Biomedicine and Biotechnology* 2012 (2012).
- [18] Wilson, David S., and Steffen Nock. "Functional protein microarrays." *Current opinion in chemical biology* 6, no. 1 (2002): 81-85.
- [19] Mueller, Claudius, Lance A. Liotta, and Virginia Espina. "Reverse phase protein microarrays advance to use in clinical trials." *Molecular oncology* 4, no. 6 (2010): 461-481.
- [20] Baldelli, Elisa, Valerie Calvert, Alex Hodge, Amy VanMeter, Emanuel F. Petricoin, and Mariaelena Pierobon. "Reverse phase protein microarrays." *Molecular profiling: methods and protocols* (2017): 149-169.
- [21] Zahedi, René P., Jan Moebius, and Albert Sickmann. "Two-dimensional BAC/SDS-PAGE for membrane proteomics." *Subcellular Proteomics: From Cell Deconstruction to System Reconstruction* (2007): 13-20.
- [22] Neilson, Karlie A., Iniga S. George, Samantha J. Emery, Sridevi Muralidharan, Mehdi Mirzaei, and Paul A. Haynes. "Analysis of rice proteins using SDS-PAGE shotgun proteomics." *Plant proteomics: Methods and protocols* (2014): 289-302.
- [23] Serrano, Solange MT, John D. Shannon, Deyu Wang, Antonio CM Camargo, and Jay W. Fox. "A multifaceted analysis of viperid snake venoms by two-dimensional gel electrophoresis: an approach to understanding venom proteomics." *Proteomics* 5, no. 2 (2005): 501-510.
- [24] Calvo, Enrique, M. Graciela Pucciarelli, Hélène Bierne, Pascale Cossart, Juan Pablo Albar, and Francisco García-del Portillo. "Analysis of the *Listeria* cell wall proteome by two-dimensional nanoliquid chromatography coupled to mass spectrometry." *Proteomics* 5, no. 2 (2005): 433-443.
- [25] Pocsfalvi, Gabriella, Giuseppina Cacace, Manuela Cuccurullo, Giovanna Serluca, Alida Sorrentino, Gitta Schlosser, Giuseppe Blaiotta, and Antonio Malorni. "Proteomic analysis of exoproteins expressed by enterotoxigenic *Staphylococcus aureus* strains." *Proteomics* 8, no. 12 (2008): 2462-2476.
- [26] Aebersold, Ruedi, and Matthias Mann. "Mass spectrometry-based proteomics." *Nature* 422, no. 6928 (2003): 198-207.
- [27] Shen, Shihua, Masami Matsubae, Toshifumi Takao, Naoki Tanaka, and Setsuko Komatsu. "A proteomic analysis of leaf sheaths from rice." *The journal of biochemistry* 132, no. 4 (2002): 613-620.
- [28] Chaube, Ruchi. "Absolute quantitation of post-translational modifications." *Frontiers in Chemistry* 2 (2014): 58.
- [29] Kumar, Chanchal, and Matthias Mann. "Bioinformatics analysis of mass spectrometry-based proteomics data sets." *FEBS letters* 583, no. 11 (2009): 1703-1712.
- [30] Rigbolt, Kristoffer TG, Jens T. Vanselow, and Blagoy Blagoev. "GProX, a user-friendly platform for bioinformatics analysis and visualization of quantitative proteomics data." *Molecular & Cellular Proteomics* 10, no. 8 (2011).
- [31] Chen, Yanmei, Yi Wang, Jun Yang, Wenbin Zhou, and Shaojun Dai. "Exploring the diversity of plant proteome." *Journal of Integrative Plant Biology* 63, no. 7 (2021): 1197-1210.
- [32] Hossain, Zahed, and Setsuko Komatsu. "Contribution of proteomic studies towards understanding plant heavy metal stress response." *Frontiers in Plant Science* 3 (2013): 310.
- [33] Sharma, Jitendra Kumar, Monika Sihmar, Anita Rani Santal, and N. P. Singh. "Impact assessment of major abiotic stresses on the proteome profiling of some important crop plants: a current update." *Biotechnology and Genetic Engineering Reviews* 35, no. 2 (2019): 126-160.

- [34] Nukarinen, Ella, Thomas Nägele, Lorenzo Pedrotti, Bernhard Wurzinger, Andrea Mair, Ramona Landgraf, Frederik Börnke et al. "Quantitative phosphoproteomics reveals the role of the AMPK plant ortholog SnRK1 as a metabolic master regulator under energy deprivation." *Scientific Reports* 6, no. 1 (2016): 31697.
- [35] Kazan, Kemal. "Diverse roles of jasmonates and ethylene in abiotic stress tolerance." *Trends in plant science* 20, no. 4 (2015): 219-229.
- [36] Jacoby, Richard P., A. Harvey Millar, and Nicolas L. Taylor. "Application of selected reaction monitoring mass spectrometry to field-grown crop plants to allow dissection of the molecular mechanisms of abiotic stress tolerance." *Frontiers in Plant Science* 4 (2013): 20.
- [37] Urban, Milan Oldřich, Jakub Vašek, Miroslav Klíma, Jana Krtková, KláraKosová, Ilja Tom Prášil, and Pavel Vítámvás. "Proteomic and physiological approach reveals drought-induced changes in rapeseeds: Water-saver and water-spender strategy." *Journal of Proteomics* 152 (2017): 188-205.
- [38] Roberts, Irma N., Carla Caputo, María Victoria Criado, and Christiane Funk. "Senescence-associated proteases in plants." *PhysiologiaPlantarum* 145, no. 1 (2012): 130-139.
- [39] Zhang, Li, Zhifang Yu, Li Jiang, Juan Jiang, Haibo Luo, and Linran Fu. "Effect of post-harvest heat treatment on proteome change of peach fruit during ripening." *Journal of proteomics* 74, no. 7 (2011): 1135-1149.
- [40] Zhang, Chong, Jiaxue Zhang, Yadi Liu, Xiatong Liu, XiaoruiGuo, Hui Li, Di Liu, and Hai Lu. "Integrated Transcriptomic and Proteomic Analysis in the Roadmap of the Xylem Development Stage in *Populustomentosa*." *Frontiers in Plant Science* 12 (2021): 724559.
- [41] Wang, Xiaoqin, Yanli Liu, and Pingfang Yang. "Proteomic studies of the abiotic stresses response in model moss–*Physcomitrella patens*." *Frontiers in plant science* 3 (2012): 258.
- [42] Alvarez, Sophie, Ellen L. Marsh, Steve G. Schroeder, and Daniel P. Schachtman. "Metabolomic and proteomic changes in the xylem sap of maize under drought." *Plant, Cell & Environment* 31, no. 3 (2008): 325-340.
- [43] Leyva-González, Marco Antonio, Enrique Ibarra-Laclette, Alfredo Cruz-Ramírez, and Luis Herrera-Estrella. "Functional and transcriptome analysis reveals an acclimatization strategy for abiotic stress tolerance mediated by *Arabidopsis* NF-YA family members." *PloS one* 7, no. 10 (2012): e48138.
- [44] Agrawal, Ganesh Kumar, Nam-Soo Jwa, Marc-Henri Lebrun, Dominique Job, and RandeepRakwal. "Plant secretome: unlocking secrets of the secreted proteins." *Proteomics* 10, no. 4 (2010): 799-827.
- [45] Hu, Junjie, Christof Rampitsch, and Natalia V. Bykova. "Advances in plant proteomics toward improvement of crop productivity and stress resistance." *Frontiers in Plant Science* 6 (2015): 209.
- [46] Fröhlich, Andreas, Frank Gaupels, HakanSarioglu, Christian Holzmeister, Manuel Spannagl, JörgDurner, and Christian Lindermayr. "Looking deep inside: detection of low-abundance proteins in leaf extracts of *Arabidopsis* and phloem exudates of pumpkin." *Plant physiology* 159, no. 3 (2012): 902-914.
- [47] Knief, Claudia, Nathanael Delmotte, and Julia A. Vorholt. "Bacterial adaptation to life in association with plants—A proteomic perspective from culture to in situ conditions." *Proteomics* 11, no. 15 (2011): 3086-3105.
- [48] Reid, Dugald E., Satomi Hayashi, Michal Lorenc, Jiri Stiller, David Edwards, Peter M. Gresshoff, and Brett J. Ferguson. "Identification of systemic responses in soybean nodulation by xylem sap feeding and complete transcriptome sequencing reveal a novel component of the autoregulation pathway." *Plant Biotechnology Journal* 10, no. 6 (2012): 680-689.
- [49] Molesini, Barbara, Daniela Cecconi, YouryPii, and TizianaPandolfini. "Local and systemic proteomic changes in *Medicago truncatula* at an early phase of *Sinorhizobium meliloti* infection." *Journal of Proteome Research* 13, no. 2 (2014): 408-421.
- [50] Chan, Zhulong. "Proteomic responses of fruits to environmental stresses." *Frontiers in Plant Science* 3 (2013): 311.
- [51] Takahashi, Daisuke, Bin Li, Takato Nakayama, Yukio Kawamura, and Matsuo Uemura. "Plant plasma membrane proteomics for improving cold tolerance." *Frontiers in plant science* 4 (2013): 90.
- [52] Beyene, Belachew, and Gizachew Haile. "Review on proteomics technologies and its application for crop improvement." *Innovative Systems Design and Engineering* (2016).
- [53] Vantini, J. S., G. C. Dedemo, D. F. JovinoGimenez, L. F. Fonseca, R. I. Tezza, M. A. Mutton, J. A. Ferro, and M. I. T. Ferro. "Differential gene expression in drought-tolerant sugarcane roots." *Genet. Mol. Res* 14, no. 2 (2015): 7196-7207.
- [54] Mitsui, Toshiaki, Takeshi Shiraya, Kentaro Kaneko, and Kaede Wada. "Proteomics of rice grain under high temperature stress." *Frontiers in Plant Science* 4 (2013): 36.

- [55] Aghaei, Keyvan, and Setsuko Komatsu. "Crop and medicinal plants proteomics in response to salt stress." *Frontiers in Plant Science* 4 (2013): 8.
- [56] Wang, Rong, FeiGao, Bing-Qian Guo, Ji-Chang Huang, Lei Wang, and Yi-Jun Zhou. "Short-term chromium-stress-induced alterations in the maize leaf proteome." *International Journal of Molecular Sciences* 14, no. 6 (2013): 11125-11144.
- [57] Singh N, Jain P, Ujwal M, Langyan S. Escalate protein plates from legumes for sustainable human nutrition. *Front Nutr.* 2022 Nov 4;9:977986.
- [58] Masclaux-Daubresse C, Daniel-Vedele F, Dechorgnat J, Chardon F, Gaufichon L, Suzuki A. Nitrogen uptake, assimilation and remobilization in plants: challenges for sustainable and productive agriculture. *Ann Bot.* 2010;105(7):1141-1157.
- [59] Mallén-Ponce MJ, Pérez-Pérez ME, Crespo JL. Photosynthetic assimilation of CO₂ regulates TOR activity. *Proc Natl AcadSci U S A.* 2022;119(2):e2115261119.
- [60] Romanov N, Kuhn M, Aebersold R, Ori A, Beck M, Bork P. Disentangling Genetic and Environmental Effects on the Proteotypes of Individuals. *Cell.* 2019;177(5):1308-1318.e10.
- [61] Enroth S, Bosdotter Enroth S, Johansson Å, Gyllensten U. Effect of genetic and environmental factors on protein biomarkers for common non-communicable disease and use of personally normalized plasma protein profiles (PNPPP). *Biomarkers.* 2015;20(6-7):355-364.
- [62] Chaudhary N, Kumar V, Sangwan P, Pant N C, Saxena A, Joshi S et al. Personalized Nutrition and -Omics. *Comprehensive Foodomics.* 2021;495-507.
- [63] Jain KK. Role of Proteomics in the Development of Personalized Medicine. *Adv Protein ChemStruct Biol.* 2016;102:41-52.
- [64] Maurya VK, Gupta SK, Sharma M, Majumder B, Deeba F, Pandey N, et al. Proteomic changes may lead to yield alteration in maize under carbon dioxide enriched condition. *3 Biotech.* 2020;10(5):203.
- [65] Wang S, Chen W, Yang C, Yao J, Xiao W, Xin Y, Qiu J, et al. Comparative proteomic analysis reveals alterations in development and photosynthesis-related proteins in diploid and triploid rice. *BMC Plant Biol.* 2016;16(1):199.
- [66] Zhang B, Kuster B. Proteomics Is Not an Island: Multi-omics Integration Is the Key to Understanding Biological Systems. *Mol Cell Proteomics.* 2019;18(8 suppl 1):S1-S4.
- [67] Jendoubi T. Approaches to Integrating Metabolomics and Multi-Omics Data: A Primer. *Metabolites.* 2021;11(3):184. Published 2021 Mar 21.
- [68] Santamaría-Hernando S, López-Maroto Á, Galvez-Roldán C, Munar-Palmer M, Monteagudo-Cascales M, Rodríguez-Herva J, et al. Pseudomonas syringae pv. tomato infection of tomato plants is mediated by GABA and l-Pro chemoperception. *Mol Plant Pathol.* 2022;23(10):1433-1445.
- [69] Zheng C, Liu Y, Sun F, Zhao L, Zhang L. Predicting Protein-Protein Interactions Between Rice and Blast Fungus Using Structure-Based Approaches. *Front Plant Sci.* 2021;12:690124.
- [70] Massoud K, Barchietto T, Le Rudulier T, Pallandre L, Didierlaurent L, Garmier M, et al. Dissecting phosphite-induced priming in Arabidopsis infected with *Hyaloperonospora arabidopsidis*. *Plant Physiol.* 2012;159(1):286-298.
- [71] Castillejo MA, Maldonado AM, Dumas-Gaudot E, Fernández-Aparicio M, Susín R, Diego R et al. Differential expression proteomics to investigate responses and resistance to *Orobanchecrenata* in *Medicago truncatula*. *BMC Genomics.* 2009;10:294.
- [72] Wen, B., Zeng, W.F., Liao, Y., Shi, Z., Savage, S.R., Jiang, W. and Zhang, B. Deep Learning in Proteomics. *Proteomics*, 2020. 20: 1900335.
- [73] Wang, S., Zhu, H., Zhou, H., Cheng, J. and Yang, H. MSpectraAI: A Powerful Platform for Deciphering Proteome Profiling of Multi-Tumor Mass Spectrometry Data by Using Deep Neural Networks. *BMC Bioinformatics*, 2020. 21(1): 1-15.
- [74] Zohora, F.T., Rahman, M.Z., Tran, N.H., Xin, L., Shan, B. and Li, M. DeepIso: A Deep Learning Model for Peptide Feature Detection from LCMS Map. *Scientific Reports*, 2019. 9(1): 1-13.
- [75] Wang, D., Liu, D., Yuchi, J., He, F., Jiang, Y., Cai, S., Li, J. and Xu, D. MusiteDeep: A Deep-Learning Based Webserver for Protein Post Translational Modification Site Prediction and Visualization. *Nucleic Acids Research*, 2020.48(W1): W140-W146.
- [76] Fu, H., Yang, Y., Wang, X., Wang, H. and Xu, Y. DeepUbi: A Deep Learning Framework for Prediction of Ubiquitination Sites in Proteins, *BMC Bioinformatics*, 2019. 20(1): 1-10.
- [77] Thapa, N., Chaudhari, M., McManus, S., Roy, K., Newman, R.H., Saigo, H. and Kc, D.B. DeepSuccinylSite: A Deep Learning Based Approach for Protein Succinylation Site Prediction. *BMC Bioinformatics*, 2020. 21: 1-10.

- [78] Guo, Y., Li, W., Wang, B., Liu, H. and Zhou, D .DeepACLST: Deep Asymmetric Convolutional Long Short-Term Memory Neural Models for Protein Secondary Structure Prediction. *BMC Bioinformatics*, 2019. 20(1):1-12.
- [79] Cai, Y., Wang, J. and Deng, L. SDN2GO: An Integrated Deep Learning Model for Protein Function Prediction. *Frontiers in Bioengineering and Biotechnology*, 2020. 8:391.
- [80] Strothoff, N., Wagner, P., Wenzel, M. and Samel, W . U D S M P r o t : Universal Deep Sequence Models for Protein Classification. *Bioinformatics*, 2020. 36(8): 2401-2409
- [81] Chang, C.Y., Hsu, T.W. and Chang, J.M. PSLCNN: Protein Subcellular Localization Prediction for Eukaryotes and Prokaryotes Using Deep Learning. In: *Proceedings of International Conference on Technologies and Applications of Artificial Intelligence, (TAAI)*, 2019. pp. 1-5.
- [82] Senior, A.W., Evans, R., Jumper, J., Kirkpatrick, J., Sifre, L., Green, T., Qin, C., Židek, A., Nelson, A.W., Bridgland, A. and Penedones, H. Protein Structure Prediction Using Multiple Deep Neural Networks in the 13th Critical Assessment of Protein Structure Prediction (CASP13). *Proteins: Structure, Function and Bioinformatics*, 2019.87(12):1141-1148.
- [83] Zheng, W., Li, Y., Zhang, C., Pearce, R., Mortuza, S.M. and Zhang, Y. Deep-Learning Contact-Map Guided Protein Structure Prediction in CASP13. *Proteins: Structure, Function and Bioinformatics*, 2019.87(12): 1149- 1164.
- [84] Tyanova, S., Temu, T. and Cox, J. The Max Quant Computational Platform for Mass Spectrometry-Based Shotgun Proteomics. *Nature Protocols*, 2016.11(12): 2301-2319.
- [85] Nesvizhskii, A.I., 2014. Proteogenomics: Concepts, Applications and Computational Strategies. *Nature Methods*, 11(11):1114-25.
- [86] Chen, L., Zhang, B., Schnaubelt, M., Shah, P., Aiyetan, P., Chan, D., Zhang, H. and Zhang, Z. MS-PyCloud: An Open-Source, Cloud Computing- Based Pipeline for LC-MS/MS Data Analysis. *bioRxiv*, 2018. [https://doi.org/ 10.1101/320887](https://doi.org/10.1101/320887).
- [87] Chiva, C., Olivella, R., Borràs, E., Espadas, G., Pastor, O., Solé, A. and Sabido, E. QCloud: A Cloud-Based Quality Control System for Mass Spectrometry-Based Proteomics Laboratories. *PLoS ONE*, 2018. 13(1): e0189209.
- [88] Feng, J., Ding, C., Qiu, N., Ni, X., Zhan, D., Liu, W., Xia, X., Li, P., Lu, B., Zhao, Q. and Nie, P. Firmiana: Towards a One-Stop Proteomic Cloud Platform for Data Processing and Analysis. *Nature Biotechnology*, 2017. 35(5): 409.
- [89] Muth, T., Peters, J., Blackburn, J., Rapp, E. and Martens, L. Proteocloud: A Full- Featured Open Source Proteomics Cloud Computing Pipeline. *Journal of Proteomics*, 2013. 88: 104-108.