

SELECTIVE BREEDING IN AQUACULTURE: CURRENT STATE OF KNOWLEDGE AND APPLICATIONS

Author

Aishwarya Sharma

Assistant Professor cum Jr. Scientist
College of Fisheries
Kishanganj, Bihar, India
aishwaryasharma974@gmail.com

I. INTRODUCTION

Aquaculture, which involves the cultivation of aquatic organisms in controlled environments, has become a crucial sector of food production [1]. This is particularly important due to concerns about the sustainability of ocean's fisheries, which are struggling to meet the increasing global demand for seafood [2]. Selective breeding has been a powerful tool in improving agricultural and livestock production for decades. The practice of selective breeding dates back to the experiments of Gregor Mendel, in the 19th century [3]. However, the adoption of selective breeding in aquaculture has been slower compared to its application in plant and terrestrial animal production [4].

Starting as early as 1919 with the work of Embury and Hyford selecting for increased survival to furunculosis in brook trout [5], and then in 1920 with the development of productive strains in common carp [6], these experiments paved the way for further research in selective breeding for desirable traits. Throughout the years, a range of selection experiments have been conducted to improve the genetic characteristics of different fish species. These studies aimed to enhance traits that are economically important in aquaculture, such as growth rate and disease resistance etc. The mentioned studies, conducted by researchers like Argue *et al.* [7]; Dunham [8]; Gitterle *et al.* [9]; Hetzel *et al.* [10]; Hussain *et al.* [11] and Langdon *et al.* [12], contributed to the knowledge surrounding selective breeding in aquaculture.

According to Mair [13] less than 20% or even less than 10%, according to Gjedrem [14] of world's aquaculture production comes from genetically improved stocks. This suggests that the majority of aquaculture production still relies on non-genetically improved populations. The reasons for this situation include insufficient information available about the reproductive cycles of many cultivated species [15]. Understanding the reproductive biology and behavior of these species is crucial for implementing effective genetic improvement programs. Many aquaculture programs still rely on capturing wild specimens for breeding purposes. This practice can hinder the domestication of the species and reliance on wild-caught individuals can introduce genetic variability that is not aligned with the goals of breeding programs. The lack of knowledge about quantitative genetics, breeding theories, and breeding programs is another hurdle. For successful genetic improvement, researchers, extensionists, and fish farmers need to understand the principles of genetics that govern traits

of interest. This knowledge gap can impede the development and implementation of effective breeding strategies.

By repeatedly selecting and breeding individuals with the desired characteristics, the frequency of those traits in the population increases over time. This approach is highly effective for improving traits like growth rate, disease resistance, and product quality [16]. Unlike some other genetic approaches (Figure 1) like hybridizations and cross-breeding, chromosome manipulation, sex control and transgenesis; the genetic improvements achieved through selective breeding can be passed down to subsequent generations without the need for continuous intervention. Selective breeding offers the advantage of scalability.




Figure 1: Breeding Techniques in Aquaculture

Thus, selective breeding is a cornerstone of genetic improvement in aquaculture. Its ability to achieve permanent, heritable gains in traits, its scalability, and its potential to reduce production costs and improve product quality make it an invaluable tool in the development of efficient and sustainable fish farming practices. Therefore, this book chapter aims to give some overview of selective breeding, certain breeding programs and its current status, outcomes of selective breeding in aquaculture species and its application to encompass the increasing demand of sourced proteins i.e., proteins from fishes for increasing human population.

II. SELECTIVE BREEDING PROGRAMS AND CURRENT STATUS IN AQUACULTURE

Several research studies have consolidated details related to the layout of aquatic species breeding schemes [17] and breeding strategies [15]. Initially, breeding programs employed the words individual selection and mass selection, regularly used interchangeably. These terms refer to selection methods based solely on an individual's phenotype and common strategy practiced in fish breeding due to its simplicity. This method doesn't require individual identification or the maintenance of pedigree records and leads to rapid improvement. However, there are potential risks associated with mass selection, particularly

the risk of inbreeding. Inbreeding can occur when the best offspring are consistently chosen from a small number of parents, especially if the progeny groups are large. This process can lead to a reduction in genetic diversity within the population, which can negatively impact the overall health and performance of the aquatic species being bred. An example is provided using *Nile Tilapia* (*Oreochromis niloticus*), where two generations of mass selection for growth rate showed no improvement over the original base population [18]. The lack of response to selection in his case was attributed to factors such as inbreeding and genetic drift, highlighting the potential limitations of simple and unstructured mass selection methods. To mitigate these issues, it is suggested that a larger number of parents be used in the breeding program, as indicated by studies by Gjerde *et al.* [19] and Villanueva *et al.* [20]. Structured breeding plans and more sophisticated selection strategies may be necessary to achieve significant and sustained improvements in aquatic species while minimizing the risks associated with inbreeding and genetic drift.

Another breeding programs use the within-family selection in improving *Tilapia* strain in SE Asian countries through selective breeding. The study was conducted by Uraiwan and Doyle in 1986 [21]. The selection program was carried out at the Freshwater Aquaculture Center (FAC) of Central Luzon State University in the Philippines. The basis for selection was the body weight of *Tilapia* at 16 weeks of age. The selection line began with a base population that consisted of four strains of *Tilapia*. Nineteen full-sib (siblings with the same parents) groups were established. To prevent inbreeding, the heaviest male from one family was mated with the two heaviest females from another family. After 12 generations of within-family selection, the genetic gain in body weight was estimated to be 12.4% per generation. This indicates a consistent improvement in body weight across generations. The study suggests that the selective breeding program resulted in a strain of *Tilapia* with improved performance characteristics. According to Camacho *et al.* [22], within-family selection was found to be easy to manage and rotational mating effectively addressed concerns related to inbreeding.

The combined selection refers to a breeding strategy that takes into account both individual performance data and the genetic information of related animals (such as full siblings, half-siblings, and progeny) to improve the accuracy of estimating the breeding values of individuals. This approach is valuable because it increases the accuracy of predicting an individual's breeding value, which is a key factor in selecting the best candidates for breeding programs. In the earlier work with fish mentioned [23], selection index theory was used to combine individual, full-sib, and half-sib information. A selection index is a mathematical tool that combines multiple sources of information, such as phenotypic data and genetic relationships, to predict the breeding value of an individual. However, while selection index theory can be effective, it has certain limitations that were addressed through the development of more advanced methodologies like Best Linear Unbiased Prediction (BLUP) [24]. One significant advantage of BLUP procedures is that they allow for the estimation of genetic gain by examining the mean of the estimated breeding values (EBVs) in each generation or year of selection. The availability of computer programs, such as PEST [25] and ASReml [26] has greatly facilitated the implementation of BLUP procedures. These software tools can handle large amounts of pedigree and phenotypic data, allowing for more accurate estimation of breeding values and better-informed selection decisions.

Several studies documented successful application of combined selection for fish improvement in developing countries as mentioned:

1. The GIFT Project (Genetically Improved Farmed Tilapia) in Philippines:

- **Species:** *Nile Tilapia (Oreochromis niloticus)*
- **Genetic Gains:** Reported genetic gains of 12-17% per generation over five generations.
- **Focus:** The main focus of selection was growth rate.
- **Reference:** Eknath *et al.*, 1998 [23].

2. The Jayanti Rohu Selective Breeding Project – India:

- **Species:** *Jayanti Rohu (Labeo rohita)*
- **Genetic Gains:** Reported a genetic gain of 17% per generation over five generations.
- **Focus:** The primary emphasis was on improving growth rate.
- **Reference:** Reddy *et al.*, 1999 [27]; Mahapatra, 2004 [28].

3. The *Oreochromis shiranus* Selection Project - Malawi:

- **Species:** *Malawian indigenous Tilapia (Oreochromis shiranus)*
- **Genetic Gains:** An accumulated gain of 13% over two generations.
- **Focus:** The selection project aimed to improve growth rates in this indigenous Tilapia species.
- **Reference:** Maluwa and Gjerde, 2007 [29].

Massive family-centered breeding programs, currently developed as the industry standard for the genetic advancement of aquaculture species, were first introduced for *salmonids* in the 1970s [30], *Nile tilapia* in 1988 [31], and marine prawn *P. vannamei* in 1993 [32]. After that, selection programs based on sib data have been utilized for a variety of species of aquaculture throughout the globe.

While there is significant potential for achieving high genetic gains in aquatic species through breeding programs, the development and implementation of these programs have been slow. A survey from 2003 (cited by Gjedrem and Baranski in 2009) indicated that there were only 60 family-based breeding programs in aquaculture, covering less than 5% of global aquaculture production at that time [33]. More recent surveys by Neira [34] and Rye *et al.* [35] suggest that the number of family-based selection programs in aquaculture has increased, surpassing 100 programs. It's also noted that there might be additional programs that have not been accounted for. The survey revealed that the highest number of breeding programs were for tilapia (27), followed by Atlantic salmon (13), and rainbow trout (13). However, among the widely farmed cyprinid species, only common carp (8) and rohu carp (1) were represented in the breeding programs. In 2010, it was estimated that around 8.2% of aquaculture production was based on genetically improved stocks. The exception was *Atlantic salmon*, where around 97% of the world's production used improved stock [33]. Considering potential gaps in the data, it's suggested that less than 10% of the global aquaculture production in

2010 was based on genetically improved stocks. The estimate might be optimistic, as some breeding programs could have been overlooked, and larger aquaculture companies might be implementing in-house selection programs that were not accounted for.

III. OUTCOMES OF SELECTIVE BREEDING IN AQUACULTURE SPECIES

The study by Schaperclaus in 1962 [36] from Germany demonstrated a significant selection response for reduced mortality due to furunculosis in common carp. These can be seen as one of the pioneering attempts at improving disease resistance through selective breeding in aquaculture. In 1920, researchers started developing two productive strains of common carp [37]. This marked the beginning of strain development in carp, an economically important species in aquaculture. Nemashev [38] and Kirpichnikov [39] reported that heritability estimates for weight in fingerlings of common carp show genetic influences on growth. The reported heritability values of 0.21 and 0.10 indicate that genetics play a moderate role in determining the weight of these young fish.

In *Brook Trout* (*Salvelinus fontinalis*), Embury and Hyford in 1925 selected surviving brook trout from endemic furunculosis population [40]. The Survival rate increased from 2% to 69% after three generations of selection. In *Rainbow Trout* (*Oncorhynchus mykiss*), Donaldson started breeding rainbow trout in 1932 and increased growth and fecundity was reported. Donaldson and Olson in 1955 [41] managed to develop the "Donaldson" strain of rainbow trout through directed individual selection. This strain is highly appreciated by trout culturists not only in the USA but also globally. Despite the success of this pioneering work, there was relatively little research conducted in the field of selective breeding for aquaculture species before 1970. In the mid-1960s, Moav and Wohlfarth [42] initiated a mass selection program in common carp. The primary objective of this program (started in 1965) was to improve the growth rate of the carp through selective breeding. However, despite their efforts, this program did not yield any significant improvements in the growth rate of the carp. Another breeding program initiated during the same period, with the goal of developing common carp that were resistant to dropsy, a serious infectious disease. This program was led by Kirpichnikov [43] and was conducted over the long term. The breeding strategy involved selection within local and Siberian wild carp populations from the river Amur, as well as crossbreeding between them. This selection effort proved successful, as resistance to dropsy was improved through selective breeding. As a result of this successful program, three distinct stocks of Krasnodar common carp were developed. Additionally, crossbreeding between the different stocks of Krasnodar common carp was carried out, resulting in heterotic crossbreds. These crossbreds showed a combination of the favorable traits from the parent stocks and were well-suited for commercial production. The commercial use of these heterotic crossbreds extended beyond the Krasnodar region to other parts of the former Soviet Union, indicating the broader impact and success of the breeding program in improving the characteristics of common carp for farming.

With *Atlantic Salmon* and *Rainbow Trout* in Norway, AKVAFORSK started breeding experiments in 1971, later transformed into a national breeding program. Traits considered for selection included growth rate, low frequency of early maturation, disease resistance, and flesh quality, color and fat percentages. Genetic gains of 13% for *salmon* and 14.4% for *trout* were reported by 1985. The Philippines National Tilapia Breeding Programme was started in 1993 with broodstock from the GIFT (Genetic Improvement of Farmed Tilapia) programme.

This resulted in significant growth improvement in GIFT *Nile Tilapia* through selective breeding. The program achieved a growth improvement ranging from 77% to 123%, which was considered superior to the results obtained from crossbreeding experiments. The program managed to achieve an impressive 11 percent genetic gain per generation in GIFT *tilapia*. This rate of genetic improvement is noteworthy, especially when compared to genetic gains obtained in other species of fish under similar breeding programs.

Collaborative research project between CIFA (India) and Institute of Aquaculture Research, (AKVAFORSK, Norway) was initiated in 1992 for Rohu (*Labeo rohita*) improvement. The fingerlings (young fish) from different Rohu populations were brought to CIFA, where they served as the base populations for the breeding program. Over time, the researchers likely selected individuals with the most desirable traits and bred them to create successive generations of fish with improved genetic characteristics for farming. Numerous selection studies and breeding schemes intended for improving rate of growth have lately been accomplished across multiple species, leading to 10-20% gain in every generation. Table 1 lists the species and nations participating in selective breeding.

Table 1: The Selection Response for Growth Rate in Aquatic Species under Selective Breeding

Species	Countries	Genetic Gain% per Generation	No. of generation	References
Channel Catfish (<i>Ictalurus punctatus</i>)	USA	13%	6	[8]
Coho Salmon (<i>Oncorhynchus kisutch</i>)	Chile	15%	4	[44]
Atlantic Salmon (<i>Salmo salar</i>)	Chile, Iceland, Ireland & Scotland	14%	6	[45]
Rainbow trout (<i>Oncorhynchus mykiss</i>)	Chile, France & USA	13%	2	[45, 46]
Common Carp (<i>Cyprinus carpio</i>)	Israel	-	-	[42]
Indian rohu carp (<i>Labeo rohita</i>)	India	29.6%	4	[28]
Silver barb (<i>Barbodes gonionotus</i>)	Bangladesh	12%	3	[11]
Tilapia (<i>Oreochromis niloticus</i>)	Philippines	12%	12	[47]

Tilapia (<i>Oreochromis niloticus</i>)	Egypt & Kenya	17%	5	[48]
Sea Bream (<i>Sparus aurata</i>)	Greece	22%	1	[40]
Shrimp (<i>Penaeus monodon</i>)	Colombia	4.2%	4-5	[9]
Oysters (<i>Crassostrea gigas</i>)	Asia, Europe & America	9.5%	1	[12]
Calico Scallop (<i>Argopecten ventricosus</i>)	Mexico	16%	1	[50]

IV. APPLICATIONS OF SELECTIVE BREEDING IN THE AQUACULTURE INDUSTRY

The role of selective breeding in the aquaculture industry, its potential benefits and challenges associated with it, and how genetic data and manipulation techniques can be employed to improve aquaculture species has been reported. The points below discuss the applications of selective breeding in the aquaculture industry and how it can contribute to the improvement of aquatic species for better husbandry and production efficiency:

1. **Selective Breeding as a Tool:** Selective breeding involves the deliberate selection of individuals with desirable traits to be used as parents for the next generation, thus increasing the occurrence of those traits over time. This process is aimed at improving the genetic makeup of aquatic species for aquaculture purposes.
2. **Human Intervention vs. Natural Selection:** The selective breeding involves human intervention in the adaptation process, which substitutes natural selection with controlled breeding practices. This alteration of the natural evolutionary process can lead to both positive and negative outcomes for aquaculture, depending on the chosen traits.
3. **Genetic Considerations and Well-Designed Programs:** Incorporating genetic considerations and well-designed selective breeding programs into aquaculture efforts is suggested to increase the likelihood of success. This involves careful planning and execution of breeding strategies to achieve desired traits.
4. **Academic vs. Industry Approach:** While academic institutions have contributed to genetic data collection and research, incorporating selective breeding programs within the aquaculture industry could be more efficient and aligned with industry goals.
5. **Adapting to Market Changes:** Selective breeding is portrayed as a means to help the aquaculture industry adapt to market dynamics and evolving demands by producing improved stocks with desired traits.

6. **Ongoing Evolution of Aquaculture:** The aquaculture industry is still developing and evolving, and selective breeding is one of the mechanisms through which it can progress toward defined production goals.
7. **Challenges and Potential:** The aquaculture industry needs more genetic data for various species to formulate effective breeding programs. While some species like rainbow trout, Atlantic salmon, channel catfish, and carp have adequate genetic data for specific traits, there's a need for more comprehensive data collection for other important aquacultural species. The Norwegian Atlantic salmon industry is cited as an example of successful industry-driven breeding programs. The aquaculture industry is becoming more competitive, requiring the development of efficient breeding programs to produce selectively bred stocks that enhance production efficiency. The advancement of genetic engineering techniques, such as chromosome manipulation and introduction of novel genes, complements selective breeding efforts and offers exciting potential for genetic improvement.
8. **Dynamic Nature of Aquaculture:** Selective breeding is viewed as a mechanism that allows the aquaculture industry to advance toward defined production goals and adapt to market demands and changes over time.

Overall, the points above emphasizes the importance of well-planned selective breeding programs, genetic data collection, and the potential integration of genetic engineering techniques to drive advancements in the aquaculture industry. Similar to agriculture, selective breeding is viewed as a major mechanism for advancing the aquaculture industry. As the market and industry demands evolve, selective breeding can help producers adapt to changes and achieve specific production goals.

REFERENCES

- [1] Gjedrem, T. (2012). Genetic improvement for the development of efficient global aquaculture: A personal opinion review. *Aquaculture*, 344–349, 12–22. <https://doi.org/10.1016/j.aquaculture.2012.03.003>.
- [2] Brander, K. M. (2007). Global fish production and climate change. *Proceedings of the National Academy of Sciences*, 104(50), 19709–19714. <https://doi.org/10.1073/pnas.0702059104>.
- [3] Mendel, G. J. (1866). Versuche über Pflanzenhybriden. *Verhandlungen Naturforschung Vereins Brünn*, 4, 3–47. <https://www.biodiversitylibrary.org/part/175272>.
- [4] Hagedoorn, A. L. (1950) Animal Breeding. Crosby Lockwood & Son Ltd., London. <https://www.nature.com/articles/hdy195142.pdf>.
- [5] Embury, G. C., & Hayford, C. O. (1925). The Advantage of Rearing Brook Trout Fingerlings from Selected Breeders. *Transactions of the American Fisheries Society*, 55(1), 135–148. [https://doi.org/10.1577/1548-8659\(1925\)55\[135:TAORBT\]2.0.CO;2](https://doi.org/10.1577/1548-8659(1925)55[135:TAORBT]2.0.CO;2).
- [6] Gjerde, B. & Korsvoll, A. (1999). Realized selection differentials for growth rate and early sexual maturity in Atlantic salmon. Abstracts, *Aquaculture Europe 99, Trondheim, Norway*, August 7–10, 73–74. <https://link.springer.com/book/10.1007/978-90-481-2773-3>.
- [7] Argue, B. J., Arce, S. M., Lotz, J. M., & Moss, S. M. (2002). Selective breeding of Pacific white shrimp (*Litopenaeus vannamei*) for growth and resistance to Taura Syndrome Virus. *Aquaculture*, 204(3–4), 447–460. [https://doi.org/10.1016/S0044-8486\(01\)00830-4](https://doi.org/10.1016/S0044-8486(01)00830-4).
- [8] Dunham, R. A. (2006). Comparison of six generations of selection, interspecific hybridization, intraspecific crossbreeding and gene transfer for growth improvement in *Ictalurus catfish*. *AGA*, 26–30 June, Montpellier, Abstract: 22. https://eja.journals.ekb.eg/article_32049_4a776074f94358c92f67f9f37ca71a2f.pdf
- [9] Gitterle, T., Salte, R., Gjerde, B., Cock, J., Johansen, H., Salazar, M., Lozano, C., & Rye, M. (2005). Genetic (co)variation in resistance to White Spot Syndrome Virus (WSSV) and harvest weight in *Penaeus*

- (Litopenaeus) vannamei. *Aquaculture*, 246(1–4), 139–149. <https://doi.org/10.1016/j.aquaculture.2005.02.011>
- [10] Hetzel, D. J. S., Crocos, P. J., Davis, G. P., Moore, S. S., & Preston, N. C. (2000). Response to selection and heritability for growth in the Kuruma prawn, *Penaeus japonicus*. *Aquaculture*, 181(3–4), 215–223. [https://doi.org/10.1016/S0044-8486\(99\)00237-9](https://doi.org/10.1016/S0044-8486(99)00237-9).
- [11] Hussain, M. G., Islam, M. S., Hossain, M. A., Wahid, M. I., Kohinoor, A. H. M., Dey, M. M., & Mazid, M. A. (2002). Stock improvement of silver barb (*Barbodes gonionotus* Bleeker) through several generations of genetic selection. *Aquaculture*, 204(3–4), 469–480. [https://doi.org/10.1016/S0044-8486\(01\)00832-8](https://doi.org/10.1016/S0044-8486(01)00832-8).
- [12] Langdon, C., Evans, F., Jacobson, D., & Blouin, M. (2003). Yields of cultured Pacific oysters *Crassostrea gigas* Thunberg improved after one generation of selection. *Aquaculture*, 220(1–4), 227–244. [https://doi.org/10.1016/S0044-8486\(02\)00621-X](https://doi.org/10.1016/S0044-8486(02)00621-X).
- [13] Mair, G. C. (2007). Genetics and breeding in seed supply for inland aquaculture. In: Bondad reantaso, m.g (ed.) Assessment of freshwater fish seed resources for sustainable aquaculture. *FAO fisheries technical paper: Rome*, pp: 519–547. <https://www.fao.org/3/a1495e/a1495e26.pdf>.
- [14] Gjedrem, T. (2012). Genetic improvement for the development of efficient global aquaculture: A personal opinion review. *Aquaculture*, 344–349, 12–22. <https://doi.org/10.1016/j.aquaculture.2012.03.003>
- [15] Gjedrem, T., Thodesen, J. (2005). Selection and breeding programs in aquaculture. Springer Berlin Heidelberg, p: 364. <https://download.e-bookshelf.de/download/0000/0032/90/L-G-0000003290-0002333212.pdf>.
- [16] Gjedrem, T. (1997). Selective breeding to improve aquaculture production. *World Aquaculture*, 28, 33–46. https://scholar.google.com/citations?view_op=view_citation&hl=en&user=W1v1W64AAAAJ&citation_f_or_view=W1v1W64AAAAJ:Y0pCki6q_DkC.
- [17] Gjerde, B. (2005). Design of Breeding Programs. In: Gjedrem, T., Ed., Selection and Breeding Programs in Aquaculture, Springer, Berlin, Heidelberg, 364. [https://www.scrip.org/\(S\(351jmbntvnsjt1aadkposzje\)\)/reference/referencespapers.aspx?referenceid=1323686](https://www.scrip.org/(S(351jmbntvnsjt1aadkposzje))/reference/referencespapers.aspx?referenceid=1323686).
- [18] Hulata, G., Wohlfarth, G. W., & Halevy, A. (1986). Mass selection for growth rate in the Nile tilapia (*Oreochromis niloticus*). *Aquaculture*, 57(1–4), 177–184. [https://doi.org/10.1016/0044-8486\(86\)90195-X](https://doi.org/10.1016/0044-8486(86)90195-X).
- [19] Gjerde, B., GjØen, H. M., & Villanueva, B. (1996). Optimum designs for fish breeding programmes with constrained inbreeding Mass selection for a normally distributed trait. *Livestock Production Science*, 47(1), 59–72. [https://doi.org/10.1016/S0301-6226\(96\)01000-7](https://doi.org/10.1016/S0301-6226(96)01000-7)
- [20] Villanueva, B., Woolliams, J. A., & Gjerde, B. (1996). Optimum designs for breeding programmes under mass selection with an application in fish breeding. *Animal Science*, 63(3), 563–576. <https://doi.org/10.1017/S1357729800015459>.
- [21] Uraivan, S., & Doyle, R. W. (1986). Replicate variance and the choice of selection procedures for tilapia (*Oreochromis niloticus*) stock improvement in Thailand. *Aquaculture*, 57(1–4), 93–98. [https://doi.org/10.1016/0044-8486\(86\)90185-7](https://doi.org/10.1016/0044-8486(86)90185-7).
- [22] Camacho, A. S., Abella, T., Tayamen, M. M. (2001). Fish genetics research and development in the Philippines. In: Gupta MV, Acosta BO (eds), Fish Genetics Research in Member Countries and Institutions of the International Network on Genetics in Aquaculture. *ICLARM Conf. Proc. 64*, Penang, Malaysia, pp. 71–76. <http://pubs.iclarm.net/libinfo/Pdf/Pub%20CP6%2064.pdf>.
- [23] Eknath, A. E., Dey, M. M., Rye, M., Gjerde, B., Abella, T. A., Sevilleja, R., Tayamen, M. M., Reyes, R. A., Bentsen, H. B. (1998). Selective breeding of Nile tilapia for Asia. 6th World Congress on Genetics Applied to Livestock Production, Vol. 27. *University of New England, Armidale, Australia*, pp. 89–96. https://www.researchgate.net/publication/279537869_Selective_breeding_of_Nile_tilapia_for_Asia
- [24] Van Vleck, L. D. (1993). Selection Index and Introduction to Mixed Model Methods. *CRC Press Inc, Boca Raton, FL, USA*. <https://lib.ugent.be/catalog/rug01:000456254>.
- [25] Groeneveld, E. (1990). PEST User's Manual. *Dept of Animal Science, University of Illinois, USA: Urbana, Illinois*. 61801, 1990.
- [26] Gilmour, A. R., Gogel, B. J., Cullis, B. R., Welham, S. J., Thompson, R. (2009). ASReml User Guide Release 1.0. VSN International Ltd, Hemel Hempstead, UK. https://www.researchgate.net/publication/228746364_Asrem1_User_Guide_Release_30.
- [27] Reddy, P. V. G. K., Gjerde, B., Mahapatra, K. D., Jana, R. K., Saha, J. N., Rye, M., Meher, P. K. (1999). Selective Breeding Procedures for Asian Carps. Central Institute of Freshwater Aquaculture, Bhubaneswar, India.

- [28] Mahapatra, K. D., Gjerde, B., Saha, J. N., Reddy, P. V. G. K., Jana, R. K., Sahoo, M. and Rye, M. (2004). Realized genetic gain for growth in rohu (*Labeo rohita*). In manuscript.
- [29] Maluwa, A. O., & Gjerde, B. (2007). Response to selection for harvest body weight of *Oreochromis shiranus*. *Aquaculture*, 273(1), 33–41. <https://doi.org/10.1016/j.aquaculture.2007.06.014>
- [30] Gjedrem, T. (1985). Improvement of productivity through breeding schemes. *GeoJournal*, 10(3), 233–241. <https://doi.org/10.1007/BF00462124>
- [31] Eknath, A. E., Bentsen, H. B., Gjerde, B., Tayamen, M. M., Abella, T. A., Gjedrem, T., Pullin, R. V. S. (1991). Approaches to national fish breeding programs: pointers from tilapia pilot study. *NAGA, The ICLARM Quarterly* No. 723, pp. 10–12. <https://ideas.repec.org/a/wfi/wfnaga/26826.html>
- [32] Fjalestad, K. T., Gjedrem, T., Carr, W. H., Sweeney, J. N. (1997). Final report: the shrimp breeding program. Selective breeding of *Penaeus vannamei*. *AKVAFORSK, Report no. 17/97*, p. 85. https://scholar.google.com/scholar_lookup?title=Final%20report%3A%20the%20shrimp%20breeding%20program.%20Selective%20breeding%20of%20Penaeus%20vannamei&publication_year=1997&author=Fjalestad%2CK%20T&author=Gjedrem%2CT&author=Carr%2CW%20H
- [33] Gjedrem, T., & Baranski, M. (2009). *Selective breeding in aquaculture: An introduction*. Springer. <https://link.springer.com/book/10.1007/978-90-481-2773-3>.
- [34] Neira, R. (2010). Breeding in aquaculture species: genetic improvement programs in developing countries. *9th World Congress on Genetics Applied to Livestock Production*, Leipzig, Germany, August 1–6, p. 8. https://www.researchgate.net/publication/265313094_Breeding_in_Aquaculture_Species_Genetic_Improvement_Programs_in_Developing_Countries.
- [35] Rye, M., Gjerde, B., Gjedrem, T. (2010). Genetic development programs for aquaculture species in developed countries. *9th World Congress on Genetics Applied to Livestock production*, Leipzig, Germany, August 1–6, p. 8.
- [36] Schaperclaus, W. (1962). *Trate de Pisciculture en Etang. Vigot Freres, Paris*, 208-227. <https://www.scirp.org/journal/paperinformation.aspx?paperid=51005>.
- [37] Kirpichnikov, V. S. (1987). Selection and New Breeds of Pond Fishes in the USSR. In: Tiewes, K. (Ed.), *Selection, Hybridization and Genetic Engineering in Aquaculture*, Volume II, 461-473.
- [38] Nemashev, G. A. (1969). Heritability of Several Traits Important in the Common Carp Selection. *Izvestija Gosudarstvennogo Nauchno-issledovatel'skogo, Instituta Ozerogo i Rechnogo Rybnogo Khosjaistva (GosNIORKh)*, Vol. 65, 185-195. <https://www.scirp.org/journal/paperinformation.aspx?paperid=51005>.
- [39] Kirpichnikov, V. S. (1972). Theory of Fish Selection. In: Cherfas, B.I., Ed., *Genetics. Selection and Hybridization of Fish, Academy of Sciences of the USSR, Israel Program for Scientific Translations, Jerusalem*, 299. https://scholar.google.com/scholar_lookup?hl=en&publication_year=1972&author=V.S.+Kirpichnikov&title=Genetics%2C+Selection+and+Hybridization+of+Fish
- [40] Embury, G. C., & Hayford, C. O. (1925). The Advantage of Rearing Brook Trout Fingerlings from Selected Breeders. *Transactions of the American Fisheries Society*, 55(1), 135–148. [https://doi.org/10.1577/1548-8659\(1925\)55\[135:TAORBT\]2.0.CO;2](https://doi.org/10.1577/1548-8659(1925)55[135:TAORBT]2.0.CO;2).
- [41] Donaldson, L. R., & Olson, P. R. (1957). Development of Rainbow Trout Brood Stock by Selective Breeding. *Transactions of the American Fisheries Society*, 85(1), 93–101. [https://doi.org/10.1577/1548-8659\(1955\)85\[93:DORTBS\]2.0.CO;2](https://doi.org/10.1577/1548-8659(1955)85[93:DORTBS]2.0.CO;2)
- [42] Moav, R., & Wohlfarth, G. (1976). TWO-WAY SELECTION FOR GROWTH RATE IN THE COMMON CARP (*CYPRINUS CARPIO* L.). *Genetics*, 82(1), 83–101. <https://doi.org/10.1093/genetics/82.1.83>.
- [43] Kirpichnikov, V. S., Ilyasov, Ju. I., Shart, L. A., Vikhman, A. A., Ganchenko, M. V., Ostashevsky, A. L., Simonov, V. M., Tikhonov, G. F., & Tjurin, V. V. (1993). Selection of Krasnodar common carp (*Cyprinus carpio* L.) for resistance to dropsy: Principal results and prospects. *Aquaculture*, 111(1–4), 7–20. [https://doi.org/10.1016/0044-8486\(93\)90020-Y](https://doi.org/10.1016/0044-8486(93)90020-Y).
- [44] Hershberger, W. K., Myers, J. M., Iwamoto, R. N., Mcauley, W. C., & Saxton, A. M. (1990). Genetic changes in the growth of coho salmon (*Oncorhynchus kisutch*) in marine net-pens, produced by ten years of selection. *Aquaculture*, 85(1–4), 187–197. [https://doi.org/10.1016/0044-8486\(90\)90018-I](https://doi.org/10.1016/0044-8486(90)90018-I).
- [45] Gjerde, B. (1986). Growth and reproduction in fish and shellfish. *Aquaculture*, 57(1–4), 37–55. [https://doi.org/10.1016/0044-8486\(86\)90179-1](https://doi.org/10.1016/0044-8486(86)90179-1)
- [46] Kincaid, H. L., Bridges, W. R. and Limbach, B. (1977). Three generations of selection for growth rate in fall spawning rainbow trout. *Transaction of the American Fisheries Society*, 106, 621–629. <https://www.usgs.gov/publications/three-generations-selection-growth-rate-fall-spawning-rainbow-trout>.

- [47] Bolivar, R. (1999). Estimation of response to within-family selection for growth rate in tilapia. *Diss. Abst. Int. Pt. B-Sci. and Eng.* 60(3): 934. <https://dalspace.library.dal.ca/xmlui/handle/10222/55584>.
- [48] Eknath, A. E., Tayamen, M. M., Palada-de Vera, M. S., Danting, J. C., Reyes, R. A., Dionisio, E. E., Capili, J. B., Bolivar, H. L., Abella, T. A., Circa, A. V., Bentsen, H. B., Gjerde, B., Gjedrem, T., & Pullin, R. S. V. (1993). Genetic improvement of farmed tilapias: The growth performance of eight strains of *Oreochromis niloticus* tested in different farm environments. *Aquaculture*, 111(1-4), 171-188. [https://doi.org/10.1016/0044-8486\(93\)90035-W](https://doi.org/10.1016/0044-8486(93)90035-W).
- [49] Thorland, I., Papaioannou, N., Kottaras, L., Refstie, T., Pappasolomontos, S., & Rye, M. (2007). Family based selection for production traits in gilthead seabream (*Sparus aurata*) and European sea bass (*Dicentrarchus labrax*) in Greece. *Aquaculture*, 272, S314. <https://doi.org/10.1016/j.aquaculture.2007.07.193>.
- [50] Ibarra, A. M., Ramirez, J. L., Ruiz, C. A., Cruz, P., & Avila, S. (1999). Realized heritabilities and genetic correlation after dual selection for total weight and shell width in catarina scallop (*Argopecten ventricosus*). *Aquaculture*, 175(3-4), 227-241. [https://doi.org/10.1016/S0044-8486\(99\)00100-3](https://doi.org/10.1016/S0044-8486(99)00100-3).