

BIOTECHNOLOGICAL APPROACH FOR BIOREMEDIATION OF TEXTILE AND PESTICIDES INDUSTRIAL POLLUTANTS

Authors

Pratiksha G. Agale

Department of Zoology and Research
Centre
Shri Shiv Chhatrapati College of Arts
Commerce and Science
Junnar, Pune, Savitribai Phule Pune
University, India.

Varsha R. Dhole

Department of Zoology and Research
Centre
Shri Shiv Chhatrapati College of Arts
Commerce and Science
Junnar, Pune, Savitribai Phule Pune
University, India.

Pramod C. Mane

Department of Zoology and Research
Centre
Shri Shiv Chhatrapati College of Arts
Commerce and Science
Junnar, Pune, Savitribai Phule Pune
University, India.

Deepali D. Kadam

Department of Zoology and Research
Centre
Shri Shiv Chhatrapati College of Arts
Commerce and Science
Junnar, Pune, Savitribai Phule Pune
University, India.

Ravindra D. Chaudhari

Department of Zoology and Research
Centre
Shri Shiv Chhatrapati College of Arts
Commerce and Science
Junnar, Pune, Savitribai Phule Pune
University, India.
rdchaudhari2004@yahoo.co.in

Water has a key role in mediating ecosystem processes at the global scale, connecting the atmosphere, lithosphere, and biosphere by transporting materials across them and facilitating chemical reactions. Natural waters are never completely pure; instead, they are a complex, dynamic blend of suspended particles, dissolved inorganic and organic compounds.

On average, water accounts for 60 to 70 percent of an organism's weight. It fills cells, giving many tissues shape and support. All of life's chemical reactions take place in the medium of water, and water actively participates in many of these events. Water is a solvent that breaks down both the nutrients that cells require for survival and the waste products that cells generate. Water is therefore necessary for the transportation of materials to and from cells. Salts and other substances are dissolved by water, creating solutions that conduct electricity. The energy that powers photosynthesis is also provided by these fluids, which are known as electrolytes.

To prevent communicable diseases and maintain a healthy lifestyle, clean drinking water and basic sanitation are essential. The ongoing use of dirty water remains one of the biggest environmental dangers to health for many of the world's poorest populations. In 2002, the UN projected that 2.4 billion people lacked access to proper sanitation and at least 1.1 billion did not have access to safe drinking water. More than 5 million people every year die as a result of these deficits, which cause hundreds of millions of cases of water-related sickness.

Water shortages are anticipated to worsen as populations increase, more people migrate into cities, and agriculture and industry struggle for dwindling water supplies. Two-thirds of the world's population will reside in water-stressed nations by 2025, as determined by the United Nations as those whose freshwater supplies are consumed at a rate greater than. Reducing the number of people without dependable access to clean water and better sanitation by half was one of the top targets outlined at the UN World Summit in Johannesburg in 2002.

The minimal essential water demands of all of their population cannot be met by 45 countries, the most of which are in Africa or the Middle East. The issue of access to clean water exists in various nations. Accessibility does not necessarily equate to affordability. For instance, a typical low-income household in Lima, Peru, uses one-sixth the amount of water that a middle-class American family does while paying three times as much for it. To buy and purify water, a poor household could spend up to one-third of their income if they followed the government's advice to boil all water to prevent cholera.

Over the past ten years, investments in rural development have resulted in notable advances. Almost 800 million people, or 13% of the world's population, now have access to clean water since 1990. The proportion of rural families that have access to clean water has increased from less than 10% to around 75%.

I. BIOREMEDIATION

Composting and wastewater treatment are well-known examples of traditional environmental biotechnologies. Environmental biotechnology is not a new field. A biotechnological procedure called bioremediation reduces or eliminates environmental contamination. 'Bio' in bioremediation refers to a living thing, and 'remediate' means to address an issue. It is a form of waste management strategy that employs living things to either remove or utilize pollutants from a polluted region.

Food, energy, and other necessities of daily life are in greater demand due to the growing global human population. These demands were met by the Industrial Revolution, which led to the mass manufacture of several organic and inorganic compounds. These procedures cause environmental pollution in a variety of ways, whether directly or indirectly. Many various procedures are employed to lessen pollution, one of which is bioremediation, in which noxious chemicals or pollutants with low toxicity are neutralized by biological agents.

Recently, it was thought that bioremediation was a solution for problems with hazardous contaminants that were emerging and involving a variety of microbes, including

both aerobic and anaerobic bacteria, fungi, algae, and both. Living things undergo a reaction as a part of their metabolic processes to change contaminants in this process (Kensa *et al.*, 2011). In this procedure, naturally occurring bacteria and fungi are occasionally employed to detoxify or breakdown chemicals that are harmful to the environment or human health. The microorganisms may be isolated from another location and delivered to the contaminated locations, or they may be native to the contaminated area.

II. BIOREMEDIATION APPLICATIONS

- 1 It is an easy and labor-efficient method.
- 2 It is a natural process that takes some time when bacteria multiply and break down contaminants, but when contaminants are broken down, the population of microbes also declines.
- 3 Sustainable and eco-friendly.
- 4 By converting harmful to safe compounds, bioremediation is useful for the total eradication of a wide range of pollutants.
- 5 Pollutants can be destroyed rather than being transferred from one environmental medium to another, such as from land to water or air.
- 6 Bioremediation can be done locally, eliminating the need to carry trash elsewhere and reducing risks to both human health and the environment while doing so.
- 7 Compared to other approaches for hazardous waste cleanup, bioremediation is less expensive.
- 8 Bioremediation preserves aesthetic qualities by keeping industry out of the environment.
- 9 Contaminants are eliminated, not just distributed throughout various environmental media.
- 10 Comparably simple implementation.
- 11 Non-intrusive, possibly enabling continuous site use.

III. TYPES OF BIOREMEDIATION MICROBIAL BIOREMEDIATION

For the removal of harmful pollutants, bacteria and fungi are used as microorganisms. When a dangerous substance is present and the temperature is below zero, microbes can proliferate. The key contributing elements for the degradation of pollutants are the microbial population, the accessibility of contaminants to the microbial population, and environmental conditions such soil type, pH, temperature, oxygen content, and nutrition levels (Sharma, 2020).

1. **Phytoremediation:** Green plants and the related microorganism are used in this procedure to purge harmful environmental contaminants from the environment. A number of processes, including phytodegradation, phytovolatilization, phytoaccumulation, and phytoextraction, are used in phytoremediation. The health and yield of soil can be improved via phytoremediation, which is more affordable than other traditional methods (Singh *et al.*, 2017).
2. **Mycoremediation:** In terms of mycoremediation, the method of employing fungi to degrade harmful compounds in the environment. Fungi have non-specific enzymes that can break down a wide variety of substances. 'White rot fungi' is the mycoremediation branch that has seen the highest development (Tomer *et al.*, 2021).

3. **Bio- Stimulation:** The ability to send the stimulus to the environment is known as bio-stimulation. One of the most established methods of bio-remediation of hydrocarbons is bio-stimulation, which has lately made advancements in geophysics, stable isotope studies, and molecular microbiology. By first providing fertilizers, growth aids, and trace minerals, then by providing other environmental factors like pH, specific nutrients are injected at the site (soil/ground water) to stimulate the activity of native microorganisms, which include naturally existing bacteria and fungus communities. Secondly, oxygen and temperature to boost their metabolism. Pollutants that are present in modest amounts also act as stimulants by activating the bioremediation enzymes. The majority of the time, nutrients and oxygen assist these routes to continue by supporting local microbes (Kensa et al., 2011).
4. **Bio-Attenuation:** The contaminants are changed to a less hazardous form during bioattenuation. These transformational processes are mostly brought on by biodegradation by microorganisms, to some extent by reactions with naturally occurring chemicals, and to some amount by sorption on geologic media. Natural attenuation is a method for treating fuel compounds that is specifically acknowledged as polluted, but not for many other groups.

Many polluted locations might not need an aggressive repair strategy. According to Maitra (2018), bioattenuation is effective and economical. Bioattenuation is dependent on natural degradation processes. In order to make sure that the concentration of contaminants at important sampling points declines over time, a technique of tracking the natural progression of degradation has been developed (Sharma *et al.*, 2020).

5. **Bio-Pile:** Because of its cost-effectiveness, this ex-situ technique enables for the effective management of operational biodegradation variables like PH, Nutrients, Temperature, and Aeration. The usage of biopiles, sometimes referred to as bio-cells, bio-heaps, bio-compounds, and compost piles, helps to lower petroleum pollutant concentrations in excavated soils while promoting biodegradation. This method includes leachate collection, bed systems for treating leachate, nutrients, irrigation, and aeration.

It is also possible to remediate volatile low molecular weight contaminants with the biopile. In order to facilitate constant air circulation in contaminated piled soil through air pump, biopile systems were connected to additional field ex-situ bioremediation techniques, such as land farming, bioventing, biosparging, robust engineering, maintenance and operation cost, and lack of power supply at remote sites.

Extreme air temperatures can cause soil to dry out and undergo bioremediation, which inhibits microbial activity and promotes volatilization rather than biodegradation. The breakdown of adsorbed petroleum pollutants increased as a result of the increased microbial activity brought on by microbial respiration (Sharma *et al.*, 2020).

6. **Bioventing:** In order to increase the activities of native microbes for bioremediation, the bioventing technique involves controlled airflow stimulation by delivering oxygen to unsaturated zones. Amendments are made by adding nutrients and moisture to increase bioremediation, which achieves microbial transformation of pollutants to harmless state.

Bioventing is used for efficient bioremediation of petroleum-contaminated soil. In unsaturated soils, bioventing can considerably lower the concentration of a variety of hydrocarbons and other organic pollutants. Systems for bioventing remediation should be planned to reduce constituent volatilization. By eliminating the requirement for off-gas treatment, it lowers remediation costs (LEE *et al.*, 1993).

- 7. Bioremediation of Water Waste of Textile and Dye Industry Textile Dye:** Commercial synthetic dyes are also use in various types of industries such as paper, printing, plastic and pharmaceutical industry, different type of paint and textile industry. The textile industry plays important role in the economic development of different countries, as China is the largest exporter of textile products, followed by India, European Union, The USA and Turkey(Sudarshan *et al.*, 2023).
- 8. Textile Waste Toxicity:** Adverse effect of Textile dyes on human health such as, some dyes cause allergic skin reaction, Numerous respiratory tract irritations, skin and mucous membrane ulceration and mental disorientation when inhaled. Improperly disposed textile dye effluent affecting photosynthetic activity, so it increases heterotrophic activity, which result lowers dissolved oxygen levels affects water Ecosystem(Sudarshan *et al.*, 2023).

IV. TYPES

- 1. Bioremediation of textile water waste by Bacteria:** Most effective degraders of synthetic dye are Bacteria and Cyanobacteria, because of their short life cycles plays important role in secondary waste generation and adaptability to variety of substrates. Microorganism reduces hazardous chemicals and transform toxic chemicals into less harmful. Some bacterial strains, such as *Bacillus cereus*, *Pseudomonas putida*, *Pseudomonas fluorescence*, *Pseudomonas desmolyticum* and *Bacillus sp.* have been used in the biodegradation of azo dyes (Sudarshan *et al.*, 2023).

Table 1: Summary of decolorization of various dyes by pure and mixed bacterial Culture

Sr. No.	Dye	Bacteria	Decolorization of textile dye (in %)	References
1.	Methyl Red	<i>Staphylococcus saprophyticus</i> AUCA SVE3	94 & 97% Decolorization within 24 & 48 hrs. Resp.	Hakimet <i>al.</i> , (2014)
2.	Reactive Violet 5	<i>Paracoccus sp.</i> GSM2	70% decolorization within 38 Hrs.	Bheemaraddiet <i>al.</i> , (2014)

3.	Acid Orange	<i>Bacillus megaterium</i> <i>PM582</i>	73% Decolorization within 38 hrs.	Shahet <i>al.</i> , (2014)
4.	Reactive Red 198	<i>Acinetobacte</i> <i>r baumannii</i>	96.20% removal after 72hrs.	Unnikrishnan <i>et</i> <i>al.</i> , (2018)
5.	Reactive Yellow 145	<i>Thiosphaera</i> <i>pantothropha</i> <i>ATCC 35512</i>	50% decolorization within 96 hrs.	Garget <i>al.</i> , (2020)
6.	Reactive Red HE8B	<i>Pseudomonas</i> <i>aeruginosa</i>	86% decolorization within 48 hrs.	Patelet <i>al.</i> , (2016)
7.	Reactive Black 5	<i>Aeromonas</i> <i>hydrophila</i>	76% decolorization within 24 hrs.	El Bouraieet <i>al.</i> , (2016)
8.	Reactive Red 120	<i>Shewanella</i> <i>haliotis</i>	99% decolorization in 2.5 hrs.	Birmoleet <i>al.</i> , (2019)
9.	Congo Red, Reactive Black 5	<i>Enterococcus</i> <i>faecalis</i> <i>R1107</i>	65.57% & 72.64 % decolorization Resp. within 48 hrs.	Wanget <i>al.</i> , (2022)
10.	Malachite Green	<i>Pandoraea</i> <i>pulmonicola</i>	85.2% decolorization	Chenet <i>al.</i> , (2009)
11.	Reactive Blue 59	<i>Bacillus</i> <i>odyssey</i> <i>SUK3</i> , <i>Morganella</i> <i>morganii</i> <i>SUK5</i> & <i>Proteus sp.</i> <i>SUK7</i>	100% decolorization within 60h, 30h. & 24h resp.	Patilet <i>al.</i> , (2008)
12.	Reactive Orange 16	<i>Pseudomonas</i> <i>sp.</i>	100% decolorization within 48 hrs.	Jadhavet <i>al.</i> , (2010)
13.	Reactive Green 19A	<i>Micrococcus</i> <i>glutamicus</i> <i>NCIM-</i> <i>2168</i>	100% decolorization within 42hrs.	Sarataleet <i>al.</i> , (2009)

14.	Direct Black 22	<i>Bacterial consortium</i>	100% decolorization within 12 hrs.	Mohana <i>et al.</i> , (2008)
15.	Metanil Yellow	<i>Bacillus sp. AK1 & Lysinibacillus sp. AK2</i>	100% decolorization within 27 hrs. & 12 hrs. Resp.	Anjaneya <i>et al.</i> , (2011)
16.	Methyl Red, Tartrazine, Ponceaus, Rea Red 35, Evans Blue, Acid Red 3R, Acid red, Methyl Orange, Reactive violet, Red AG	<i>Nesterenkonia lacusekhoensis EMLA3</i>	<90% decolorization within 72 -192 hrs.	Prabhakaret <i>al.</i> , (2022)
17.	Acid Black, Congo red, Acid red 27,	<i>Bacillus</i>	51.2%, 1.9%, 32.05%, 36.2%	Kesebir <i>et al.</i> , (2021)
	Reactive black, Methylene Blue	<i>licheniformis</i>	decolorization resp.	
18.	Congo red, brilliant blue & Bromophenol blue, Crystal violet	<i>Staphylococcus haemolyticus</i>	80% and 40% decolorization resp. within 3 hrs.	Li <i>et al.</i> , (2020)
19.	Malachite Green	<i>Dietzia sp.</i>	72.05% decolorization	Beraet <i>al.</i> , (2016)
20.	Amido Black 10B	<i>Chroococcus minutus</i>	55% decolorization	Parikh <i>et al.</i> , (2005)
21.	Reactive Dark blue	<i>Exiguobacterium sp.</i>	97% decolorization within 24 hrs.	Qu <i>et al.</i> , (2010)

V. UTILIZING MICROALGAE FOR TEXTILE WASTE WATER BIOREMEDIATION

If discharged without adequate treatment, waste water from the textile industry contains a variety of pollutants, the majority of which are dyes and have negative effects on aesthetics, eutrophication, a reduction in photosynthetic activity, and bioaccumulation of toxins in aquatic ecosystems.

A viable alternative to the current standard method of waste water treatment is the growing of microalgae in the textile dye effluent. The conventional treatment process by using microalgae for bioremediation of textile effluents provided valuable biomass that can be processed into bioproducts, biofuels, and bioenergy. The treatment using microalgae reduces color and nutrient load of textile effluent, which reduces numerous negative environmental impacts caused by its discharge into natural environment (Premarathe *et al.*, 2021).

Table 2: Summary of Some Recent Studies on Phycoremediation of Textile Dye Wastewater Using Microalgae

Sr. No	Textile dye	Decolorizing Microalgae	Decolorization Removal percentage	References
1.	Indigo Blue	Scenedesmus quadricauda ABU12	100% decolorization within 4 days	Chiaet <i>et al.</i> , (2014)
2.	Congo Red	Chlorella	98%	Mahalakshmi <i>et al.</i> , (2015)
		vulgaris	Decolorization	
3.	Direct Red 5B	Comamonas sp. UVS	100% decolorization	Jadhav <i>et al.</i> , (2008)
4.	Congo Red	Haematococcus sp.	98% Decolorization	Mahalakshmi <i>et al.</i> , (2015)
5.	Azo dyes	Nostoc muscourm	68% Decolorization in 6 days	Omar <i>et al.</i> , (2008)
6.	Methylene Blue & Malachite Green	Desmodesmus sp.	98.6% decolorization in 6 days	Bera <i>et al.</i> , (2016)
7.	Direct Red 31	Chlorella pyrenoidosa	80.12 % decolorization within 180 min	Behl <i>et al.</i> , (2019)

8.	Indigo Blue	<i>Chlorella vulgaris</i>	49.03 % decolorization within 24 hrs.	Revathi <i>et al.</i> , (2017)
9.	Disperse orange 2RL	<i>Scenedesmus obliquus</i>	98.14% Decolorization	Hamouda <i>et al.</i> , (2022)
10.	CI Reactive Red 66	<i>Shewanella algae B29</i>	91.04% Decolorization	Chaiebet <i>et al.</i> , (2008)
11.	Remazol Black 5, Reactive Blue	<i>Chlamydomon areinhardtii</i>	72.97% Decolorization	San <i>et al.</i> , (2015)
12.	Remazol Black B	<i>Phormidiuman imale</i>	99.96 % decolorization	Bayazit <i>et al.</i> , (2020)

VI. UTILIZING FUNGI FOR BIOREMEDIATION OF TEXTILE WASTE WATER

The biological method—which employs a variety of microorganisms and fungi—is thought to be the most efficient and least energy-intensive way to remove the majority of pollutants from water.

Industrial dyes are removed by fungus through an adsorption mechanism, however in some fungi, such as White Rot fungus, both adsorption and degradation can take place at the same time. The decolorization of textile colors using *Funaliatrogii* pellets, a white rot fungus. The dye concentration, amount of pellet, temperature, and media agitation all had a substantial impact on the decolorization activity.

White rot fungus, which can release ligninolytic enzymes that bind to non-specific substrates and then degrade a wide range of refractory compounds (i.e., pollutants including dyes), can deculturate dyes (Jebapriya *et al.*, 2013).

Table 3: Summary of Decolorization of Various Dyes by Fungi

Sr. No.	Species	Dye	Percentage Removal	References
1.	<i>Aspergillus versicolor</i>	Reactive Black 5	98% decolorization within 420 min	Huang <i>et al.</i> , (2016)
2.	<i>Pleurotus eryngii</i>	Reactive Black 5	93.57 % decolorization within 72 hrs.	Hadibarataet <i>al.</i> , (2013)
3.	<i>Funalia trogii</i>	Reactive Black 5	100% decolorization within 48 hrs.	Mazmanciet <i>al.</i> , (2005)

4.	<i>Pleurotus eryngii</i>	Methyl Orange	43% decolorization with 5 min treatment.	Akpinaret <i>et al.</i> , (2017)
5.	<i>Coriolopsis gallica</i>	Reactive Black 5	82% decolorization within 120min	Ben <i>et al.</i> , (2022)
6.	<i>Penicillium sp. QQ</i>	Reactive dark blue	97% decolorization within 24 hrs	Qu <i>et al.</i> , (2010)
7.	<i>Penicillium oxalicum</i>	Methylene Blue	99.17% decolorization within 6hrs	Mathur <i>et al.</i> , (2021)
8.	<i>Penicillium simplicissimum</i>	Reactive Black 5	92% decolorization	Muthukumaran <i>et al.</i> , (2017)
9.	<i>Penicillium chrysosporium</i>	Reactive lack 5 & Direct red 81	88% decolorization	Muthukumaran <i>et al.</i> , (2017)
10.	<i>Penicillium sp. YW01</i>	Malachite Green	98.23 % decolorization within 6 days	Yang <i>et al.</i> , (2011)
11.	<i>Umbelopsis isabellinna & Penicillium geastrivorous</i>	Reactive Black 5	100% decolorization within 16-48 hrs.	Yang <i>et al.</i> , (2003)
12.	<i>Aspergillus niger</i>	Cibacron Black	33% decolorization	Biyik <i>et al.</i> , (2012)
		W-NN	decolorization	
13.	<i>Cyathus bulleri</i>	Reactive Red 198, Reactive Orange	80 % decolorization	Chhabra <i>et al.</i> , (2008)

Various dyes like Malachite Green, Commercial Xanthene, Rhodamine B, Brilliant Green, Azo dyes, Metanil Yellow and Methyl Orange leads to Carcinogenic, Genotoxic, Mutagenic and Neurotoxic against human health and other living organism, also affect immune system and reproductive as well as respiratory system of living organism (Sudarshan *et al.*, 2023).

VII. PESTICIDE BIOREMEDIATION

In agriculture, the use of pesticides boosts agricultural output and lowers crop loss. Agricultural discharges of pesticides into water increase their toxicity and harm aquatic life (Singhal *et al.*, 2021). According to the FAO (2018), Asia uses 52.8% of the world's pesticides, followed by the USA (30.0%), Europe (13.7%), Africa (2.2%), and Oceania (1.3%).

Pesticide removal is influenced by two factors: the first is the biome's ideal conditions for survival and activity, and the second is the pesticide's chemical composition and factors related to organisms (microalgae), such as the quantity of suitable organisms, the biological substrate, the availability of water, oxygen tension and redox potential, surface bonding, the presence of substitute carbon substrates, and other electron acceptors. (Nieet *al.*,2020).

Pesticides have significantly increased crop yields from agriculture by helping to manage pests on a global scale, yet applying pesticides heavily to agricultural land has negative effects on the environment, the human body, and human health. For this reason, the development of a quick method of pesticide detoxification is particularly crucial. The detoxification of environmental pesticide residues is greatly aided by bioremediation. Pesticides can be detoxified or degraded by a variety of microorganisms, including fungi, bacteria, and algae.

The composition of pesticides in contaminated wastewater, treatment costs, and ease of use are the main factors influencing pesticide treatment methods. In order to construct treatment facilities that are intended to remove emerging pollutants like pesticides from wastewater, a thorough investigation of influent characteristics and the coupling of the best treatment technology are necessary. For the elimination of pesticide degradation in aqueous medium, physical, chemical, and biological approaches have been widely applied (Nieet *al.*, 2020).

VIII.EFFECTS OF PESTICIDES AND HEAVY METALS ON HUMAN HEALTH

Organophorus (op), Carbamate (CB), and OC pesticides are among the most harmful because they work by interfering with the nervous system's normal operation (Riodolfiet *al.*, 2014). These pesticides lead to plenty of hazardous effects on human, animals, plants and environment. Table 4 depicts examples of some pesticides with their adverse effects on human health.

Table 4: Harmful Effect of Pesticides on Human Health

Sr no.	Pesticides	Health effects	
1.	Aldrin	Nervous system effects. Probable carcinogen.	Green <i>etal.</i> , (2004)
2.	Dichlorodiphenyltrichloroethane (DDT)	Nervous system effects (tremors, seizures). Probable carcinogen	
3.	Chlordane	Nervous system, digestive system, liver effects, Headaches, irritability, confusion, weakness, vision problems, vomiting, stomach cramps, diarrhoea, and jaundice for lower doses.	
4.	Dieldrin	Nervous system effects.	

		Probable carcinogen. Uncontrolled muscle Movement.
5.	Heptachlor	Nervous system damage, liver and adrenal gland Damage, tremors.

IX. MICROALGAL BIOREMEDIATION OF CONTAMINATED BY PESTICIDES

Algae likely make up to 27% of the total microbial biomass in the soil, making them a significant part of the soil microflora. It is crucial for the nitrogen economy of soils and helps sustain soil fertility and oxygen generation. Algae increase BOD by fixing carbon dioxide (CO₂) and releasing oxygen (O₂) during photosynthesis. Algae are used as biofertilizers or soil conditioners. contribute significantly to the biomonitoring and regulation of organic pollutants in aquatic ecosystems (Nie *et al.*, 2020). There are several pesticide elimination mechanisms used in bioremediation, including bio adsorption, bioaccumulation, and biodegradation. 2020 (Nie *et al.*).

The method of bio adsorption is passive (Ardal *et al.*, 2014). According to a recent study (Mishaqa *et al.*, 2017), grown algae were able to remove 87-96% of a variety of pesticides from aqueous phase, including atrazine, simazine, molinate, isoproturn, carbofuran, propanil, dimethoate, metolachlor, pendimethalin, and pyriproxin.

According to Ardalet *al.* (2014), bioaccumulation is an active process that can be expressed by the bio-concentration factor (BCF). According to Wang *et al.* (2014), variations in the bioconcentration mechanism, bioavailability of chemicals, physical barriers, methods of determining the BCF, dissolved organic matter, metabolism, ionization of ionizable compounds, and environmental conditions have a significant impact on the values of the BCF. According to additional data, BCF values vary depending on the concentration. Additionally, pyrometryne BCF values at 2.5 (or 5.0) g/L concentrations were higher than those at 10.0 (or 12.5) g/L concentrations in green algae (Jinet *al.*, 2012).

Pesticides in the environment undergo biodegradation as a result of different enzymes' metabolism. Pesticide degradation is a multi-step process that involves enzyme metabolism. Steps include (i) activating pesticides without functional groups by cytochrome P450 through oxidation, reduction, and hydroxylation reactions to produce more hydrophilic, soluble, degradable, and less toxic compounds; (ii) transferring enzymes in the cytosol to pesticides that are activated functional groups forming conjugation with glutathione, glucose, and malonate; and (iii) Glutathione transporters are responsible for moving these conjugates into vacuoles (Ghasemi *et al.*, 2011; Kumar and Singh, 2017; Laura *et al.*, 2013; Mau *et al.*, 2017).

Studies have been done on the co-culture of microalgae and beneficial bacteria for pollutant removal. The ability of microalgae to produce oxygen for photosynthesis to support bacterial development and microalgae to use carbon dioxide produced by bacterial metabolism as,

Table 5: Summary of Bioremediation Potential of Pesticides by Algae

Sr. No.	Microorganism	Pesticides	Percentage of Removal	References
1.	<i>Streptomyces sp. ML</i> <i>Streptomyces sp. OV</i>	Sole carbon thiamethoxam Dichlorophenol	84% o 40%	Bouferach <i>et al.</i> , (2022)
2.	<i>Monoraphidium braunii</i>	Bisphenol	48%	Gattullaet <i>al.</i> , (2012)
3.	<i>Scenedemusa quadricauda</i>	Dimethomorph pyrimethanil Isoproturon	24% 10%	Oletteet <i>al.</i> , (2010)
			58%	
4.	<i>Selenastrum capricornutum</i> <i>Scenedesmus acutus</i>	benzo(a)pyrene	99% 95%	Laseraet <i>al.</i> , (2016)
5.	<i>Nannochloris oculata</i>	Lindane	73% -68.2%	Perez- legaspriet <i>al.</i> , (2016)
6.	<i>Chlamydomonas reinhardtii</i>	Organophosphru sTrichloforn	100%	Wanet <i>al.</i> ,(2020)
7.	<i>Chalamydomona s reinhardtii</i>	Fluroxypyr	57%	Zhang <i>et al.</i> , (2011)
8.	<i>Chlamydomonas reinhardtii</i>	Trichlorforn	51.3%	Wan <i>et al.</i> ,(2020)
9.	<i>Chlorella vulgaris</i> <i>Scenedesmus quadricuda</i> <i>Spirulina platensis</i>	Malathion Nickle Lead Cadmium	99% 95% 89% 88%	Abdel-razek <i>et al.</i> , (2019)
10.	<i>Nostoc muscorum</i> <i>S. platensis</i>	Malathion	91%	Ibrahim <i>et al.</i> ,(2014)
11.	<i>Serratia marcescens</i>	Chlorpyrifos Fenitrothion Parathion	58.9% 70.5% 82.5%	Cyconet <i>al.</i> , (2013)

12.	<i>Serratia liquefaciens</i> <i>Serratia marcescens</i> <i>Pseudomonas sp.</i>	Diazin	80%-92%	Cyconet <i>al.</i> ,(2009)
-----	--	--------	---------	----------------------------

X. DIFFERENT PESTICIDES CAN BE BROKEN DOWN BY BACTERIA IN BOTH LIQUID AND SOIL ENVIRONMENTS.

Potentials for bacterial bioremediation are advantageous from an environmental and financial standpoint. The parent ingredient of a pesticide must be completely oxidized in order to produce carbon dioxide and water, which gives microorganisms energy. Pesticides are discovered to be degraded by Bacterium Raoultella sp. (Uquabet *al.*, 2016). Group of bacteria are present in high concentration in soils called actinobacteria (Alvarez *et al.*, 2017). Most representative pesticide-degrading genera of actinobacteria such as, Arthrobacter, *Rhodococcus*, *Streptomyces*, *Frankia*, *Janibacter*, *Kokuria*, *Mycobacterium*, *Nocardia*, and *Psuedonocardia* (Alvarez *et al.*, 2017).

Table 6: Summary of Bioremediation Potential of Pesticides by Microorganism

Sr. No.	Microorganism	Pesticides	Percentage of Removal	References
1.	White rot fungi	Aldicarb Atrazine Alacholar	47% 98% 62%	Haieet <i>al.</i> , (2011)
2.	<i>Pseudomonas</i>	Crude oil	73.7%	Maganet <i>al.</i> , (2010)
3.	<i>B. cereus</i> <i>B. safensis</i>	Methomyl	88.25% 77.5%	Roy <i>et al.</i> , (2017)
4.	<i>Bacillus sefensis</i>	Diazinon	63%	Aly <i>et al.</i> ,(2017)
5.	<i>Phanerochaete velutina</i>	Polyaromatic Hydrocarbons PHAs	96%	Winqvist <i>et al.</i> , (2014)
6.	<i>Pleurotus ostreatus</i>	Polychlorinatrd biphenyls (PCBs)	50.5%	Stella <i>et al.</i> , (2017)

7.	<i>Rhizopus sp.</i> <i>Penicilliumfuniculosm</i> <i>Aspergillus sydowi</i>	Petroleum hydrocarbon (TPH) Aliphatic hydrocarbons(AH) Polycyclic aromatic hydrocarbon(PAH)	36% 30% 17%	Lopez <i>et al.</i> ,(2008)
8.	<i>T. versicolor</i> (R26 and R101) <i>P. ostreatus</i>	Dieldrin Trifluralin Simazine	80%	Fragoeiro <i>et al.</i> ,(2005)
9.	<i>Pleurotus cystidiosus</i> <i>Pleurotus sajor-caju</i> <i>Trametes socotrana</i> <i>Polystictus sanguine aus</i> <i>Trametes veriscolar</i> <i>Phanerochaete chrysosporium</i>	Simazine Trifluranin Dieldrin	50%	Maganet <i>et al.</i> ,(2010)
10.	<i>Novosphingobium Strain DY4</i>	2,4-dichlorophenoxyacetic acid	50and 95%	Diaet <i>et al.</i> ,(2005)
11.	<i>Pseudomonas</i>	Atrazine Carbofuran Glyphosate	90%	Echeverria <i>et al.</i> , (2020)
12.	<i>Trichoderma</i>	Dichlorvos Glyphosate	100%	Poveda <i>et al.</i> , (2022)
13.	<i>Aspergillus oryzae</i> <i>Penicillium</i> <i>Trichoderma</i>	Glycophosate	60%	Correa <i>et al.</i> , (2019)
14.	<i>Stenotrophomonas sp.</i>	DDT DDE	81% 55%	Xie <i>et al.</i> , (2022)
15.	<i>Sphingomonas trueperi</i> CW3	Allethrin	93%	Bhatt <i>et al.</i> ,(2020)
16.	<i>Brucella spp.</i>	Dimethoate	83%	Ahmad <i>et al.</i> ,(2022)

In comparison to other bacteria, fungi are more significant to pollution because they can quickly colonize and their hyphae can penetrate soil to access contaminants faster (Readdy and Mathew 2002; Harms *et al.*, 2011).

Fungal enzymes like lignin, degrading enzymes, laccase, oxidoreductases, and peroxidases have the notable ability to remove the pesticides and insecticides residue from contaminated soil. Fungi are eukaryotic organisms that are diverse throughout the world in any environmental condition. They also have a high bioremediation potential to degrade

pesticide residue. Pesticide degradation is influenced by soil's physical and chemical characteristics, contaminated microorganism kinds, and concentration levels(Khatoon *et al.*,2021).

REFERENCES

- [1] Abdel-Razek, M. A., Abozeid, A. M., Eltholth, M. M., Abouelenien, F. A., El-Midany, S. A., Moustafa, N. Y., & Mohamed, R. A. (2019). Bioremediation of a pesticide and selected heavy metals in wastewater from various sources using a consortium of microalgae and cyanobacteria. *Slov vet*, 56(Suppl 22), 61-73.
- [2] Ahmad, S., Chaudhary, H. J., & Damalas, C. A. (2022). Microbial detoxification of dimethoate through mediated hydrolysis by *Brucella* sp. PS4: Molecular profiling and plant growth-promoting traits. *Environmental Science and Pollution Research*, 29(2), 2420-2431.
- [3] Akpinar, M., & Ozturk Urek, R. (2017). Induction of fungal laccase production under solid state bioprocessing of new agroindustrial waste and its application on dye decolorization. *3 Biotech*, 7, 1-10.
- [4] Aly, M. M., Al-aidaroos, B.A., &Alfassi, F. A. (2017). Factors Affecting Biodegradation of the Organophosphorus Insecticide Diazinon by Bacterial Mono- Culture of *Bacillus Sefensis* 7 Isolated from the Rhizosphere of Date Palm tree. *IOSR-JPBS*, 12(3), 28-26.
- [5] Anjaneya, O., Souche, S. Y., Santoshkumar, M., &Karegoudar, T. B. (2011). Decolorization of sulfonated azo dye Metanil Yellow by newly isolated bacterial strains: *Bacillus* sp. strain Ak1 and *Lysinibacillus* sp. strain AK2. *Journal of Hazardous Materials*, 190(1-3), 351-358.
- [6] Ardal, E. (2014). Phycoremediation of pesticides using microalgae.
- [7] Asad, M. A. U., Lavoie, M., Song, H., Jin, Y., Fu, Z., & Qian, H.(2017). Intraction of chiral herbicides with soil microorganism, algae and vascular plants. *Science of the Total Environment* ,580, 1287-1299
- [8] Bayazit, G., ERTIT TASTAN, B. U. R. C. U., & Gul, U. D. (2020). Biosorption, isotherm and kinetic properties of common textile dye by *Phormidiumanimale*. *Global Nest Journal*, 22(1).
- [9] Behl, K., Sinha, S., Sharma, M., Singh, R., Joshi, M., Bhatnagar, A., & Nigam, S. (2019). One-time cultivation of *Chlorella pyrenoidosa* in aqueous dye solution supplemented with biochar for microalgal growth, dye decolorization and lipid production. *Chemical Engineering Journal*, 364, 552-561.
- [10] Ben Ayed, A., Hadrich, B., Sciara, G., Lomascolo, A., Bertrand, E., Faulds, C. B., ... &Mechichi, T. (2022). Optimization of the decolorization of the reactive black 5 by a Laccase-like active cell-free supernatant from *Corioliopsis gallica*. *Microorganisms*, 10(6), 1137.
- [11] Bera, S., Sharma, V. P., Dutta, S., & Dutta, D. (2016). Biological decolorization and detoxification of malachite green from aqueous solution by *Dietziamaris* NIT-D. *Journal of the Taiwan Institute of Chemical Engineers*, 67, 271-284.
- [12] Bhatt, P., Huang, Y., Rene, E. R., Kumar, A.j., & Chen, S. (2020). Mechanism of allethrin biodegradation by a newly isolated *Sphingomonas* strain CW3 from wastewater sludge. *Bioresource Technology*, 305,123074.
- [13] Bheemaraddi, M. C., Patil, S., Shivannavar, C. T., &Gaddad, S. M. (2014). Isolation and characterization of *Paracoccus* sp. GSM2 capable of degrading textile azo dye reactive violet 5. *The Scientific world Journal*, 2014.
- [14] Birmole, R., & Aruna, K. (2019). Optimisation studies of reactive red 120 decolorisation by *Shewanellahaliois* RDB_1. *J Glob Biosci*, 8, 6324-67.
- [15] Biyik, H., Basbulbul, G., Kalyoncu, F., Kalmis, E., &Oryasin, E. (2012). Biological decolorization of textile dyes from isolated microfungi. *Journal of Environmental Biology*, 33(3), 667.
- [16] Boufercha, O., Moreira, I. S., Castro, P. M., &Boudemagh, A. (2022). Actinobacteria isolated from wastewater treatment plants located in the east-north of Algeria able to degrade pesticides. *World Journal of Microbiology and Biotechnology* 38, no. 6(2022): 105.
- [17] Chaieb, K., Kouidhi, B., Ayed, L., Hosawi, S. B., Abdulhakim, J. A., Hajri, A., &Altayb, H. N. (2023). Enhanced textile dye removal from wastewater using natural biosorbent and *Shewanella* algae B29: Application of Box Behnken design and genomic approach. *Bioresource Technology*, 374, 128755.
- [19] Chen, C. C., Liao, H. J., Cheng, C. Y., Yen, C. Y., & Chung, Y.C. (2007). Biodegradation of crystal violet by *Pseudomonas putida*. *Biotechnology letters*, 29, 391-396.
- [20] Chen, C. Y., Kuo, J. T., Cheng, C. Y., Huang, Y. T., Ho, I. H., & Chung, Y. C. (2009). Biological decolorization of dye solution containing malachite green by *Pandoraeapulmonicola* YC32 using a batch and continuous system. *Journal of hazardous materials*, 172(2-3), 1439-1445.

- [21] Chhabra, M., Mishra, S., & Sreekrishnan, T. R. (2008). Mediator-assisted decolorization and detoxification of textile dyes/dye mixture by *Cyathobulleri laccase*. *Applied biochemistry and biotechnology*, 151, 587-598.
- [22] Chia, M. A., Odoh, O. A., & Ladan, Z. (2014). The indigo blue decolorization potential of immobilized *Scenedesmus quadricauda*. *Water, Air, & Soil pollution*, 225, 1-9.
- [23] Correa, L. O., Bezerra, A. F. M., Honorato, L. R. S., Cortez, A. C. A., Souza, J. V. B., & Souza, E. S. (2021). Amazonian soil fungi are efficient degraders of glyphosate herbicide; novel isolates of *Penicillium*, *Aspergillus*, and *Trichoderma*. *Brazilian journal of biology*, 83, e242830.
- [24] Cycon, M., Wojcik, M., & Piotrowska-Segat, Z. (2009). Biodegradation of the organophosphorus insecticide diazinon by *Serratia* sp. and *Pseudomonas* sp. and their use in bioremediation of contaminated soil. *Chemosphere*, 76(4), 494-501.
- [25] Cycon, M., Zmijowska, A., Wojcik, M., & Piotrowska-Segat, Z. (2013). Biodegradation and bioremediation potential of diazinon-degrading *Serratia marcescens* to remove other organophosphorus pesticides from soils. *Journal of Environmental Management* 117, 7- 16.
- [26] Dai, Y., Li, N., Zhao, Q., & Xie, S. (2015). Bioremediation using *Novosphingobium* strain DY4 for 2,4-dichlorophenoxyacetic acid- contaminated soil and impact on microbial community structure. *Biodegradation* 26, 161-170.
- [27] Devonshire, A. L., & Field, L. M. (1991) Gene amplification and insecticide resistance.
- [28] *Annual review of entomology*, 36 (1), 1-21.
- [29] Dosnon-Olette, R., Trotel-Aziz, P., Couderchet, M., & Eullaffroy, P. (2010). Fungicides and herbicide removal in *Scenedesmus* cell suspensions. *Chemosphere*, 79(2), 117-123.
- [30] Eggen, T. (1999). Application of fungal substrate from commercial mushroom production-*Pleurotus ostreatus*-for bioremediation of creosote contaminated soil. *International Biodeterioration & Biodegradation*, 44(2-3), 117-126.
- [31] El Bouraie, M., & El Din, W. S. (2016). Biodegradation of Reactive Black 5 by *Aeromonas hydrophila* strain isolated from dye-contaminated textile wastewater. *Sustainable Environment Research*, 26(5), 209-216.
- [32] FAO, F. (2018). Food and agriculture organization of the United Nations. Rome, URL://faostat. Fao. Org.
- [33] Fragoeiro, S. (2005). Use of fungi in bioremediation of pesticides.
- [34] Garg, N., Garg, A., & Mukherji, S. (2020). Eco-friendly decolorization and degradation of reactive yellow 145 textile dye by *Pseudomonas aeruginosa* and *Thiosphaerapantotropha*. *Journal of environmental management*, 263, 110383.
- [35] Gattullo, C. E., Bahrs, H., Steinberg, C. E., & Lofferdo, E. (2012). Removal of bisphenol A by the freshwater green alga *Monoraphidium braunii* and the role of natural organic matter. *Science of the Total Environment*, 416, 501-506.
- [36] Ghasemi, Y., Rasoul-Amini, S., & Fotooh-Abadi, E. (2011). The biotransformation, biodegradation, and bioremediation of organic compounds by microalgae 1. *Journal of phycology*, 47(5), 969-980.
- [37] Gongora-Echeverria, V. R., Garcia-Escalante, R., Rojas-Herrera, R., Giacoman-Vallejos, G., & Ponce-Caballero, C. (2020). Pesticide bioremediation in liquid media using a microbial consortium and bacteria -pure strains isolated from a biomixture used in agricultural areas. *Ecotoxicology and Environmental Safety*, 200, 110734.
- [38] Green, C., & Hoffnagle, A. (2004). *Phytoremediation field studies database for chlorinated solvents, pesticides, explosives, and metals* (p. 168). Washington, DC: US Environmental Protection Agency.
- [39] Hadibarata, T., Adnan, L. A., Yusoff, A. R. M., Yuniarto, A., Rubiyatno, Zubir, M. M. F. A., ... & Naser, M. A. (2013). Microbial decolorization of an azo dye reactive black 5 using white-rot fungus *Pleurotus eryngii* F032. *Water, Air, & Soil Pollution*, 224, 1-9.
- [40] Hai, F. I., Modin, O., Yamamoto, K., Fukushi, K., Nakajima, F., & Nghiem, L. D. (2012). Pesticide removal by a mixed culture of bacteria and white-rot fungi. *Journal of the Taiwan Institute of Chemical Engineers*, 43(3), 459-462.
- [41] Hakim, A., Hoque, S., & Ullah, S. M. (2014). Decolorization of Methyl Red by *Staphylococcus saprophyticus* strain AUCASVE3 Isolated from Textile Effluent. *Environment Asia*, 7(1).
- [42] Hamouda, R. A., El-Naggar, N. E. A., & Abou-El-Souod, G. W. (2022). Simultaneous bioremediation of Disperse orange-2RL Azo dye and fatty acids production by *Scenedesmus obliquus* cultured under mixotrophic and heterotrophic conditions. *Scientific Reports*, 12(1), 20768.
- [43] Harmss, H., Schlosser, D., & Wick, L. Y. (2011). Untapped potential: exploiting fungi in bioremediation of hazardous chemicals. *Nature Reviews Microbiology*, 9(3), 177-192.
- [44] Huang, J., Liu, D., Lu, J., Wang, H., Wei, X., & Liu, J. (2016). Biosorption of reactive black 5 by

- modified *Aspergillus versicolor* biomass: kinetics, capacity and mechanism studies. *Colloids and Surfaces A: Physicochemical and Engineering Aspects*, 492, 242- 248.
- [45] Ibrahim, W. M., Karam, M. A., El-Shahat, R.M., &Adway, A. A. (2014). Biodegradation and utilization of organophosphorus pesticide malathion by cyanobacteria. *BioMed research international*, 2014.
- [46] Jadhav, J. P., Kalyani, D. C., Telke, A. A., Phugare, S. S., &Govindwar, S. P. (2010). Evaluation of the efficacy of a bacterial consortium for the removal of color, reduction of heavy metals, and toxicity from textile dye effluent. *BioresourceTechnology*, 101(1), 165- 173.
- [47] Jadhav, U. U., Dawkar, V. V., Ghodake, G. S., &Govindwar, S. P. (2008). Biodegradation of Direct Red 5B, a textile dye by newly isolated *Comamonas* sp. UVS. *Journal of Hazardous Materials*, 158(2-3), 507-516.
- [48] Jebapriya, G. R., &Gnanadoss, J. J. (2013). Bioremediation of textile dye using white rot fungi: a Review. *International Journal of Current Research and Review*, 5(3), 1.
- [49] Jin, Z. P., Luo, K., Zhang, S., Zheng, Q., & Yang, H. (2012). Bioaccumulation and catabolism of prometryne in green algae. *Chemosphere*, 87(3), 278-284.
- [50] Kensa, V.M. (2011). Bioremediation- an overview. *Journal of Industrial Pollution Control*, 27(2), 161-168.
- [51] Kesebir, A. O., Kilic, D., Sisecioglu, M., Adiguzel, A., &Kufrevioglu, O. I. (2021). Recombinant laccase production from *Bacillus licheniformis* O12: Characterization and its application for dye decolorization. *Biologia*, 76(11), 3429-3438.
- [52] Khatoon, H., Rai, J. P. N., &Jillani, A. (2021). Role of fungi in bioremediation of contaminated soil. In *Fungi bioprospects in sustainable agriculture, environment and nano-technology* (pp. 121-156). Academic press.
- [53] Kumar, A., & Singh, J. S. (2017). Cyanoremediation: a green-clean tool for decontamination of synthetic pesticides from agro-and aquatic ecosystems. *Agro- Environmental Sustainability: Volume 2: Managing Environmental Pollution*, 59-83.
- [54] LEE, M. D., & SWINDOLL, C. M. (1993). Bioventing for in situ remediation.
- [55] *Hydrological sciences journal*, 38(4), 273-282.
- [56] Li, X., Liu, D., Wu, Z., Li, D., Cai, Y., Lu, Y., ... & Xue, H. (2020). Multiple Tolerances and Dye Decolorization Ability of a Novel Laccase Identified from *Staphylococcus Haemolyticus*. *Journal of Microbiology and Biotechnology*, 30(4), 615.
- [57] Magan, N., Fragoerio, S., & Bastos, C. (2010). Environmental factors and bioremediation of xenobiotics using white rot fungi. *Mycobiology*, 38 (4), 238-248.
- [58] Mahalakshmi, S., Lakshmi, D., & Menaga, U. (2015). Biodegradation of different concentration of dye (Congo red dye) by using green and blue green algae. *Int J Environ Res*, 9(2), 735-744.
- [59] Maitra, S. (2018). In situ bioremediation-An overview. *Responsibility of Life Science Informatics Publications*.
- [60] Mali, H., Shah, C., Patel, D. H., Trivedi, U., & Subramanian, R.B. (2022). Degradation insight of organophosphate pesticide chlorpyrifos through novel intermediate 2, 6- dihydroxypyridine by *Arthrobacter* sp. HM01. *Bioresources and Bioprocessing*, 9(1), 1- 14.
- [61] Mancera-Lopez, M. E., Esparza-Garcia, F., Chavez-Gomez, B., Rodriguez-Vazquez, R., Saucedo-Castaneda, G., & Barrera-Cortes, J. (2008). Bioremediation of an aged hydrocarbon contaminated soil by a combined system of biostimulation- bioaugmentation with filamentous fungi. *International Biodeterioration & Biodegradation*, 61(2), 151- 160.
- [62] Mathur, P., Saini, S., Paul, E., Sharma, C., & Mehtani, P. (2021). Endophytic fungi mediated synthesis of iron nanoparticles: Characterization and application in methylene blue decolorization. *Current Research in Green and Sustainable Chemistry*, 4, 100053.
- [63] Mazmanci, M. A., &Unyayar, A. (2005). Decolourisation of Reactive Black 5 by *Funaliatrogii* immobilised on *Luffa cylindrica* sponge. *Process Biochemistry*, 40(1), 337- 342.
- [64] Mohana, S., Shrivastava, S., Divecha, J., &Madamwar, D. (2008). Response surface methodology for optimization of medium for decolorization of textile dye Direct Black 22 by a novel bacterial consortium. *Bioresource technology*, 99(3), 562-569.
- [65] Muthukumar, P., Paramasivan, T., Priya, M., Abarna, S., Ramalingam, P., & Saraswathy, N. (2017). Biodecolorization of textile reactive dyes by *Penicillium chrysosporium* and *Penicillium simplicissimum*- A comparative study. *Research Journal of Engineering and Technology*, 8(1), 33-38.
- [66] Nie, J., Sun, Y., Zhou, Y., Kumar, M., Usman, M., Li, J., ... & Tsang, D. C. (2020). Bioremediation of water containing pesticides by microalgae: Mechanism, methods, and prospects for future research, *Science of the Total Environment*, 707, 136080.

- [67] Omar, H. H. (2008). Algal decolorization and degradation of monoazo and diazo dyes. *Pak J Biol Sci*, 11(10), 1310-1316.
- [68] Ortiz-Hernandez, M. L., Sanchez-Salinas, E., Dantan-Gonzalez, E., & Castrejón-Godínez, M. L. (2013). Pesticide biodegradation: mechanisms, genetics and strategies to enhance the process. *Biodegradation-life of science*, 10, 251-287.
- [70] Parikh, A., & Madamwar, D. (2005). Textile dye decolorization using cyanobacteria. *Biotechnology letters*, 27, 323-326.
- [71] Patel, Y., & Gupte, A. (2016). Evaluation of bioremediation potential of isolated bacterial culture YPAG-9 (*Pseudomonas aeruginosa*) for decolorization of sulfonated di-azodye reactive red HE8B under optimized culture conditions. *Int J Curr Microbiol Appl Sci*, 5(8), 258-272.
- [72] Patil, P. S., Shedbalkar, U. U., Kalyani, D. C., & Jadhav, J. P. (2008). Biodegradation of Reactive Blue 59 by isolated bacterial consortium PMB11. *Journal of Industrial Microbiology and Biotechnology*, 35(10), 1181-1190.
- [73] Perez-Legaspi, I. A., Ortega-Clemente, L. A., Moha-Leon, J. D., Rios-Leal, E., Gutierrez, S. C. R., & Rubio-Franchini, I. (2016). Effect of the pesticide lindane on the biomass of the microalgae *Nannochloris oculata*. *Journal of Environmental Science and Health, Part B*, 51(2), 103-106.
- [74] Poveda, J. (2022). Trichoderma role in anthropogenic pollutions mycoremediation: pesticides and heavy metals. In *Advances in Trichoderma Biology for Agricultural Applications* (pp.465-497). Cham: Springer International Publishing.
- [75] Prabhakar, Y., Gupta, A., & Kaushik, A. (2022). Using indigenous bacterial isolate *Nesterenkonia lacusekhoensis* for removal of azo dyes: A low-cost ecofriendly approach for bioremediation of textile wastewaters. *Environment, Development and Sustainability*, 24(4), 5344-5367.
- [76] Premaratne, M., Nishshanka, G. K. S. H., Liyanaarachchi, V. C., Nimarshana, P. H. V., & Ariyadasa, T. U. (2021). Bioremediation of textile dye wastewater using microalgae: current trends and future perspectives. *Journal of Chemical Technology & Biotechnology*, 96(12), 3249-3258.
- [77] Qu, Y., Shi, S., Ma, F., & Yan, B. (2010). Decolorization of reactive dark blue KR by the synergism of fungus and bacterium using response surface methodology. *Bioresource technology*, 101(21), 8016-8023.
- [78] Reddy, C. A., & Mathew, Z. A. C. H. A. R. I. A. (2001, November). Bioremediation potential of white rot fungi. In *British mycological society symposium series* (Vol. 23, pp. 52-78).
- [79] Revathi, S., Kumar, S. M., Santhanam, P., Kumar, S. D., Son, N., & Kim, M. K. (2017). Bioremoval of the indigo blue dye by immobilized *Chlorella vulgaris* (PSBDU06).
- [80] Ridolfi, A. S., Alvarez, G. B., & Rodriguez Girault, M. E. (2014). Organochlorinated contaminants in general population of Argentina and other Latin America: *Current Research and Perspectives*, 17-40.
- [81] Roy, T., & Das, N. (2017). Isolation, characterization, and identification of two methomyl-degradating bacteria from a pesticide-treated crop field in West Bengal, India. *Microbiology*, 86, 753-764.
- [82] San Keskin, N. O., Celebioglu, A., Uyar, T., & Tekinay, T. (2015). Microalgae immobilized by nanofibrous web for removal of reactive dyes from wastewater. *Industrial & Engineering Chemistry Research*, 54(21), 5802-5809.
- [83] Saratale, R. G., Saratale, G. D., Chang, J. S., & Govindwar, S. P. (2009). Ecofriendly degradation of sulfonated diazo dye CI Reactive Green 19A using *Micrococcus glutamicus* NCIM-2168. *Bioresource technology*, 100(17), 3897-3905.
- [84] Shah, M. P. (2014). Bioremediation application of *Bacillus megaterium* PMS82 in microbial degradation of acid orange dye. *Int J Environ Bioremediat Biodegrad*, 2, 93-99.
- [85] Sharma, I. (2020). Bioremediation techniques for polluted environment: concept, advantages, limitations, and prospects. In *Trace metals in the environment-new approaches and recent advances*. IntechOpen.
- [86] Sharma, I. (2020). Bioremediation techniques for polluted environment: concept, advantages, and prospects. In *Trace metals in the environment- new approaches and recent advances*. IntechOpen.
- [87] Singh, T., & Singh, D. K. (2017). Phytoremediation of organochlorine pesticides: concept, method, and recent developments. *International journal of phytoremediation*, 19(9), 834-843.
- [88] Singhal, M., Jadhav, S., Sonone, S. S., Sankhla, M. S., & Kumar, R. (2022). Microalgae based sustainable bioremediation of water contaminated by pesticides. *Biointerface Res. Appl. Chem*, 12(1), 149-169.
- [89] Stella, T., Covino, S., Cvanarova, M., Filipova, A., Petruccioli, M., D'Annibale, A., & Cajthaml, T. (2017). Bioremediation of long-term PCB-contaminated soil by white-rot fungi. *Journal of Hazardous Materials*, 324, 701-710.
- [90] Sudarshan, S., Harikrishnan, S., Rathi Bhuvaneswari, G., Alamelu, V., Aanand, S., Rajasekar, A., & Govarthanan, M. (2023). Impact of textile dyes on human health and bioremediation of textile industry effluent using microorganisms: current status and future prospects. *Journal of Applied Microbiology*,

- 134(2), Ixac064.
- [93] Tomer, A., Singh, R., Singh, S.K., Dwivedi, S.A., Reddy, C.U., Keloth, M. R. A., & Rachel, R. (2021). Role of fungi in bioremediation and environmental sustainability. *Mycoremediation and Environmental Sustainability: Volume 3*, 187-200.
- [94] Unnikrishnan, S., Kha, M. H., & Ramaligam, K. (2018). Dye-tolerant marine *Acinetobacter baumannii*-mediated biodegradation of reactive red. *Water Science and Engineering*, 11(4), 265-275.
- [95] Uquab, B., Mudasir, S., & Nazir, R. (2016). Review on bioremediation of pesticides. *J Bioremediat Biodegrad*, 7(343), 2.
- [96] Wan, L., Wu, Y., Ding, H., & Zhang, W. (2020). Toxicity, biodegradation, and metabolic fate of organophosphorus pesticide trichlorfon on the freshwater algae *Chlamydomonas reinhardtii*. *Journal of agricultural and food chemistry*, 68(6), 1645-1653.
- [97] Wang, R., Li, H., Liu, Y., Chen, J., Peng, F., Jiang, Z., ... & Song, H. (2022). Efficient removal of azo dyes by *Enterococcus faecalis* R1107 and its application in stimulated textile effluent treatment. *Ecotoxicology and Environmental Safety*, 238, 113577.
- [98] Wang, Y., Mu, W., Sun, X., Sun, X., Lu, X., Fan, Y., & Liu, Y. (2020). Physiological response and removal ability of freshwater diatom *Nitzschia palea* to two organophosphorus pesticides. *Chemistry and Ecology*, 36(9), 881-902.
- [99] Wang, Y., Wen, T., Li, J. J., He, J., Qin, W. C., Su, L. M., & Zhao, Y. H. (2014). Investigation on the relationship between bioconcentration factor and distribution coefficient based on class-based compounds: The factors that affect bioconcentration. *Environmental toxicology and pharmacology*, 38(2), 388-396.
- [100] Winquist, E., Bjorklof, k., Schultz, E., Rasanen, M., Salonen, K., Anasonye, F., ... & Tuomela, M. (2014). Bioremediation of PAH-contaminated soil with fungi- From laboratory to field scale. *International biodeterioration & Biodegradation* 86 (2014): 238- 247.
- [101] Xie, H., Liu, R., Xu, Y., Liu, X., Sun, F., Ma, Y., & Wang, Y. (2022). Effect of In Situ Bioremediation of soil Contaminated with DDT and DDE by *Stenotrophomonas* sp. Strain DXZ9 and Ryegrass on Soil Microorganism. *Microbiology Research*, 13(1), 64-86.
- [102] Yang, Q., Yang, M., Pritsch, K., Yediler, A., Hagn, A., Schloter, M., & Kettrup, A. (2003). Decolorization of synthetic dyes and production of manganese-dependent peroxidase by new fungal isolates. *Biotechnology Letters*, 25, 709-713.
- [103] Yang, Y., Wang, G., Wang, B., Du, L., Jia, X., & Zhao, Y. (2011). Decolorization of malachite green by a newly isolated *Penicillium* sp. YW 01 and optimization of decolorization parameters. *Environmental Engineering Science*, 28(8), 555-562.
- [104] Zhang, S., Qiu, C. B., Zhou, Y., Jin, Z. p., & Yang, H. (2011). Bioaccumulation and degradation of pesticide fluroxypyr are associated with toxic tolerance in green alga *Chlamydomonas reinhardtii*. *Ecotoxicology*, 20, 337-347.