

AN OPTIMUM SOLUTION WITH HEXAGONAL NUMBERS BY USING DOMINANCE PROPERTY

Abstract

In this paper approaches the method to solve dominance property consider the matrix game taken as Hexagonal fuzzy numbers and using ranking technique. It has different ways to this matrix game solved by dominance property. The intent of this research is to identify the value of the game.

Keywords: Hexagonal Numbers, dominance property, Graphical method.

Author

K. Preethi
Research Scholar
Department of Mathematics
Vellalar College for Women
Erode, Tamil Nadu, India.
preethikips@gmail.com

I. INTRODUCTION

In today's real life, there are many complex situations in engineering and business, in which experts and decision makers struggle with uncertainty and hesitation. In many practical situations, collection of crisp data of various parameters is difficult due to lack of exact communications, error in data, market knowledge and customer's satisfaction. The information available is sometimes vague and insufficient. While a decision maker is used to characterize a real-world issue involving uncertainty, fuzzy sets are the result. The exact evaluation of membership values cannot be feasible due to insufficient data. Additionally, it is unattainable to evaluate non-membership values. In order to cope with unreliable data when making decisions as well as thrive in an indeterministic surroundings, Bellman and Zadeh [1] established the idea of fuzziness.

Study on hexagonal fuzzy number with R. Pavithra and Rosario[2]. Intuitionistic fuzzy set theory gave more information about the problem than fuzzy set theory did. As an outcome, intuitionistic fuzzy set theory has been applied by many academics in a variety of academic fields [1–9]. It is often hard to fully convey the statistical information underlying a decision-making dilemma. Thus, it might be represented by fuzzy numbers. For expressing fuzzy numbers, membership functions' triangular and trapezoidal geometries are commonly used in research [10–14]. In order to arrive at an entirely ambiguous agreement, Srinivasan and Karthikeyan [15] explored a two-stage cost-limiting fuzzy transportation problem when inventories and demand both involve fuzzy numbers. In challenging circumstances where things are unclear. To address transportation concerns, Srinivasan et al. have implemented a suggested technique employing trapezoidal fuzzy numbers [10]. Contrary to this, intuitionistic fuzzy numbers in dubious problem scenarios provide more details and imply significant ambiguity.

The hexagonal fuzzy number having its dominant attribute is formulated in this study as a cost to account for both supply and demand ambiguity and hesitancy. The new ranking measure proposed in this paper proves to be efficient over the other fuzzy ranking existing techniques.

II. PRELIMINARIES

- 1. Definition 1:** Fuzzy numbers are defined as a) a normal and convex fuzzy set; b) a support that is bounded; and c) a fuzzy set A of the real line R with membership function $A(x): R \rightarrow [0,1]$. Each in $[0,1]$ has an interval that should be closed.
- 2. Definition 2:** The hexagonal fuzzy number H is a fuzzy integer. The function of membership of the hexagonal fuzzy number AH , denoted by $(a, b, c, d, e, f; 1)$, is shown the following:

$$\mu_{AH}(x) = \begin{cases} \frac{y-a}{b-a}, & a \leq y \leq b \\ 1, & b \leq y \leq c \\ \frac{d-y}{d-c}, & c \leq y \leq d \\ 0, & \text{otherwise} \\ \frac{y-c}{d-c}, & c \leq y \leq d \\ 1, & d \leq y \leq e \\ \frac{f-y}{f-c}, & e \leq y \leq f \\ 0, & \text{otherwise} \end{cases}$$




Figure 1: Hexagonal Fuzzy

3. Definition 3: Assuming $A_H=(a_1, a_2, a_3, a_4, a_5, a_6)$ and $B_H=(b_1, b_2, b_3, b_4, b_5, b_6)$ are two fuzzy hexagonal numbers that can be combined in a variety of ways.

- Addition $A_H + B_H = (a_1 + b_1, a_2 + b_2, a_3 + b_3, a_4 + b_4, a_5 + b_5, a_6 + b_6)$
- Subtraction $A_H - B_H = (a_1 - b_6, a_2 - b_5, a_3 - b_4, a_4 - b_3, a_5 - b_2, a_6 - b_1)$
- Multiplication $A_H * B_H = (a_1 * b_1, a_2 * b_2, a_3 * b_3, a_4 * b_4, a_5 * b_5, a_6 * b_6)$

4. Definition 4: The extent to which a using Hexagonal fuzzy number $A_H=(a_1, a_2, a_3, a_4, a_5, a_6)$ is specified as the ranking methodology depending on the CC approach applying the definition given above.

$$R(CC_H) = \left(\frac{2a_1 + 3a_2 + 4a_3 + 3a_4 + 3a_5 + 2a_6}{18}, \frac{5}{18} \right)$$

III. DOMINANCE PROPERTY

The more effective approaches are deemed to predominate the lesser ones when it appears that a single of each player's purest approaches remains better than a minimum one of the others.

1. General Rules: If each request in a single row, say k^{th} , is fewer than (or) equivalent to each component in another row, say r^{th} , then k^{th} row is overpowered by r^{th} row. Similarly, if every component in a column, say k^{th} , exceeds than (or) equivalent to every component in another column, say r^{th} , then k^{th} column is predominated by r^{th} column. (ii) Dominated rows (or) columns may be removed to make the payoff matrix smaller while maintaining the effectiveness of the best possible techniques.

2. The Modified Dominance Property: The supremacy of pure tactics is not the only foundation for the prevailing attribute. If two or more different pure strategies are included into a particular approach, it is also referred to as being dominant.

In a broader sense, the i^{th} row can be eliminated if any convex linear combination of certain rows exceeds the i^{th} row. Similar rationales apply to columns.

3. Numerical Example: Consider the following Fuzzy Transportation problem where supply and demand are given as Hexagonal Fuzzy Numbers[7]

	d_1	d_2	d_3	d_4	Supply
o_1	(3,7,11,15,19,24)	(13,18,23,28,33,40)	(6,13,20,28,36,45)	(15,20,25,31,38,45)	(6,8,11,14,19,25)
o_2	(16,19,24,29,34,39)	(3,5,7,9,10,12)	(5,7,10,13,17,21)	(20,23,26,30,35,40)	(9,11,13,15,18,20)
o_3	(11,14,17,21,25,30)	(7,9,11,14,18,22)	(2,3,4,6,7,9)	(5,7,8,11,14,17)	(7,9,11,13,16,20)
Demand	(3,4,5,6,8,10)	(3,5,7,9,12,15)	(6,7,9,11,13,16)	(10,12,14,16,20,24)	(2,3,4,5,6,7)


Solution: Using ranking technique the rank of hexagonal fuzzy cost matrix is obtained.

$$\begin{pmatrix} 3.87 & 6.71 & 6.37 & 7.52 & 3.53 \\ 6.98 & 2.01 & 3.09 & 7.53 & 4.19 \\ 5.09 & 3.47 & 1.33 & 2.65 & 3.27 \\ 1.54 & 2.19 & 2.67 & 4.14 & 1.17 \end{pmatrix}$$

Proceeding by dominance property, the original 5×4 matrix can be reduced to 3×2 matrix, ie. The payoff matrix,

$$\begin{pmatrix} 6.71 & 6.37 & 3.53 \\ 2.01 & 3.09 & 4.19 \end{pmatrix}$$

Applying graphical method,



The cost matrix 3×2 can be reduced by 2×2 matrix, then determine the game's worth.

$$\begin{pmatrix} 6.71 & 3.53 \\ 2.01 & 4.19 \end{pmatrix}$$

By using the mixed strategy method formula we get,
 Then the value of the game $\gamma = 3.92$

IV. CONCLUSION

This paper proposes a value of the game whose costs are taken as hexagonal fuzzy numbers. For future research we propose effective implementation of the hexagonal fuzzy numbers in all fuzzy problems. Further operational research studies might use a similar method for resolving intuitionistic fuzzy situations.

REFERENCES

- [1] Biswas, P., Pramanik, S., Giri, B.C, “Cosine similarity measure based multiattribute decision-making with trapezoidal fuzzy neutrosophic numbers”, *Neutrosophic Sets and Systems* 8, 46–56 (2014).
- [2] Chen, S, “Operations on fuzzy numbers with function principal. Tamkang”, *Journal of management sciences* 6, 13–25 (1985).
- [3] Dong, J., Yang, D.Y., Wan, S.P, “Trapezoidal intuitionistic fuzzy prioritized aggregation operators and application to multi-attribute decision making”, *Iranian Journal of Fuzzy Systems* 12, 1–32 (2015).
- [4] Dubey, D., Mehra, A, “Linear programming with triangular intuitionistic fuzzy number”, *Advances in Intelligent Systems Research* 1, 563–569 (2011).
- [5] Ejegwa, P.A., Akubo, A.J., Joshua, O.M, “Intuitionistic fuzzy set and its application in career determination via normalized euclidean distance method”, *European Scientific Journal* 10, 529–536 (2014).
- [6] Ionescu, C., Rezapour, S., Samei, M.E, “Fixed points of some new contractions on intuitionistic fuzzy metric spaces”, *Fixed Point Theory and Algorithms for Sciences and Engineering* 2013, 168.
- [7] Pavithra.R, G.M. Rosario, “A study on Fuzzy transportation problem using hexagonal fuzzy number”.
- [8] Rezapour, S., Samei, M.E, “Some fixed point results for α -contractive type mappings on intuitionistic fuzzy metric spaces”, *Journal of Advanced Mathematical Studies* 7(1), 176–178 (2014).
- [9] Samei, M.E, “Convergence of an iterative scheme for multifunctions on fuzzy metric spaces”. *Sahand Communications in Mathematical Analysis* 15(1), 91–106 (2019).
- [10] Samei, M.E, “Fixed point theorem for mappings satisfying contractive condition of integral type on intuitionistic fuzzy metric space”. *Journal of Linear and Topological Algebra* 7(3), 183–199 (2018).
- [11] Samei, M.E, “Some fixed point results on intuitionistic fuzzy metric spaces with a graph”, *Sahand Communications in Mathematical Analysis* 13(1), 141–152 (2019).
- [12] Srinivasan, R., Karthikeyan, N., Renganathan, K., Vijayan, D.V.: Method for solving fully fuzzy transportation problem to transform the materials. *Materials Today: Proceedings* (2020).
- [13] Szmidt, E., Kacprzyk, J, “Intuitionistic fuzzy sets in some medical applications”, *Notes on Intuitionistic Fuzzy Sets* 7, 58–64 (2001).
- [14] Wan, S.P., Li, D.F., Rui, Z.F, “Possibility mean, variance and covariance of triangular intuitionistic fuzzy numbers”, *Journal of Intelligent and Fuzzy Systems* 24, 847–858 (2013).
- [15] Zadeh L.A, “Fuzzy sets *Inf Control*” 8,338-353(1965).