

# OPTIMIZING STABILITY OF PLANT EXTRACTS USING LYOPHILIZATION

## Abstract

The growing need for bioactive metabolites necessitates an efficient method of extracting and processing diverse plant extracts. The majority of the existing literature focused on determining the influence of extraction procedure on plant extracts. However, the process used to dry plant extracts after extraction is also critical. The "gold standard" for drying plant extracts is lyophilization, also known as freeze drying, which is crucial for maintaining their quality and increasing their shelf life. A literature search using Pub Med and Google Scholar was done, and a summary of all the particular issues associated with lyophilization was drawn up. Due to the low process temperature, freeze-drying is used as a microencapsulation technology and is especially ideal for the entrapment and protection of delicate bioactive chemicals. The zeta potential, poly dispersity index (PDI), and minimum particle size are typically used to optimize the stability of lyophilized extracts. In this review, the impact of lyophilization on the stability of plant extracts has been explored.

**Keywords:** Lyophilization, Stability, Plant extracts, Encapsulation, Zeta Potential, Poly dispersity Index.

## Authors

### Meenaloshini G

Saveetha College of Pharmacy  
Saveetha Institute of Medical and  
Technical Sciences  
Saveetha University  
Thandalam , Chennai, TamilNadu, India.

### Karthikha V. S

Saveetha College of Pharmacy  
Saveetha Institute of Medical and  
Technical Sciences  
Saveetha University  
Thandalam , Chennai, TamilNadu, India.

### Vaishnavi V

Saveetha College of Pharmacy  
Saveetha Institute of Medical and  
Technical Sciences  
Saveetha University  
Thandalam , Chennai, TamilNadu, India.

### Dravid Ragul Y

Saveetha College of Pharmacy  
Saveetha Institute of Medical and  
Technical Sciences  
Saveetha University  
Thandalam , Chennai, TamilNadu, India.

### Hemalatha K

Saveetha College of Pharmacy  
Saveetha Institute of Medical and  
Technical Sciences  
Saveetha University  
Thandalam , Chennai, TamilNadu, India.  
hemalatha6585@gmail.com

## I. INTRODUCTION

Many drug discovery efforts have used plant extracts as a significant source of bioactive chemicals and some significant medications have been extracted and discovered from plants [1]. Small biomolecules, biologics produced by plants and a recently established third category of medications known as phytopharmaceuticals makeup this class of medications [2]. Around 80% of the world's population, according to the World Health Organization (WHO), relied on medicinal plants as their main source of treatment in 1985 [3]. Various medicinal preparations, including powders, tinctures, tablets, etc., use plant extracts in crude form or standardized fractions [4]. Plant extracts contain a variety of bioactive substances with strong biological effects, such as polyphenols, alkaloids, and terpenes [5]. An adequate strategy is needed to effectively extract and process different plant extracts due to the growing demand for these phytochemicals [6]. Polar solvents (such as water and alcohols), intermediate polar solvents (such as acetone and dichloromethane), and nonpolar solvents (such as n-hexane, ether, and chloroform) are frequently used in the extraction of medicinal plants. Optimal extraction and the ability to maintain the stability of the chemical structure of desired substances are two qualities that define a good solvent [7]. The image illustrate show these bioactive chemicals are now subjected to a variety of thermal, non thermal, and physicochemical variables, including temperature, pH, pressure, sonication, radiation, electric field, humidity, oxygen, and light, as a result of advancements in extraction technology (Figure1) [8]. These processing and storage parameters have been thoroughly investigated by researchers, who have found that they have a considerable impact on the concentration, bioactivity, functionality, bioavailability, and overall stability of the active chemicals [9].




**Figure 1:** Factors Affecting Extraction Technology

## II. BACKGROUND

Enhancing the quality of plants and enhancing the safety, quality, and consistency of the finished products need the adoption of optimized high-quality agricultural and biomass processing processes [10]. The primary active substances utilized in the creation of natural products, which are often offered in the form of powders or tablets in order to ensure good stability and appropriate dosage, are dried plant extracts high in bioactive metabolites [11]. In order to manufacture these natural compounds, it is crucial to optimize the extraction

protocol, which comprises the solvent choice, solvent to biomass ratio, extraction time, and extraction method choice [12]. The majority of the material that was published compared different extraction methods utilized for plant extracts. But it's also crucial to consider the process for drying plant extracts after extraction [13]. Especially because current research indicates that the drying technique used to prepare plant biomass prior to extraction influences their bioactivity and chemical makeup [14]. Similarly, the method of post-extraction drying used has an impact on the chemical and biological characteristics of plant extracts [15]. A mind map enumerating several drying techniques is represented as a figure (Figure 2). Lyophilization is one of the more modern and inventive post-extraction techniques. The "gold standard" for drying plant extracts is lyophilization, also known as freeze drying, which is crucial for maintaining their quality and increasing their shelf life [16]. With the fore mentioned context in mind, this review attempts to provide insight into the various ways that lyophilization can be used to maintain the stability of plant extracts.

## DRYING METHODS




**Figure 2:** Methods of Drying

### III. METHODOLOGY

A PubMed and Google Scholar literature search was conducted using the key terms “lyophilization”, “freeze-drying”, “plant post-extraction methods”, “lyophilized plant extract”, “Stability studies” and “preserving bioactive compounds” to gather the information for this manuscript. An outline of all the special issues exclusive to lyophilization was prepared. All of the data cited has been written entirely in English.

**1. State of the Art: Lyophilization:** Lyophilization in the pharmaceutical industry has undergone continuous growth and steady expansion as we approach a new century. In the freeze-drying process, water is sublimated directly from the solid state (ice) to the vapor state, omitting the liquid state, and is subsequently reabsorbed from the "dry" layer [17]. Although lyophilization is a more recent invention than freeze-drying, they have only been around for about 100 years [18]. The procedure preserves the dried product's quality (including its biological, dietary, and oil factory qualities) [19]. Lyophilization is always used to increase the stability of a product that is sensitive to moisture or to make the product simpler to store or transport [20]. Three basic processes are involved in the

pharmaceutical lyophilization, as depicted in the picture (figure3). The first step is to freeze the product in solution to create a matrix of ice along with additional crystallizable excipients while concentrating other solutes and the active pharmaceutical ingredient (API) inside the interstitial voids [21]. Aqueous medicinal formulations can be frozen at temperatures below  $-35^{\circ}\text{C}$  [22]. The second phase is primary drying, which involves submerging ice in a vacuum at a low temperature. Traditionally, this process is performed at chamber pressures of 40-400 Torr at shelf temperatures ranging from  $-30^{\circ}\text{C}$  to  $-10^{\circ}\text{C}$  [23]. After the primary freeze-drying process is completed and all ice has sublimed, bound moisture remains in the product. Although the product seems dry, the residual moisture level could be as high as 7-8% [24]. As a result, a last process called secondary drying at a warmer temperature is required to remove the residual moisture content to optimal levels. This is known as 'Isothermal Desorption' because the bound water is desorbed from the product [25]. By increasing the shelf temperature above ambient levels, this process is completed. The shelf temperature can be increased to  $15-300^{\circ}\text{C}$  to allow for vacuum-induced desorption of water molecules [26]. Four standard parts are typically included in freeze-dryers: a drying chamber, a vacuum pump, a heat source, and a condenser [27]. The tools and procedures are created to guarantee that the sterility of the product is preserved throughout the lyophilization process [28].



**Figure 3:** Lyophilization Principle and Process Lyophilization and its Multidisciplinary Application

Lyophilization is a common technique used to concentrate extracts or stabilize biological samples before storage [29]. On a laboratory and industrial scale, the freeze-drying method is mostly employed for delicate, temperature-sensitive plant materials [15, 30]. By using the freeze-drying method, plant samples are highly effective in removing water while retaining bioactive components, such as antioxidants [31]. The freeze-drying procedure is frequently employed to preserve and stabilize biological materials because of its benefits [32]. It guards against both the spread of microbes and the deterioration of biological materials linked to the activity of bacteria [33]. In order to prepare plant and food samples for the identification of physiologically active chemicals, the freeze-drying

method are also used [34]. It is used to preserve the chemical makeup of the plant material while stabilizing, enhancing, and/or extending its life [35]. Due to the low process temperature, freeze-drying is used as a micro encapsulation technology, making it especially ideal for the entrapment and protection of delicate bioactive chemicals [36]. In freeze-drying-based encapsulation, the target component and the encapsulating ingredients create an emulsion solution, which is then used to create microcapsules using the freeze-drying method [37]. Additionally, due to the few stages required, the freeze-drying technique is simpler than other micro encapsulation techniques [38].

- 2. Effect of Lyophilization on Stability of Plant Extracts:** In order to maximize zeta potential and minimize polydispersity index (PDI), lyophilized extract stability is typically optimized [39]. The effects of lyophilization technique on stability of plant extracts are enlisted as table (Table1). The poly dispersity index is a size-based indicator of a sample's heterogeneity [40]. Size distribution in a sample, as well as aggregation or agglomeration of the sample during isolation or analysis, can lead to poly dispersity [41]. Good homogeneity is indicated by a PDI value below 0.3 [42]. The zeta potential's size reveals the strength of electrostatic attraction between nearby, similarly charged particles in dispersion [43]. For sufficiently small molecules and particles, a high zeta potential will impart stability, which means that the solution or dispersion will fend off aggregation. Zeta potential readings exceeding 30 mV are often indicative of high stability [44]. Plant extracts' zeta potential is influenced by the extraction process and, consequently, by the extracted chemicals [45]. The thickness of frozen material and its surface area are additional factors. The rate of lyophilization increases with the frozen material's surface area. In contrast, the rate of lyophilization is slower the thicker the frozen substance. The ability of a sample to absorb and transfer heat to the surface undergoing sublimation is influenced by sample thickness [46].

**Table1: Effect of Lyophilization on Stability of Plant Extracts**

S.No	Plant Extract	Stability studies	Source
1	<i>Serpylliherba</i>	Zeta potential at pH6:- 18.7±0.7 mV	[47]
2	<i>Alliumcepa</i> L.	Particle size:177.73 nmPDI:0.45	[48]
3	<i>Artemisiaabsinthium</i>	Zeta potential:-11.9 Particle size: 253.8 nmPDI:0.258	[49]
4	<i>Betavulgaris</i> L. Leaf hydro alcoholic extract nanogel	Zeta Potential:+28.8mV Particle size:247nm PDI:0.259	[50]
5	<i>Anoectochillus burmannicus</i> Ethanollic Extract-Synthesized Selenium Nanoparticles	Zeta Potential:-24.5±1.9m VPDI:0.366±0.074	[51]

6	<i>Hippophaes rhamnoides</i> aqueous extract-containing nanoemulsion	Zeta Potential (mV): -30.11±2.02 Droplet Size (nm):183.07±9.53 PDI:0.295± 0.045	[52]
7	<i>Punica granatum</i> L. Loaded nanophytosomes	Particle size:166.70–144.40nm PDI:<0.5	[53]
8	Chitosan-Coated- <i>Aloeperryi</i> -Solid Lipid Nanoparticles(C-ALP-SLNs)	Zeta Potential:13.6 ±1.1mV Particle size:173.6 ±11.3nm PDI:0.21 ±0.02	[54]
9	<i>Eupatoriumadenophorum</i> Leaf extract-Silver Nanoparticles	Zeta Potential:- 33.4mV Particle size:117.75nm	[55]
10	<i>Calotropis gigantean</i> -loaded poliglusam-silver nanomatrices	Zeta Potential:42.42mV Particle size: 178.5 nmPDI:2.97	[56]

#### IV. CONCLUSION

Researchers have been inspired by the increased demand for plant extracts to create novel drying processes that preserve high concentrations of essential oils and other physiologically active substances without compromising the dried plant material's flavor, color, or texture. Lyophilization appears to be a useful approach for conserving the volatile profile of plants, although its effects depend on the species of plant (the location of volatile accumulation, the type of volatile components, etc.). The lyophilization process' technical specifications also have a big impact on the end product's quality. In order to more accurately examine the therapeutic benefits of plant extracts, additional systematic research is required on the effects of freeze-drying in comparison to alternative preparation methods, such as alcohol extractions of fresh plant material.

- **Competing Interests:** The authors declare no competing interests.
- **Ethics Approval:** There are no studies on human volunteers or animals in this article.
- **Consent to Participate:** The manuscript would not have been possible without the contributions of all of the authors.
- **Consent for Publication:** The decision to publish the manuscript was approved by all of the authors.
- **Conflict of Interest:** There is no financial or other conflict of interest between the authors.
- **Funding:** This work isn't funded.

- **Acknowledgement:** We thank “Saveetha Institute of Medical and Technical Sciences” management for their support and encouragement.

## REFERENCES

- [1] Aware, C. B., Patil, D. N., Suryawanshi, S. S., Mali, P. R., Rane, M. R., Gurav, R. G., & Jadhav, J. P. (2022). Natural bioactive products as promising therapeutics: A review of natural product-based drug development. *South African Journal of Botany*, 151, 512–528.
- [2] Nasim, N., Sandeep, I. S., & Mohanty, S. (2022). Plant-derived natural products for drug discovery: current approaches and prospects. *TheNucleus*, 65(3), 399–411.
- [3] Rajčević, N., Bukvički, D., Dodoš, T., &Marin, P.D.(2022).Interactions between Natural Products-A Review. *Metabolites*, 12(12), 1256.
- [4] Rao, K.S., Haran, R.H., & Rajpoot, V.S.(2022). Value addition: Anovel strategyfor quality enhancement of medicinal and aromatic plants. *Journal of Applied Research on Medicinal and Aromatic Plants*, 31, 100415.
- [5] Teneva, D., & Denev, P.(2023).Biologically Active Compounds from Probiotic Microorganisms and Plant Extracts Used as Bio preservatives. *Microorganisms*, 11(8), 1896.
- [6] Zakeri, A., Bahmani, E., &Aghdam, A. S. R. (2022). Plant extracts as sustainable and green corrosion inhibitors for protection of ferrous metals in corrosive media: A mini review. *Corrosion*
- [7] Abubakar, A., & Haque,M.(2020).Preparation of medicinal plants: Basic extraction and fractionation procedures for experimental purposes. *Journal of Pharmacy and Bio allied Sciences*, 12(1), 1.
- [8] Nelluri, P., Venkatesh, T., Kothakota, A., Pandiselvam, R., Garg, R., Eswaran, V., Mousavi Khaneghah, A. (2022). Recent advances innon-thermal andthermal processing of jackfruit (*Artocarpus heterophyllus* Lam): An updatedreview. *Journal of Food Processing and Preservation*, 46(9).
- [9] Nishimoto-Sauceda, D., Romero-Robles, L. E., & Antunes-Ricardo, M. (2021). Biopolymer nanoparticles: a strategy to enhance stability, bioavailability, and biological effects of phenolic compounds as functional ingredients. *Journal of the Science of Food and Agriculture*, 102(1), 41–52.
- [10] ElNaker, N. A., Daou, M., Ochsenkühn, M. A., Amin, S. A., Yousef, A. F., & Yousef, L. F. (2021). A metabolomics approach to evaluate the effect of lyophilization versus oven drying on the chemical composition of plant extracts. *Scientific Reports*, 11(1).
- [11] Fotsing Yannick Stéphane, F., Kezetas Jean Jules, B., El-Saber Batiha, G., Ali, I., &Ndjakou Bruno, L. (2022). Extraction of Bioactive Compounds from Medicinal Plants and Herbs. *Intech Open*.
- [12] Sánchez- Camargo, A.D.P., Bueno, M.,Parada-Alfonso, F.,Cifuentes, A., & Ibáñez, E.(2019).Hansen solubility parameters for selection of green extraction 0solvents.*Trends in Analytical Chemistry*, 118, 227–237.
- [13] Lazarjani, M. P., Young, O., Kebede, L., &Seyfoddin, A. (2021). Processing and extraction methods of medicinal cannabis: a narrative review.*Journal of Cannabis Research*, 3(1).
- [14] Daud, N. M., Putra, N. R., Jamaludin, R., Md Norodin, N. S., Sarkawi, N. S., Hamzah, M. H. S., . . . Md Salleh, L. (2022). Valorisation of plant seed as natural bioactive compounds by various extraction methods: Areview. *Trends in Food Science & Technology*, 119, 201–214.
- [15] Krakowska-Sieprawska, A., Kielbasa, A., Rafińska, K., Ligor, M., &Buszewski, B. (2022). Modern Methods of Pre-Treatment of Plant Material for the Extraction of Bioactive Compounds. *Molecules (Basel, Switzerland)*, 27(3), 730.
- [16] Sharma, Ashutosh & Khamar, Dikshitkumar & Hayden, Ambrose & Hughes, Helen.(2021).Innovative Drying Technologies for Biopharmaceuticals. *International Journal of Pharmaceutics*.609.121115.
- [17] Vareltzis, P., Stergiou, A., Kalinderi, K., &Chamilaki, M. (2023). Antioxidant Potential of Spray-and Freeze- Dried Extract from Oregano Processing Wastes, Using an Optimized Ultrasound-Assisted Method. *Foods*, 12(13), 2628.
- [18] Juckers, A., Knerr, P., Harms, F., & Strube,J.(2022). Emerging PAT for Freeze-Drying Processes for Advanced Process Control. *Processes*, 10(10), 2059.
- [19] Çetin, N., Ropelewska, E., &Sabanci, K. (2023). The innovative approach to the assessment of differences in image textures between wind fall apple samples dried using non-thermal and thermal techniques without and with ultrasound pretreatment. *Expert Systems with Applications*, 232, 120917.
- [20] Pardeshi, S. R., Kole, E. B., Kapare, H. S., Chandankar, S. M., Shinde, P. J., Boisa, G. S., ... & Junnuthula, V. (2022). Progress on Thin Film Freezing Technology for Dry Powder Inhalation Formulations. *Pharmaceutics*, 14(12), 2632.
- [21] Jameel, F., Alexeenko, A., Bhambhani, A., Sacha, G., Zhu, T., Tchessalov, S., Kumar, L., Sharma, P.,

- Moussa, E., Iyer, L., Fang, R., Srinivasan, J., Tharp, T., Azzarella, J., Kazarin, P., & Jalal, M. (2021). Recommended Best Practices for Lyophilization Validation-2021 Part I: Process Design and Modeling. *AAPS PharmSciTech*, 22(7), 221.
- [22] Yamashita, Y., Ohzuno, Y., Saito, Y., Fujiwara, Y., Yoshida, M., & Takei, T. (2023). Autoclaving-Triggered Hydrogelation of Chitosan-Gluconic acid Conjugate Aqueous Solution for Wound Healing. *Gels*, 9(4), 280.
- [23] Nowak, D., & Jakubczyk, E. (2020). The Freeze-Drying of Foods-The Characteristic of the Process Course and the Effect of Its Parameters on the Physical Properties of Food Materials. *Foods (Basel, Switzerland)*, 9(10), 1488.
- [24] Adali, M. B., Barresi, A. A., Boccardo, G., & Pisano, R. (2020). Spray freeze-drying as a solution to continuous manufacturing of pharmaceutical products in bulk. *Processes*, 8(6), 709.
- [25] Degobert, G., & Aydin, D. (2021). Lyophilization of nanocapsules: instability sources, formulation and process parameters. *Pharmaceutics*, 13(8), 1112.
- [26] Merivaara, A., Zini, J., Koivunotko, E., Valkonen, S., Korhonen, O., Fernandes, F. M., & Yliperttula, M. (2021). Preservation of bio materials and cells by freeze-drying: Change of paradigm. *Journal of Controlled Release*, 336, 480-498.
- [27] Ghaemmaghmanian, Z., Zarghami, R., Walker, G., O'Reilly, E., & Ziaee, A. (2022). Stabilizing vaccines via drying: Quality by design considerations. *Advanced Drug Delivery Reviews*, 187, 114313.
- [28] Kommineni, N., Butreddy, A., Jyothi, V. G. S., & Angsantikul, P. (2022). Freeze Drying for the Preservation of Immuno engineering Products. *I science*.
- [29] Gregorio, N. E., Kao, W. Y., Williams, L. C., Hight, C. M., Patel, P., Watts, K. R., & Oza, J. P. (2020). Unlocking applications of cell-free biotechnology through enhanced shelf life and productivity of *E. coli* extracts. *ACS Synthetic Biology*, 9(4), 766-778.
- [30] Thorat, B.N., Sett, A., & Mujumdar, A.S. (2022). Drying of vaccines and biomolecules. *Drying Technology*, 40(3), 461-483.
- [31] Mustafa, I., & Chin, N.L. (2023). Antioxidant Properties of Dried Ginger (*Zingiber officinale* Roscoe) var. Bentong. *Foods*, 12(1), 178.
- [32] Rockinger, U., Funk, M., & Winter, G. (2021). Current approaches of preservation of cells during (freeze-) drying. *Journal of Pharmaceutical Sciences*, 110(8), 2873-2893.
- [33] Thalen, M., Debrie, A. S., Coutte, L., Raze, D., Solovay, K., Rubin, K., Mielcarek, N., & Loch, C. (2020). Manufacture of a Stable Lyophilized Formulation of the Live Attenuated Pertussis Vaccine BPZE1. *Vaccines*, 8(3), 523.
- [34] Chumroenphat, T., Somboonwatthanakul, I., Saensouk, S., & Siriamornpun, S. (2021). Changes in curcuminoids and chemical components of turmeric (*Curcuma longa* L.) under freeze-drying and low-temperature drying methods. *Food Chemistry*, 339, 128121.
- [35] Zabot, G. L., Schaefer Rodrigues, F., Polano Ody, L., Vinicius Tres, M., Herrera, E., Palacin, H., & Olivera-Montenegro, L. (2022). Encapsulation of bioactive compounds for food and agricultural applications. *Polymers*, 14(19), 4194.
- [36] Raj, G. B., & Dash, K. K. (2022). Microencapsulation of Dragon Fruit Peel Extract by Freeze-Drying Using Hydrocolloids: Optimization by Hybrid Artificial Neural Network and Genetic Algorithm. *Food and Bioprocess Technology*, 15(9), 2035-2049.
- [37] Carpena, M., Garcia-Oliveira, P., Lourenço-Lopes, C., Pereira, A. G., Fraga-Corral, M., Prieto, M. A., & Simal-Gandara, J. (2021). Freeze-Drying Encapsulation as a Mechanism of Choice in Oils: Methods and Mechanism.
- [38] Pudziuvelyte, L., Marksa, M., Sosnowska, K., Winnicka, K., Morkuniene, R., & Bernatoniene, J. (2020). Freeze-drying technique for microencapsulation of *Elsholtzia ciliate* ethanolic extract using different coating materials. *Molecules*, 25(9), 2237.
- [39] Badawi, N., El-Say, K., Attia, D., El-Nabarawi, M., Elmazar, M., & Teaima, M. (2020). Development of Pomegranate Extract-Loaded Solid Lipid Nanoparticles: Quality by Design Approach to Screen the Variables Affecting the Quality Attributes and Characterization. *A C Somega*, 5(34), 21712-21721.
- [40] Pavaloiu, R. D., Shaat, F., Eremia, M., Hlevca, C., Petrescu, M., SHA'AT, M. O. U. S. A., & Sevcenco, C. (2022). Formulation and Characterization of Pullulan Acetate Nanoparticles Loaded with 5-Fluorouracil. *Materiale Plastice*, 59(2), 138-144.
- [41] Mitrović, J. R., Žiberna, M. B., Vukadinović, A., Knutson, D. E., Sharmin, D., Kremenović, A., ... & Savić, S. D. (2023). Freeze-dried nanocrystal dispersion of novel deuterated pyrazoloquinolinone ligand (DK-I-56-1): Process parameters and lyoprotectant selection through the stability study. *European Journal of Pharmaceutical Sciences*, 106557.
- [42] Zhao, Y., Wang, Z., Liu, S., Xie, S., Xie, Y., Li, R., & Oda, H. (2023). Lyoprotectant Formulation and



- Optimization of the J-Aggregates Astaxanthin /BSA/Chitosan Nanosuspension. *Biomolecules*, 13(3), 496.
- [43] Dembek, M., Bocian, S., & Buszewski, B. (2022). Solvent influence on zeta potential of stationary phase—mobile phase interface. *Molecules*, 27(3), 968.
- [44] Schwan, J., Markert, S., Rosenfeldt, S., Schüller, D., Mickoleit, F., & Schenk, A. S. (2023). Comparing the Colloidal Stabilities of Commercial and Biogenic Iron Oxide Nanoparticles That Have Potential In Vitro / In Vivo Applications. *Molecules*, 28(13), 4895.
- [45] Pungle, R., Nile, S. H., Makwana, N., Singh, R., Singh, R. P., & Kharat, A. S. (2022). Green synthesis of silver nanoparticles using the Tridax Procumbens plant extract and screening of its antimicrobial and anticancer activities. *Oxidative Medicine and Cellular Longevity*, 2022.
- [46] Hossain, M. S., & Nakane, K. (2022). Development of a high-performance heat dissipation sheet with three-dimensional alumina fibrous structure fabricated by freeze-drying. *Results in Materials*, 13, 100241.
- [47] Jovanović, A., Volić, M., Obradović, N., & Bugraka, B. (2023). COMPARISON OF LIQUID AND LYOPHILIZED SERPYLLI HERBA WASTE EXTRACTS PREPARED AT DIFFERENT pH VALUES. In *Proceedings of the 2<sup>nd</sup> International Conference on Advanced Production and Processing ICAPP 2022, Novi Sad* (pp.166-175). University of Novi Sad, Faculty of Technology Novi Sad.
- [48] Sahin, O. I., Dundar, A. N., Uzuner, K., Parlak, M. E., Dagdelen, A. F., & Saricaoglu, F. T. (2023). Lyophilized nano-liposomal system for red onion (*Allium cepa*L.) peel anthocyanin: Characterization, bioaccessibility and release kinetics. *Food Bioscience*, 53, 102702.
- [49] Jahan, N., Kousar, F., Rahman, K. U., Touqeer, S. I., & Abbas, N. (2023). Development of Nanosuspension of Artemisia absinthium Extract as Novel Drug Delivery System to Enhance Its Bioavailability and Hepato protective Potential. *Journal of Functional Biomaterials*, 14(8), 433.
- [50] Singh, S., Sindhu, R. K., Alsayegh, A. A., Batiha, G. E. S., Alotaibi, S. S., Albogami, S. M., & Conte-Junior, C. A. (2023). Formulation Development and Investigations on Therapeutic Potential of Nanogel from Beta vulgaris L. Extract in Testosterone-Induced Alopecia. *Bio Med Research International*, 2023.
- [51] Buacheen, P., Chaipuang, A., Karinchai, J., Nuchuchua, O., Imsumran, A., Wongnoppavich, A., & Pitchakarn, P. (2023). Stabilization of Antioxidant and Anti-Inflammatory Activities of Nano-Selenium Using Anoectochilus burmannicus Extract as a Potential Novel Functional Ingredient. *Nutrients*, 15(4), 1018.
- [52] ÇETİN, S., KILINBOZ, Y. F., KAPLAN, A. B. U., & ÇETİN, M. (2023). Development and In Vitro Characterization of Nanoemulsion Containing Hippophaes Rhamnoides Aqueous Extract. *International Journal of Pharm ATA*, 3(2), 31-34.
- [53] Dundar, A. N., Ozdemir, S., Uzuner, K., Parlak, M. E., Sahin, O. I., Dagdelen, A. F., & Saricaoglu, F. T. (2023). Characterization of pomegranate peel extract loaded nano phytosomes and the enhancement of bio-accessibility and storage stability. *Food Chemistry*, 398, 133921.
- [54] Aldayel, T. S., M. Badran, M., H. Alomrani, A., AlFaris, N. A., Z. Altamimi, J., S. Alqahtani, A. & Orfali, R. (2023). Chitosan-Coated Solid Lipid Nanoparticles as an Efficient Avenue for Boosted Biological Activities of Aloe perryi: Antioxidant, Antibacterial, and Anticancer Potential. *Molecules*, 28(8), 3569.
- [55] Dua, T. K., Giri, S., Nandi, G., Sahu, R., Shaw, T. K., & Paul, P. (2023). Green synthesis of silver nanoparticles using Eupatorium adenophorum leaf extract: characterizations, antioxidant, antibacterial and photocatalytic activities. *Chemical Papers*, 1-10.
- [56] Aftab, A., Ahmad, B., Bashir, S., Rafique, S., Bashir, M., Ghani, T., & Sajini, A. A. (2023). Comparative study of micro scale and macro scale technique for encapsulation of Calotropis gigantea extract in metal-conjugated nanomatrices for invasive ductal carcinoma. *Scientific Reports*, 13(1), 13474.