

Local Isolate Domination In Graphs

S.V.Padmavathi¹

¹P.G AndResearch Department Of Mathematics
Saraswathi Narayanan College, Perungudi Madurai-625 022,
Tamilnadu, India.

Abstract

In this Communication a new parameter called Local Isolate Domination in graphs is defined and Studied. A Dominating Set S is a Local Dominating set iff for each u in S , $\langle N(u) \rangle$ has an isolate.

AMSSubjectCode:05C69

Key Words: Dominating Set, Isolate Dominating Set, Local Isolate Dominating Set, Local Isolate Domination Number.

1 Introduction

Throughout this paper , Simple Finite Graphs without loops and multiple edges are considered. For terminologies and notations refer Chartrand And Lesniak[3].

Domination and related topics are dealt in [1 , 4, 5, 6].

A subset S of the vertex set $V(G)$ is a Dominating set if each vertex in the set $V \setminus S$ is adjacent to a vertex in S . Minimum cardinality of a minimal dominating set is the Domination number of a Graph denoted by $\gamma(G)$.

It is an isolate dominating set if the induced graph $\langle S \rangle$ contains an isolate and is introduced and studied in [7].

A Dominating set S is called a Doubly Isolate Dominating set if both the induced graphs $\langle S \rangle$ and $\langle V \setminus S \rangle$ have isolates. Doubly Isolate Dominating set is introduced and studied in [2].

*Corresponding Author.E-Mail Address: svsripadhma@gmail.com

Also when the concept of Isolate Dominating set is localized to the Neighbour set we arrive at a new variant called Local Isolate Domination in Graphs. This motivated us to define a new parameter that is introduced and studied in this communication.

2 Preliminary Results

Theorem 2.1 “For a Graph G with order atleast 3, $\Delta(G) = n-1$ and minimum degree atleast 2, G has no Local Isolate Dominating Set.”

Proof: From the hypothesis we observe that G is a graph without isolates and with Domination number as one and hence this dominating set is not a Local Isolate Dominating set of G . “Suppose S is any dominating set of G and $S \setminus \{v\} \neq \emptyset$ where $\{v\}$ is a full degree vertex of G . Now for each u in $\{S \setminus \{v\}\}$ the induced graph $\langle N(u) \rangle$ has no isolated vertex. Hence S is not a Local Isolate Dominating Set of G .

Corollary 2.2 “The Local Isolate Dominating Set Does Not Exist For The Following Graphs:

- (i) Complete Graph K_n .
- (ii) Wheel Graph W_n .
- (iii) Fan Graph F_n .”

Observation 2.3 “The Local Isolate Dominating Set Does Not Exist For Complete r -Partite K_{n_1, n_2, \dots, n_r} , $r \geq 3$ Graph.”

3. Main results

Proposition 3.1

“(i) For the Paths P_n and the Cycles C_n we have $\gamma_{lo}(P_n) = \gamma_{lo}(C_n) = \left\lceil \frac{n}{3} \right\rceil$, $\Gamma_{lo}(P_n) = \left\lceil \frac{n}{2} \right\rceil$

and $\Gamma_{lo}(C_n) = \left\lceil \frac{n}{2} \right\rceil$

(ii) If G is a Graph of Order n , Then $\gamma(G^+) = \Gamma_{lo}(G^+) = n$, Where G^+ is the Graph that was produced from G by attaching at each of G 's vertex's e Edges.”

Proof.(i) “Obviously $\gamma_{lo}(P_4) = 2$ and when $n \neq 4$, any γ -set of P_n is a local isolate dominating set as well, so that $\gamma_{lo}(P_n) \leq \gamma(P_n)$.” Every Local Isolate D dominating set is

a dominating set so $\gamma(P_n) \leq \gamma_{lo}(P_n)$ thus $\gamma_{lo}(P_n) = \gamma(P_n)$ and so $\gamma_{lo}(P_n) = \left\lceil \frac{n}{3} \right\rceil$ as

$\gamma(P_n) \left\lceil \frac{n}{3} \right\rceil$. Now if $P_n = \{v_1, v_2, v_3, \dots, v_n\}$ then the set $s = \left\{ v_{2i-1} \mid 1 \leq i \leq \left\lceil \frac{n}{2} \right\rceil \right\}$ is a minimal

isolate dominating set so that $\Gamma_{lo}(P_n) \geq \left\lceil \frac{n}{2} \right\rceil$. Additionally, since any set that contains

more than vertices of P_n is no longer able to be a minimal isolation dominating, we

have $\Gamma_{lo}(P_n) = \left\lceil \frac{n}{2} \right\rceil$. Similar to this, one may prove $\gamma_{lo}(C_n) = \left\lceil \frac{n}{3} \right\rceil$ and $\Gamma_{lo}(C_n) = \left\lceil \frac{n}{2} \right\rceil$

(ii) Each pendant vertex is required to be present in any minimal isolate-dominating set S of G^+ or one of its neighbours, in order to have at least n vertices. “Further, if $|S| > n$, S must consequently include a pendant vertex along with its support and so $S - \{v\}$, where v is the support, is an isolate dominating set of G^+ , a contradiction to the minimality of S .” Hence $|S| = n$.

Theorem 3.2 “For a Graph G of order at least 2, $\gamma_{lo}(G) = 1$ iff there exists a pair $u, v \in V(G)$, $\deg_G(u) = 1$ and $\deg_G(v) = n - 1$.”

Proof: Let G be a graph with $n \geq 2$. Suppose $\gamma_{lo}(G) = 1$. Let $S = \{v\}$ be a Local Isolate Dominating set of G . “Since S is a dominating set and $|V(G) \setminus S| = n - 1$, $\deg_G(v) = n - 1$.” Also since S is a γ_{lo} -set of G , $\langle n(v) \rangle$ has an isolate vertex, say u . Therefore u is a pendent vertex of G . Hence $\deg_G(u) = 1$. Conversely, $\{V\}$ is a dominant set of G since there is a vertex v with $\deg_G(v) = n$. Since “ $\deg_G(u) = 1$, u is a isolate vertex in $\langle n(v) \rangle$ ”, thus $\gamma_{lo}(G) = 1$.

Corollary 3.3 “For a Star Graph S_n with $n \geq 2$, $\gamma_{lo}(S_n) = 1$.”

Theorem 3.4 “If G is a Tree with $n \geq 2$ then G has a Local Isolate Dominating Set.”

Proof : Let G be a Tree of Order $n \geq 2$ and S be any Dominating Set of G . “Suppose G has no Local Isolate Dominating Set, there exist a vertex $v \in S$, and $\langle n(v) \rangle$ has no isolate vertex.” Thus $\langle n(v) \rangle$ is a connected Graph. This implies $\langle n[v] \rangle$ contains a cycle, which contradicts that G is a Tree. Therefore G has a Local Isolate Dominating Set.

Corollary 3.5 “For any Tree T , $\gamma(T) = \gamma_o(T) = \gamma_{lo}(T)$ ”.

Theorem 3.6 “Let S be any Local Isolate Dominating Set of a Graph G and $U \in S$. Then there exist a vertex $v \in V(G)$ such that $uv \in E(G)$ and $N(U) \cap N(V) = \emptyset$.”

Theorem 3.7 “For a Complete bipartite Graph $K_{m,n}$, $\gamma_{lo}(K_{m,n}) = 2$, $m \geq 2$, $n \geq 2$.”

Theorem 3.8 “A Local isolate dominating set S of a graph G is minimal iff it is 1-minimal.”

proof : “Let S be a 1-minimal Local Isolate Dominating Set of a graph G . Suppose there exists a $S' \subset S$ that is also a Local Isolated Dominating Set of G , then for all v in S' , $\langle n(v) \rangle$ has an isolate vertex”. Since S' is a Dominating Set, for all vertex in u in $S \setminus S'$ is adjacent to at least one vertex in S' and either u is an isolate vertex in $\langle n(v) \rangle$, $v \in S'$ or $\langle n(v) \rangle$ has an isolated vertex in $V \setminus S$.

case(i): u is an isolate vertex in $\langle n(v) \rangle$, $v \in S'$ then $S' \setminus \{v\}$ is Local Isolate Dominating Set of G which contradicts the “1-minimality of S ”.

case(ii): $\langle n(v) \rangle$ has an isolated vertex in $V \setminus S$. Let $w \in \langle n(v) \rangle$ be isolate vertex in $V \setminus S$ then $S \setminus \{u\}$ is Local Isolate Dominating Set of G which contradicts the 1-minimality of S . Hence S is minimal.

Converse is obvious.

Theorem 3.9 “A Local Isolate Dominating Set S of a Graph G is Minimal iff every vertex in S has a Private Neighbor with respect to S .”

Corollary 3.10 “A minimal Local Isolate Dominating Set S of a Graph G is also a minimal Dominating Set of a Graph G .”

4. Join of Graphs

Observation 4.1 “Let G and H be any two Graphs of order $m, n \geq 3$ with isolate vertex and S be a Local Isolate Dominating Set of $G+H$. Then $S \cap V(G) \neq \emptyset$ and $S \cap V(H) \neq \emptyset$.”

Theorem 4.2 “Let G and H be any two Graphs. Then S a subset of $V(G+H)$ is a Local Isolate Dominating Set of $G+H$ iff G and H have isolated vertices.”

Proof “Let G and H be any two Graphs and $S \subseteq V(G+H)$ be a Local Isolate Dominating Set of $G+H$.” Suppose G and H have no isolated vertex then for each $u \in S$, $\langle n(u) \rangle$ is connected, which is a contradiction. Therefore there are isolated vertex in both G and H .

Conversely, U and V be isolated vertices of G and H respectively, “Then $S = \{u, v\}$ is a Dominating Set of $G + H$ and also $n(u) \geq V(H)$ and $n(v) \geq V(G)$.” Thus $\langle n(u) \rangle$ and $\langle n(v) \rangle$ have isolated vertex. Therefore “ S is a Local Isolate Dominating Set of $G+H$.”

Corollary 4.3 “Let G and H be any Graphs with isolated vertex, Then $\gamma_{1o}(G+H) \leq 2$.”

Proof. “Let G and H be Graphs with isolated vertex.” Suppose either $G=K_1$ or $H=K_1$ or $G=H=K_1$ Then Clearly, $\gamma_{1o}(G+H) = 1$. Suppose $G \neq K_1$ and $H \neq K_1$, by the theorem 4.2, $\gamma_{1o}(G+H) = 2$. Thus $\gamma_{1o}(G+H) \leq 2$.

References

- [1] Benjier H. Arriola, Isolate Domination In The Join And Corona Of Graphs, *Appl.Math.Sci.*9(2015)1543-1549.
- [2] Benjier H. Arriola, Doubly Isolate Domination In Graphs, *International Journal Of Mathematical Analysis*, Vol. 9, 2015, No. 57, 27932798.
- [3] G. Chartrand, Lesniak, *Graphs And Digraphs*, Fourthed., Crcpress, Boca Raton, 2005.
- [4] T.W. Haynes, S.T. Hedetniemi, P.J. Slater, *Domination In Graphs: Advanced topics*, Marcel Dekker, Newyork, 1998.
- [5] T.W. Haynes, S.T. Hedetniemi, P.J. Slater, *Fundamentals Of Domination In Graphs*, Marcel Dekker, New York, 1998.
- [6] S.T. Hedetniemi, R. Laskar(Eds.), *Topics In Domination In Graphs*, *Discrete Math.* 86 (1990).
- [7] I. Sahulhamid, S. Balamurugan, *Isolate Domination In Graphs*, *Arab.J.Math Sci*22 (2016) 232-241.

About the Author(s)



Dr.S.V.Padmavathi, is currently an Associate Professor in PG and Research Department of Mathematics, Saraswathi Narayanan College (Autonomous), Madurai. Earlier she has experience as Assistant Professor in Lady Doak College and Fatima College, Madurai. She has 20 years of teaching experience. She completed her Ph.D degree at Saraswathi Narayanan College under UGC-JRF-NET scholarship in the year 2008. She received her M.Phil (Mathematics) degree at Madurai Kamaraj University, Madurai and completed M.Sc (Mathematics) degree at Lady Doak College, Madurai. Her area of research is Graph Theory. She published 12 papers in refereed journals. Participated and presented number of papers in various National and International conferences or seminars in Mathematics. More than 45 M.Phil Scholars have been awarded under her guidance. Currently 4 Ph.D Scholars are pursuing research.