

Introduction

The most popular method of treating pulp necrosis is root canal treatment (RCT). However, it neither restores the vitality nor the neovascularity of the pulpal tissue. Therefore, with the advancement of tissue engineering, regenerative endodontic therapy (RET) has garnered more interest^[1]. RET uses tissue engineering to revascularize, innervate, and restore the odontoblastic layers and fills the cleansed canals with vital tissue as opposed to biocompatible, nonvital foreign substance eg. gutta percha being used in root canal therapy^[2,3].

Definition

Revascularization - The engraftment of regenerated pulpal tissue to host vasculature in root canals or invagination of undifferentiated periodontal cells from the apical region in immature teeth^[4].

Regeneration - The replacement of damaged tissue by cells identical to the lost tissue, leading to the complete reestablishment of biological function that is, both revascularization and revitalization, including dentin, root structure and pulp-dentin complex^[5].

Regenerative Endodontics (according to the American Association of Endodontists) - biologically based procedures designed to replace damaged structures, including dentin and root structures, as well as cells of the pulp–dentin complex^[6].

Scaffolds




Figure 1: Various platelet products

Scaffolds are essential to regenerative endodontic therapy because they create an environment that encourages the stem cell population to migrate, proliferate and undergo differentiation^[7]. Blood clot revascularization is the most commonly used technique in which a K or H file is inserted beyond the apical foramen, thus inducing a periapical bleed resulting in a blood clot capable of serving as a scaffold^[8].

Platelet products are commonly being used now because concentrated autologous

platelets and growth factors can be extracted from the plasma and they

replicate the final stage of the coagulation cascade. The inclusion of leukocytes was done for their immunological and antimicrobial capabilities, role in the healing of wounds by control of regional factors, release of growth factors and promoting cell division^[9]. Variable centrifugal time periods and rate, chemical alterations, and choice of supernatants and precipitates have led to the synthesis of

several formulations of fibrin network configurations, white blood cell content and cytokines, resulting in different physical and physiological properties of platelet products and its applications ^[10] [FIGURE 1].

Stem cells –

Stem cells are an intriguing biological foundation for regenerative therapies due to their capacity for self-renewal. Mesenchymal populations are assumed to be the source of dental stem cells. The most




Figure 2:Steps to obtain DPSC

prevalent dental stem cells are: Dental pulp stem cells (DPSC), stem cells from exfoliated deciduous teeth (SHED), stem cells from apical papilla (SCAP) and those from periodontal ligament. From the standpoint of endodontic regeneration, DPSC are multipotent cells that can transform into osteoblasts, adipocytes, and neural cells ^[11]. Its applications are supported by their inherent role in the generation of odontoblasts that produce reparative dentin. Transplantation of DPSC acquired from teeth with irreversible pulpitis necessitating root canal therapy is another method to achieve

effective pulp regeneration in the clinical outlook. The simplicity with which DPSCs may

Clinical considerations for regenerative endodontics –




Figure 3: Case selection

Case selection is crucial when selecting treatment procedure for a specific pulpal disease [FIGURE 3].

Criteria for RET:

1. Degree of infection in the canal - If the canals are effectively disinfected (using triple antibiotic paste (TAP), double antibiotic paste (DAP), calcium hydroxide or formocresol) and the coronal access is effectively sealed, regenerative endodontic treatment should occur.
2. Apex diameter - 1.1 mm or larger (Now a days, RET is also being performed on teeth with a mature apex)

3. Patients age – 8 to 16 years ^[13] [FIGURE 4]

Specific disinfection and irrigation protocols are required in order to entirely eliminate tissue remnants and bacteria in the root canal. ^[14].

TAP consisting of ciprofloxacin, metronidazole, and minocycline was suggested by Hoshino et al. in 1996 to decontaminate the canal and can successfully be used in RET ^[15]. Contrastingly, Chueh et al. demonstrated calcium hydroxide to completely cleanse the canal and promote regeneration ^[16]. Bose et al. conducted a global analysis of RET cases in 2009 which concluded that RET with the TAP, calcium




Figure 4: Standard protocol for RET according to American Association of Endodontics

hydroxide, and formocresol significantly increased root length and width. However, TAP substantially increased dentin wall thickness without altering root width ^[17].

Drawbacks of TAP:

According to Ding et al., two patients experienced pain following the administration of TAP and thus, were not included in the study. As reported by Jung et al., utilising TAP resulted in a chronic sinus tract and when switched out for calcium hydroxide, and the patient's problems subsided ^[18,19]. Minocycline, one of the ingredients in TAP, is the main

contributor to tooth discolouration, claim Kim et al. Through dentinal tubules, minocycline can enter the tooth and combine with the tooth's crystal structure. Before applying the TAP, the coronal dentin wall can be sealed with flowable composite resin. The other procedure is retrograde antibiotic filling with a 20 gauge needle to reduce contact with the coronal area of the tooth. According to case studies, cefaclor and amoxicillin can be substituted for minocycline. However, amoxicillin can cause allergic reactions as a side effect ^[20].

Double antibiotic paste is thought to be an efficient medication for removing harmful germs from root canals since it contains both metronidazole and ciprofloxacin. The problem of tooth discolouration brought on by the presence of minocycline in the TAP formulation is currently being addressed in endodontics with DAP ^[21].

Other methods of regenerative endodontics -

- **Pulp autotransplantation -**

The ideal "scaffold" for DPSC differentiation in their natural habitat might be produced by transplanting the complete pulp. Additionally, majority of the nerves and vessels are already developed, which facilitates the revascularization of grafted pulp tissue. Overall, this approach modality might be achieved in clinical settings without the necessity for an extension of the in vitro DPSC laboratory [22].

- **Cell homing technique -**

Cell homing technique is based upon the biological principles of typical tissue wound healing. Cell recruitment and differentiation are two discrete biological processes that need to take place. Over-instrumentation results in bleeding, which is used to start cell homing therapy. Endogenous cells and growth factors required for RET are present in the blood clots that follow, filling the pulp space, as well as a biological structure supporting cellular activity [23].

Histological aspect of RET -

Investigations on animals showed that platelet concentrates do not produce healthy, sound pulp tissue; instead, they generate bone-like tissue in the root canal [24]. Numerous histological investigations regarding regenerative endodontic treatments have demonstrated that the tissues developed in the root canal space were mineralized akin to cementum and bone as well as fibrous connective tissue similar to periodontal ligament [25]. Without histological proof, radiographic thickening of the canal walls and ongoing root maturation of young permanent teeth with necrotic pulps following regenerative endodontic operations should not be interpreted as the regeneration of dentin-pulp complex [26].

Conclusion –

The goal of tissue engineering via regenerative endodontics is to re-generate the pulp-dentin complex of immature young permanent teeth thereby resuming root formation. RET has been able to meet the main objectives of root canal therapy that are: to treat apical periodontitis in addition to eradicating the patient's clinical signs, symptoms and indicators. Although further development of the root is unpredictable, RET has the prospective to encourage further root formation as opposed to apexification.

Instead of pulp-like tissue, RET has shown to grow periodontal-like tissues in the canal space. Despite the vitality of the tooth being restored, the biological functions of dental pulp cease to exist. Nevertheless, young permanent teeth with necrotic pulpal tissues can be effectively and consistently treated with RET, resulting in patient-focused results like tooth preservation, symptom-free conditions, and appealing aesthetics.

References -

1. Kim SG, Malek M, Sigurdsson A, Lin LM, Kahler B. Regenerative endodontics: a comprehensive review. *Int Endod J*. 2018;**51**(12):1367–1388. doi: 10.1111/iej.12954.
2. Adnan S, Ullah R. Top-cited articles in regenerative endodontics: a bibliometric analysis. *J Endod*. 2018;**44**(11):1650–1664. doi: 10.1016/j.joen.2018.07.015.
3. Saoud TM, Martin G, Chen YH, et al. Treatment of mature permanent teeth with necrotic pulps and apical periodontitis using regenerative endodontic procedures: a case series. *J Endod*. 2016;**42**(1):57–65. doi: 10.1016/j.joen.2015.09.015.
4. Kumar V, Abbas AK, Fausto N, et al. Robbins and cotran pathologic basis of disease. Philadelphia,PA: Saunders, 9th ed. 2014.
5. Majno G, Joris I. Cells, tissues, and disease. Oxford, London: Oxford University Press, 2nd ed. 2004.
6. Murray P.E., Garcia-Godoy F., Hargreaves K.M. Regenerative Endodontics: A Review of Current Status and a Call for Action. *J. Endod*. 2007;**33**:377–390.
7. Torabinejad M, Turman M. Revitalization of tooth with necrotic pulp and open apex by using platelet-rich plasma: a case report. *J Endod*. 2011;**37**(2): 265–8. <https://doi.org/10.1016/j.joen.2010.11.004>.
8. Murray PE. Platelet-rich plasma and platelet-rich fibrin can induce apical closure more frequently than blood-clot revascularization for the regeneration of immature permanent teeth: a meta-analysis of clinical efficacy. *Front Bioeng Biotechnol*. 2018;**6**:139. <https://doi.org/10.3389/fbioe.2018.00139>.
9. Sushma Naag, Abhishek Savirmath, Butchi Babu Kalakonda, Uday Kiran Uppada, Shekar Kamisetty, et al. (2015) Platelet concentrates: Bioengineering dentistry's regenerative dreams. *J Dent Res Rev* 2(2): 23-29.
10. Dohan Ehrenfest DM, Pinto NR, Pereda A (2018) The impact of the centrifuge characteristics and centrifugation protocols on the cells, growth factors, and fibrin architecture of a leukocyte – and platelet – rich fibrin (L-PRF) clot and membrane. *Platelets* 29(2): 171 – 184.
11. Huang GT, Gronthos S, Shi S. Mesenchymal stem cells derived from dental tissues vs. those from other sources: their biology and role in regenerative medicine. *J Dent Res* 2009;**88**:792–806.
12. Tatullo M, Marrelli M, Shakesheff KM, White LJ. Dental pulp stem cells: function, isolation and applications in regenerative medicine. *J Tissue Eng Regen Med*. 2015 Nov;**9**(11):1205-16. doi: 10.1002/term.1899. Epub 2014 May 21. PMID: 24850632.
13. Garcia-Godoy F, Murray PE. Recommendations for using regenerative endodontic procedures in permanent immature traumatized teeth. *Dent Traumatol* 2012;**28**:33-41.

14. Mohammadi Z, Jafarzadeh H, Shalavi S, Yaripour S, Sharifi F, Kinoshita JI. A Review on Triple Antibiotic Paste as a Suitable Material Used in Regenerative Endodontics. *Iran Endod J*. 2018 Winter;13(1):1-6. doi: 10.22037/iej.v13i1.17941. PMID: 29692827; PMCID: PMC5800433.
15. Hoshino E, Kurihara-Ando N, Sato I, Uematsu H, Sato M, Kota K, Iwaku M. *In-vitro* antibacterial susceptibility of bacteria taken from infected root dentine to a mixture of ciprofloxacin, metronidazole and minocycline. *Int Endod J* 1996;29:125-130.
16. Chueh LH, Ho YC, Kuo TC, Lai WH, Chen YH, Chiang CP. Regenerative endodontic treatment for necrotic immature permanent teeth. *J Endod* 2009;35:160-164.
17. Bose R, Nummikoski P, Hargreaves K. A retrospective evaluation of radiographic outcomes in immature teeth with necrotic root canal systems treated with regenerative endodontic procedures. *J Endod* 2009;35: 1343-1349.
18. Ding RY, Cheung GS, Chen J, Yin XZ, Wang QQ, Zhang CF. Pulp revascularization of immature teeth with apical periodontitis: a clinical study. *J Endod* 2009;35:745-749.
19. Jung IY, Lee SJ, Hargreaves KM. Biologically based treatment of immature permanent teeth with pulpal necrosis: a case series. *J Endod* 2008;34:876-887.
20. Kim JH, Kim Y, Shin SJ, Park JW, Jung IY. Tooth discoloration of immature permanent incisor associated with triple antibiotic therapy: a case report. *J Endod* 2010;36:1086-1091.
21. Sabbagh, Sedigheh & Sarraf Shirazi, Alireza & Torabzadeh, Hassan. (2018). Double Antibiotic Paste for Management of External Inflammatory Root Resorption. *Iranian Endodontic Journal*. 13. 569-572. 10.22037/iej.v13i4.22893.
22. Nakashima M, Iohara K, Bottino MC, et al. Animal models for stem cell-based pulp regeneration: foundation for human clinical applications. *Tissue Eng Part B Rev* 2019;25:100–13.
23. Yan, Hongji et al. "Regenerative Endodontics by Cell Homing: A Review of Recent Clinical trials." *Journal of endodontics* vol. 49,1 (2023): 4-17. doi:10.1016/j.joen.2022.09.008
24. Meza G, Urrejola D, Saint Jean N, et al. Personalized cell therapy for pulpitis using autologous dental pulp stem cells and leukocyte platelet-rich fibrin: a case report. *J Endod* 2019;45:144–9.
25. Lei L, Chen Y, Zhou R, Huang X, Cai Z. Histologic and immunohistochemical findings of a human immature permanent tooth with apical periodontitis after regenerative endodontic treatment. *J Endod*. 2015. July;41(7):1172–9. 10.1016/j.joen.2015.03.012
26. Lin LM, Kahler B. A review of regenerative endodontics: current protocols and future directions. *J Istanbul Univ Fac Dent*. 2017 Dec 2;51(3 Suppl 1):S41-S51. doi: 10.17096/jiufd.53911. PMID: 29354308; PMCID: PMC5750827.