

Cross-kingdom RNA interference: A novel strategy for plant disease management

Gopal Chowdhury
Dept. of Plant Pathology
Bidhan Chandra Krishi Viswavidyalaya
Nadia, West Bengal, India
gchowdhury.bckv@gmail.com

Abstract

Small RNAs (sRNAs) are short non-coding RNA molecules that play a crucial role in regulating gene expression. They include microRNAs (miRNAs) and small interfering RNAs (siRNAs) and are involved in controlling plant immunity against pathogens and pests. Recent research has revealed that sRNAs can move between different species and silence genes, a process called cross-kingdom RNA interference (RNAi). This discovery has led to the development of host-induced gene silencing (HIGS), where transgenic plants are engineered to produce double-stranded RNA (dsRNA) that targets specific pathogen genes. This RNA-based approach offers a promising alternative to agrochemicals and transgenic crops, which have potential health and environmental risks. By applying dsRNAs or sRNAs directly onto host plants or post-harvest products, a technique known as spray-induced gene silencing (SIGS), effective disease management can be achieved without the need for transgenic methods. This review explores recent advancements in using cross-kingdom RNAi for managing plant diseases.

Key words- Cross kingdom RNA interference, small interfering RNA, microRNAs (miRNAs), host-induced gene silencing, spray-induced gene silencing, disease management.

I. Introduction

Small RNAs, including small interfering RNAs (siRNAs) and microRNAs (miRNAs), are a class of non-coding RNAs that typically ranges from 20 to 30 nucleotides in length. They play important roles in regulating gene expression [1]. siRNAs, which are about 20-24 nucleotides long, are double-stranded RNA molecules produced from longer precursor molecules derived from the organism's genome or external sources like viruses and transgenic transcripts [2-3]. Conversely, miRNAs are single-stranded non-coding RNAs that are approximately 20 to 22 nucleotides long. They are derived from primary miRNAs (pri-miRNAs), which are mainly transcribed from regions located between protein-coding genes. These pri-miRNAs have a distinct stem-loop structure [4-5].

The regulation of plant immunity against pests and pathogens (such as fungi, bacteria, oomycetes, and viruses) relies significantly on small RNA-mediated RNA interference (RNAi). Additionally, sRNAs originating from pests and diseases play a crucial role in monitoring their virulence. While it was previously understood that sRNAs could move within the cells and tissues of a single organism, recent evidence supports the idea that some of these sRNAs can transfer across different species, leading to the silencing of genes in the recipient organism. This phenomenon is known as 'cross-kingdom RNAi.' By targeting the genes responsible for pathogenicity in eukaryotic pathogens and pests, sRNAs produced within the plant system can effectively suppress them. This discovery has paved the way for the development of RNA-based pesticide formulations that are highly specific and adaptable for controlling multiple diseases simultaneously using cross-kingdom RNAi.

Certain pathogens possess the ability to uptake RNAs from the environment, a process known as 'environmental RNAi' [6]. Leveraging this knowledge, host-induced gene silencing (HIGS) technology has emerged, enabling the evolution of a recent approach to plant protection called transgene-mediated cross-kingdom RNAi or 'spray-induced gene silencing' (SIGS). In SIGS, dsRNAs and sRNAs targeting the genes of pathogens are sprayed onto plant surfaces to suppress their virulence [7]. This review provides a comprehensive summary of our current understanding and the practical applications of cross-kingdom RNAi.

To sum up, the current understanding of cross-kingdom RNAi has revealed the potential of sRNAs in regulating gene expression across different organisms. This knowledge has led to the development of innovative

RNA-based pesticide formulations and approaches like SIGS, which offer effective means of controlling pests and diseases in the realm of agriculture.

II. The cross-kingdom RNA interference

RNAi is a universally observed biological mechanism that controls the expression of protein-coding genes in response to the presence of double-stranded RNA (dsRNA). Evidence suggests that RNAi can be both pathogen as well as host induced. Naturally in most of the plant-microbe interactions the cross-kingdom RNAi occurs [8]. Pathogenic small RNAs play a role in cross-kingdom RNAi within plant hosts, serving as a unique type of pathogen effectors that hinder host immunity to facilitate successful infection [9-10]. Cross-kingdom transfer of small RNA (sRNA) and the subsequent silencing phenomenon are observed in various plant-patho systems, including interactions with fungal pathogens such as *Botrytis cinerea*, *Verticillium dahliae*, *Blumeria graminis*, *Sclerotinia sclerotiorum*, and *Phytophthora infestans*.

Botrytis cinerea, an important fungal pathogen causing pre and as post-harvest diseases in many crops delivers sRNAs (Bc-sRNAs) that bind to the host argonaute-RISC component 1 (AGO1) gene and hijack the host RNAi pathways and suppresses the immunity system [8,11]. Furthermore, the sRNA Bc-siR37 derived from *Botrytis cinerea* has the ability to suppress the plant's defense mechanisms by reducing the expression of several target genes within the host [12]. *Sclerotinia sclerotiorum* produces at least 374 distinct sRNAs during infection which target the functional domains associated with immunity of *Arabidopsis thaliana* [13]. Zhang et al. [14] showed that when cotton plants are infected by *Verticillium dahliae*, they enhance the production of microRNA166 (miR166) and miR159, which are then transported to the fungal hyphae to facilitate targeted silencing. The expression of two critical genes in *Verticillium dahliae*, namely Clp-1 (encoding a Ca²⁺-dependent cysteine protease) and HiC-15 (encoding an isotrichodermin C-15 hydroxylase), which are essential for the pathogenicity of the fungus, is specifically inhibited by miR166 and miR159, respectively. In experimental conditions, the silencing of these pathogenic genes by small RNAs produced by the host plant has been demonstrated to limit fungal growth, as observed in powdery mildew studies [15]. On the other hand, the oomycete pathogen *Phytophthora infestans* employs RxLR-type effectors that possess RNA silencing suppressor capabilities [16-17]. Hence, the cross-kingdom transfer and silencing of RNA appear to play a crucial role in the pathogenicity of various fungal and filamentous pathogens, as well as in plant resistance mechanisms.

III. Formation of small RNAs in plants

In the plant system, miRNAs are produced from hairpin-shaped precursors that contain 64 to 303 nucleotides, whereas in animals, miRNA precursors are usually 60 to 70 nucleotides in size [18-19]. These miRNAs are generated from specific genes called MIR genes. The first step involves the synthesis of primary miRNAs (pri-miRNAs) by an enzyme called RNA polymerase II (RNA pol II) within the nucleus. The processing of pri-miRNAs into mature miRNA duplexes occurs through the action of Dicer-like 1 (DCL1), a ribonuclease III-like enzyme, along with the involvement of Histonically Nucleated Leaves 1 (HYL1) protein. The resulting miRNA duplex consists of two strands: miRNA and miRNA*. The miRNA* strand possesses a 2-nucleotide overhang at its 3' end [20]. The miRNA duplex is methylated by a methyl-esterase known as Hua Enhancer-1 (HEN1), which is present in both plants and animals [21]. Methylation plays a crucial role in protecting RNA from degradation by exonucleases and from uridylation, which involves the addition of a short poly-U tail to unmethylated miRNAs. This uridylation decreases the stability of miRNAs, making them more prone to decay. Once the miRNA duplex matures, it is transported to the cytoplasm with the assistance of a nuclear membrane protein known as HASTY (HST) [22-25]. Following that, an unidentified helicase protein unwinds the miRNA-miRNA* duplex, exposing the mature miRNA to the RNA-induced silencing complex (RISC). The mature miRNA, bound to RISC containing the AGO protein, then either cleaves or represses the translation of specific target messenger RNAs (mRNAs) by guiding the RISC to the target mRNA [26-27].

siRNAs and miRNAs have a close connection but differ in their origin, structure, associated effector proteins, and mechanisms of action [4, 28]. There are two types of siRNAs: Ta-siRNAs and Ra-siRNAs, which vary in their precursor molecules and synthesis steps. Ta-siRNAs originate from non-coding RNA (ncRNA) precursors, whereas Ra-siRNAs are produced from transposable elements and repetitive sequences. The synthesis of ssRNA precursor by miRNA-dependent processing represents the most distinctive aspect of Ta-siRNA biogenesis [29-31] on the other hand, ssRNA for Ra-siRNAs is synthesised by DNA-dependent RNA polymerase IV from the heterochromatic locus. Later, RNA-dependent RNA polymerases (RDRPs) duplicate the ssRNA

precursors in both siRNAs to produce dsRNA precursors. The rest of the process of siRNA synthesis is the same as that of miRNA, with the exception of the types of enzymes and proteins involved [32].




Figure 1. Mechanism of miRNAs synthesis in plants

IV. Cross kingdom movement of sRNAs

The movement of sRNAs within organisms or between different kingdoms is crucial for regulating the expression of specific target genes. This movement allows the transmission of RNA-silencing signals from one cell type to another, even across long distances, thereby influencing gene expression in distant cells and tissues. This phenomenon is referred to as "systemic acquired silencing" (SAS). It has been extensively studied and documented [33-35]. The sRNA population involved in this mobile transfer often exhibits differences compared to the total sRNA population within cells, indicating that there is a selective process in place during the transfer to ensure the inclusion of potentially functional sRNAs [36-38]. This selection process may involve various factors such as RNA-binding proteins (RBPs), AGOs, packaging into the extracellular vesicles (EVs), or other mechanisms of transport [9]. Recently, a gating mechanism has been discovered at specific cell-cell interfaces that precisely regulate the transport of miRNAs, preventing their long-distance movement from shoot to root [39].

The movement of sRNAs within an organism, specifically between cells, is of utmost importance in governing gene expression within the organism, thereby playing a pivotal role in its regulatory mechanisms. However, the cross-kingdom mobility of these RNAs is of significant importance to both hosts and pathogens, as it influences resistance mechanisms in the host and pathogenicity in the pathogens.

A. Transfer of sRNAs from pathogens to host plants

Crop yield losses and their impact on global food security are primarily caused by plant pathogenic fungi. Plants are vulnerable to a wide range of fungal pathogens. In nature, resistance to these pathogens is the norm, while disease development is the exception. For a pathogen to successfully cause disease, it must possess sophisticated mechanisms to overcome the defense barriers of the host.

Pathogens have evolved mechanisms to manipulate host gene expression and evade defense mechanisms. One such mechanism is called pathogen-modulated RNA interference, where small RNA molecules originating from pathogens can act as effectors to alter host gene expression. These sRNAs can undermine the host's defense strategies by targeting specific genes. For example, a study by Wang et al. [40] found that a novel microRNA (miRNA) called Pst-miR1 in *Puccinia striiformis* f. sp. *tritici* targets the pathogenesis-related 2 (PR2) gene in wheat, potentially suppressing the host's defense strategy. Similarly, sRNAs derived from *Botrytis cinerea*, known as Bc-sRNAs, bind to Argonaute 1 (AGO1) and hijack the host RNAi machinery, leading to the selective silencing of host immunity genes [11]. This indicates that *B. cinerea* transfers virulent sRNA molecules into host plant cells

to suppress host immunity as a counter-defense strategy for successful infection. Furthermore, bidirectional interactions between pathogens and hosts have been identified. For instance, sRNAs produced by *Plasmopara viticola* trigger the cleavage of grapevine (*Vitis vinifera*) genes, while grapevine-derived sRNAs target *P. viticola* mRNAs [41]. These examples highlight the intricate interplay between pathogens and their host plants, where sRNAs play a crucial role in modulating gene expression to facilitate infection or counter host defense mechanisms.

B. Transfer of sRNAs from host plants to pathogens

Transfer of miRNAs between different kingdoms has been observed during interactions between hosts and pathogens, resulting in the inhibition of pathogen's invasive capabilities [42]. Plants encounter attacks from various pathogens, such as bacteria, fungi, mycoplasma, nematodes, viruses, viroids, and parasites, and have developed defense strategies to counteract them [43-44]. In a sophisticated manner, plants employ miRNA-mediated transcriptional or post-transcriptional silencing of pathogenic mRNA, specifically targeting virulence genes. This mechanism forms a part of the plant's resistance strategy against pathogens.

The host organisms possess RNA interference (RNAi) silencing machinery that can directly target the RNA genome and related transcripts of various pathogens like viruses, virus satellites, and viroids to regulate their accumulation [45]. This silencing process involves the transportation of specific small RNAs (sRNAs) from plants, including miRNAs, which induce gene silencing in pathogenic fungi, thereby enhancing disease resistance [46-47]. For example, in *Arabidopsis*, siRNAs can enter *Phytophthora* through extracellular vesicles during infection, contributing to the development of resistance [46]. Additionally, *Arabidopsis* miR166 has been observed to be exported to *V. dahliae* fungal hyphae, resulting in the suppression of pathogenicity [48]. In cotton plants infected with *V. dahliae*, the accumulation of miR166 and miR159 has been observed, which targets *V. dahliae* genes encoding a Ca²⁺ dependent cysteine protease (Clp-1) and an isotrichodermin C-15 hydroxylase (HiC-15), respectively [14]. Wheat plants restrict disease development by targeting specific genes of *Puccinia striiformis* f. sp. *tritici*, such as Calcineurin homologs Pscn1/Pscn1 and MAPK kinase gene PsFU27, resulting in reduced hyphal extension, decreased uredospore production, and induction of necrosis in plant cells [49-50]. Targeting the PsCK1 gene (a PKA catalytic subunit gene) of *P. striiformis* f. sp. *tritici* also leads to a reduction in disease progression, significantly reducing the length of infection hyphae and disease symptoms [50]. Various other interactions involving RNA and siRNA between plants and fungi have been summarized in Table 1, resulting in either resistance or disease outcomes.

Table 1: cross kingdom RNA interaction between plants and fungi

From	To	Target Genes	Involved sRNA	Reference
<i>B. cinerea</i>	<i>S. lycopersicum</i>	<i>MAPKKK4</i>	Bc-siR3.2	[11]
<i>B. cinerea</i>	<i>A. thaliana</i>	<i>PRXIIIF</i>	Bc-siR3.1	[11]
<i>B. cinerea</i>	<i>A. thaliana</i>	<i>WAK</i>	Bc-siR5	[11]
<i>B. cinerea</i>	<i>A. thaliana</i>	<i>MPK2</i> and <i>MPK1</i>	Bc-siR3.2	[11]
<i>G. hirsutum</i>	<i>V. dahliae</i>	<i>Clp-1</i>	miR166	[14]
<i>G. hirsutum</i>	<i>V. dahliae</i>	<i>Clp-1</i>	miR166	[14]
<i>P. striiformis</i> f. sp. <i>tritici</i>	<i>T. aestivum</i>	<i>PR2</i>	Pst-milR1	[40]
<i>B. cinerea</i>	<i>A. thaliana</i>	<i>WRKY7</i> , <i>PMR6</i> & <i>FEI2</i>	Bc-siR37	[40]
GRSPaV	<i>V. vinifera</i>	<i>VPS55</i>	vsR6978	[51]
GFkV	<i>V. vinifera</i>	<i>S2P</i> metalloprotease	vsR1378	[51]

V. Potential of cross kingdom RNAi application in plant disease management

The transfer of miRNAs between different kingdoms presents promising opportunities for the development of eco-friendly crop protection strategies. For instance, the transfer of miRNA159 and miRNA166 from cotton to the pathogenic fungus *Verticillium dahliae* has been demonstrated to provide resistance [15]. As mentioned earlier, numerous studies have highlighted the horizontal transfer of miRNAs among plants, animals,

and microbes. This transfer facilitates the regulation of gene expression in both host and pathogenic organisms, thereby contributing to the protection of crops in agricultural production. The primary objective of miRNA transfer from pathogens to hosts is to suppress plant defense mechanisms, acting as a counter-defense strategy employed by the pathogens. Wang et al. [12] demonstrated bidirectional RNAi, indicating that the cross-kingdom transfer of miRNAs can suppress the virulence of plant pathogens and protect crops. Another study by Wang et al. [12] revealed that fungal pathogens can uptake sRNAs from the environment, leading to a loss of virulence. For instance, the fungal pathogen *Botrytis cinerea*, responsible for grey mold disease, has been targeted using externally applied sRNAs and dsRNA through spraying on the surface of fruits, vegetables, and flowers, resulting in the suppression of fungal pathogenic genes and protection against plant infections. This discovery opens up new possibilities for managing plant diseases through spray-induced gene silencing (SIGS).

A. Spray induced gene silencing (SIGS) in plant disease management

Crop production currently relies heavily on chemical fungicides, which can leave pesticide residues and contribute to the development of resistance in pathogens. Therefore, there is a growing need for an alternative and environmentally friendly method to protect crops. The third green revolution in modern agriculture is on the horizon, and understanding reverse genetics in gene functional characterization holds significant potential for managing agricultural pests. The RNA interference (RNAi) technology offers a practical approach to plant protection, with spray induced gene silencing (SIGS) being the most effective method for crop protection [52]. SIGS has been observed to successfully silence target genes in viruses [53] and fungi [7, 54-55]. Notably, *Fusarium graminearum* and *Botrytis cinerea* have been effectively controlled using SIGS, which mimics host-induced gene silencing (HIGS) without requiring the modification of plant genomes. The sprayed RNAs can enter fungal cells through two possible pathways: either through plant cells and transferred to the pathogenic fungi, or directly taken up by fungal cells. These RNAs then exert their effects in two ways: by activating the plant's RNAi machinery in plant cells or by directly activating the fungal RNAi machinery in fungal cells. Koch et al. [54] demonstrated successful suppression of CYP51 gene in barley, providing resistance against *F. graminearum* through SIGS. They also demonstrated the effectiveness of silencing green fluorescent protein (GFP) expression in a GFP-expressing strain of *Fusarium graminearum* by spraying RNA fragments derived from jellyfish GFP onto barley leaves. This finding indicates that small RNA-induced gene silencing (SIGS) is not restricted to specific sequences and has the potential to target essential genes in different interacting pathogens. Additionally, McLoughlin et al. [55] found that spraying dsRNA can inhibit the growth of *Botrytis cinerea* and *Sclerotinia sclerotiorum* on *Brassica napus*.

SIGS offers a technologically simple and socially accepted approach to crop protection, as it does not involve the generation of genetically modified organisms (GMOs). It is considered safe for consumption and allows effective control of fungi even after harvesting [8, 56]. SIGS has the advantage of simultaneously targeting multiple pathogens, making it practical for managing pests and diseases that affect various crops [57]. Another benefit is that RNA degrades relatively quickly, so any potential accumulation in the environment is not expected to be hazardous, considering that nucleic acids are naturally present [58]. Therefore, RNAi-based pesticides offer an environmentally friendly alternative to conventional agrochemical products and genetically engineered plant solutions [12].

Table 2. SIGS in plant-pathogen interactions.

Host Plant	Target Pathogens	Target Genes	Concentration of dsRNA	Reference
<i>T. aestivum</i>	<i>F. asiaticum</i>	<i>Myosin 5</i>	0.1 pM	[59]
<i>A. thaliana</i>	<i>V. dahliae</i>	<i>DCL</i>	20 ng/μL	[8]
<i>A. thaliana</i> , etc.	<i>B. cinerea</i>	<i>DCL1/2</i>	20 ng/μL	[8]
<i>H. vulgare</i>	<i>F. graminearum</i>	<i>CYP51</i>	20 ng/μL	[54]
<i>B. napus</i>	<i>B. cinerea</i>	BC1G_04955, etc.	42 ng/μL	[55]
<i>B. napus</i>	<i>S. sclerotiorum</i>	SS1G_01703, etc.	20 ng/μL *	[55]

<i>A. thaliana</i>	<i>S. sclerotiorum</i>	SS1G_03208, etc.	20 ng/μL	[55]
--------------------	------------------------	------------------	----------	------

VI. Constraints

Although RNAi-based bio-fungicides hold great potential for managing pests and fungal pathogens, there are challenges that need to be overcome for their large-scale production and commercialization. One major challenge is the uptake efficiency of the bio-fungicides. Qiao et al. [60] showed that not all pathogens have the ability to effectively capture long and/or small double-stranded RNAs (dsRNAs), which is crucial for their control through RNAi. This factor plays a determining role in the success of RNAi-based bio-fungicides in controlling specific pathogens.

VII. Summary

Cross-kingdom RNA interference (RNAi) is a process in which small RNA molecules regulate gene expression between different organisms. In the context of plants and pathogens, small RNAs known as microRNAs (miRNAs) and small interfering RNAs (siRNAs) play a significant role. Pathogens can produce sRNAs to interfere with host immunity and facilitate infection, while host plants can produce sRNAs to target pathogen genes and suppress their virulence. The biogenesis of sRNAs in plants involves processing primary miRNAs (pri-miRNAs) into mature miRNAs and siRNAs, which then bind to the RNA-induced silencing complex (RISC) to degrade or repress target messenger RNAs (mRNAs). The movement of sRNAs within organisms or between kingdoms is crucial for gene regulation, allowing the RNAi signal to propagate over long distances. Factors such as argonaute proteins, RNA-binding proteins, and packaging into extracellular vesicles influence the selection and transport of sRNAs during their movement. Cross-kingdom RNAi impacts host-pathogen interactions, where sRNAs can be transferred between pathogens and host plants to modulate gene expression, suppress immunity, and inhibit pathogen virulence. Understanding the mechanisms and dynamics of cross-kingdom RNAi has important implications for plant protection. It offers potential applications in developing RNA-based pesticides to control diseases and pests in crops. Harnessing the power of RNA-based approaches can provide new strategies for managing plant diseases and improving crop yield.

References

- [1] J. Zeng, V.K. Gupta, Y., Jiang, B. Yang, L. Gong, and H. Zhu, "Cross-kingdom small RNAs among animals, plants and microbes", *Cells*, vol. 8, p. 371, April 2019.
- [2] S.M. Hammond, E. Bernstein, D. Beach, and G.J. Hannon, "An RNA-directed nuclease mediates post-transcriptional gene silencing in *Drosophila* cells", *Nature*, vol. 404, pp. 293-296, March 2000.
- [3] E. Bernstein, A.A. Caudy, S.M. Hammond, and G.J. Hannon, "Role for a bidentate ribonuclease in the initiation step of RNA interference", *Nature*, vol. 409, pp. 363-366, January 2001.
- [4] D.P. Bartel, "MicroRNAs: genomics, biogenesis, mechanism, and function", *Cell*, vol. 116, pp. 281-297, January 2004.
- [5] L. He, and G.J. Hannon, "MicroRNAs: small RNAs with a big role in gene regulation", *Nat. Rev. Genet.*, vol. 5, pp. 522-531, July 2004.
- [6] Q. Cai, B. He, K.H. Kogel, and H. Jin, "Cross-kingdom RNA trafficking and environmental RNAi—nature's blueprint for modern crop protection strategies", *Current opinion in microbiology*, vol. 46, pp.58-64, December 2018.
- [7] M. Wang, and H. Jin, "Spray-induced gene silencing: a powerful innovative strategy for crop protection" *Trends in microbiology*, vol. 25, pp. 4-6, January 2017.
- [8] M. Wang, A. Weiberg, F.M. Lin, B.P. Thomma, H.D. Huang, and H. Jin, "Bidirectional cross-kingdom RNAi and fungal uptake of external RNAs confer plant protection", *Nature plants*, vol. 2, pp.1-10, September 2016.
- [9] A. Weiberg, and H. Jin, "Small RNAs—the secret agents in the plant-pathogen interactions" *Current opinion in plant biology*, vol. 26, pp. 87-94, August 2015.
- [10] A. Weiberg, M. Bellinger, and H. Jin, "Conversations between kingdoms: small RNAs", *Current opinion in biotechnology*, vol. 32, pp. 207-215, April 2015.
- [11] A. Weiberg, M. Wang, F.M. Lin, H. Zhao, Z. Zhang, I. Kaloshian, H.D. Huang, and H. Jin, "Fungal small RNAs suppress plant immunity by hijacking host RNA interference pathways" *Science*, vol. 342, pp. 118-123, October 2013.
- [12] M. Wang, N. Thomas, and H. Jin, "Cross-kingdom RNA trafficking and environmental RNAi for powerful innovative pre-and post-harvest plant protection", *Current opinion in plant biology*, vol. 38, pp. 133-141, August 2017.
- [13] M. Derbyshire, M. Mbengue, M. Barascud, O. Navaud, and S. Raffaele, "Small RNAs from the plant pathogenic fungus *Sclerotinia sclerotiorum* highlight host candidate genes associated with quantitative disease resistance", *Molecular Plant Pathology*, vol. 20, pp. 1279-1297, September 2019.
- [14] T. Zhang, Y.L. Zhao, J.H. Zhao, S. Wang, Y. Jin, Z.Q. Chen, Y.Y. Fang, C.L. Hua, S.W. Ding, and H.S. Guo, "Cotton plants export microRNAs to inhibit virulence gene expression in a fungal pathogen", *Nature plants*, vol. 2, pp. 1-6, September 2016.
- [15] D. Nowara, A. Gay, C. Lacomme, J. Shaw, C. Ridout, D. Douchkov, G. Hensel, J. Kumlehn, and P. Schweizer, "HIGS: host-induced gene silencing in the obligate biotrophic fungal pathogen *Blumeria graminis*", *The Plant Cell*, vol. 22, pp. 3130-3141, September 2010.
- [16] Y. Qiao, L. Liu, Q. Xiong, C. Flores, J. Wong, J. Shi, X. Wang, X. Liu, Q. Xiang, S. Jiang, and F. Zhang, "Oomycete pathogens encode RNA silencing suppressors", *Nature genetics*, vol. 45, pp. 330-333, March 2013.

- [17] Q. Xiong, W. Ye, D. Choi, J. Wong, Y. Qiao, K. Tao, Y. Wang, and W. Ma, “*Phytophthora* suppressor of RNA silencing 2 is a conserved RxLR effector that promotes infection in soybean and *Arabidopsis thaliana*”, *Molecular Plant-Microbe Interactions*, vol. 27, pp. 1379-1389, December 2014.
- [18] L.P. Lim, N.C. Lau, E.G. Weinstein, A. Abdelhakim, S. Yekta, M.W. Rhoades, C.B. Burge, and D.P. Bartel, “The microRNAs of *Caenorhabditis elegans*”, *Genes & development*, vol. 17, pp. 991-1008, April 2003.
- [19] B. Bartel, and D.P. Bartel, “MicroRNAs: at the root of plant development?”, *Plant physiology*, vol. 132, pp. 709-717, June 2003.
- [20] I. Papp, M.F. Mette, W. Aufsatz, L. Daxinger, S.E. Schauer, A. Ray, J. Van Der Winden, M. Matzke, and A.J. Matzke, “Evidence for nuclear processing of plant micro RNA and short interfering RNA precursors”, *Plant physiology*, vol. 132, pp. 1382-1390, July 2003.
- [21] M.D. Horwich, C. Li, C. Matranga, V. Vagin, G. Farley, P. Wang, and P.D. Zamore, “The *Drosophila* RNA methyltransferase, DmHen1, modifies germline piRNAs and single-stranded siRNAs in RISC”, *Current biology*, vol. 17, pp. 1265-1272, July 2007.
- [22] R. Sunkar and J.K. Zhu, “Micro RNAs and short-interfering RNAs in plants”, *J. Integr. Plant Biol.*, vol. 49, pp. 817-826, June 2007.
- [23] G. Tang, “siRNA and miRNA: an insight into RISCs”, *Trends in biochemical sciences*, vol. 30, pp. 106-114, February 2005.
- [24] B. Yu, Z. Yang, J. Li, S. Minakhina, M. Yang, R.W. Padgett, R. Steward, and X. Chen, “Methylation as a crucial step in plant microRNA biogenesis”, *Science*, vol. 307, pp. 932-935, February 2005.
- [25] J. Li, Z. Yang, B. Yu, J. Liu, and X. Chen, “Methylation protects miRNAs and siRNAs from a 3'-end uridylation activity in *Arabidopsis*”, *Current biology*, vol. 15, pp. 1501-1507, August 2005.
- [26] Y. Mao, X. Xue, and X. Chen, “Are small RNAs a big help to plants?”, *Science in China Series C: Life Sciences*, vol. 52, pp. 212-223, March 2009.
- [27] M.A. Carmell, Z. Xuan, M.Q. Zhang, and G.J. Hannon, “The Argonaute family: tentacles that reach into RNAi, developmental control, stem cell maintenance, and tumorigenesis” *Genes & development*, vol. 16, pp. 2733-2742, November 2002.
- [28] R.W. Carthew, and E.J. Sontheimer, “Origins and mechanisms of miRNAs and siRNAs”, *Cell*, vol. 136, pp. 642-655, February 2009.
- [29] X. Chen, “Small RNAs-secrets and surprises of the genome”, *The plant journal*, vol. 61, pp. 941-958, March 2010.
- [30] F. Vazquez, H. Vaucheret, R. Rajagopalan, C. Lepers, V. Gascioli, A.C. Mallory, J.L. Hilbert, D.P. Bartel, and P. Crété, “Endogenous trans-acting siRNAs regulate the accumulation of *Arabidopsis* mRNAs”, *Molecular cell*, vol. 16, pp. 69-79, October 2004.
- [31] A. Peragine, M. Yoshikawa, G. Wu, H.L. Albrecht, and R.S. Poethig, “SGS3 and SGS2/SDE1/RDR6 are required for juvenile development and the production of trans-acting siRNAs in *Arabidopsis*”, *Genes & development*, vol. 18, pp. 2368-2379, October 2004.
- [32] P. Guleria, M. Mahajan, J. Bhardwaj, and S.K. Yadav, “Plant small RNAs: biogenesis, mode of action and their roles in abiotic stresses”, *Genomics, proteomics & bioinformatics*, vol. 9, pp. 183-199, December 2011.
- [33] C.A. Brosnan, and O. Voinnet, “Cell-to-cell and long-distance siRNA movement in plants: mechanisms and biological implications”, *Current opinion in plant biology*, vol. 14, pp. 580-587, October 2011.
- [34] J.C. Palauqui, T. Elmayer, J.M. Pollien, and H. Vaucheret, “Systemic acquired silencing: transgene-specific post-transcriptional silencing is transmitted by grafting from silenced stocks to non-silenced scions”, *The EMBO journal*, vol. 16, pp. 4738-4745, August 1997.
- [35] P. Sarkies, and E.A. Miska, Small RNAs break out: the molecular cell biology of mobile small RNAs”, *Nature Reviews Molecular Cell Biology*, vol. 15, pp. 525-535, August 2014.
- [36] E.N. Nolte-’t Hoen, H.P. Buermans, M. Waasdorp, W. Stoorvogel, M.H. Wauben, and P.A. ’t Hoen “Deep sequencing of RNA from immune cell-derived vesicles uncovers the selective incorporation of small non-coding RNA biotypes with potential regulatory functions”, *Nucleic acids research*, vol. 40, pp. 9272-9285, October 2012.
- [37] M. Colombo, G. Raposo, and C. Théry, “Biogenesis, secretion, and intercellular interactions of exosomes and other extracellular vesicles” *Annual review of cell and developmental biology*, vol. 30, pp. 255-289, October 2014.
- [38] J. Guduric-Fuchs, A. O’Connor, B. Camp, C.L. O’Neill, R.J. Medina, and D.A. Simpson, “Selective extracellular vesicle-mediated export of an overlapping set of microRNAs from multiple cell types”, *BMC genomics*, vol. 13, pp. 1-14, December 2012.
- [39] D.S. Skopelitis, K. Hill, S. Klesen, C.F. Marco, P. von Born, D.H. Chitwood, and M.C. Timmermans, “Gating of miRNA movement at defined cell-cell interfaces governs their impact as positional signals”, *Nature communications*, vol. 9, p. 3107, August 2018.
- [40] B. Wang, Y. Sun, N. Song, M. Zhao, R. Liu, H. Feng, X. Wang, and Z. Kang, “*Puccinia striiformis* f. sp. *tritici* microRNA-like RNA 1 (Pst-miR1), an important pathogenicity factor of *Pst*, impairs wheat resistance to *Pst* by suppressing the wheat pathogenesis-related 2 gene”, *New Phytologist*, vol. 215, pp. 338-350, July 2017.
- [41] M. Brill, E. Asquini, M. Moser, P.L. Bianchedi, M. Perazzolli, and A. Si-Ammour, “A multi-omics study of the grapevine-downy mildew (*Plasmopara viticola*) pathosystem unveils a complex protein coding-and noncoding-based arms race during infection”, *Scientific Reports*, vol. 8, pp. 1-12, January 2018.
- [42] C. Gualtieri, P. Leonetti, and A. Macovei, “Plant miRNA cross-kingdom transfer targeting parasitic and mutualistic organisms as a tool to advance modern agriculture”, *Frontiers in Plant Science*, vol. 11, June 2020.
- [43] W. Islam, A. Noman, M. Qasim, and L. Wang, “Plant responses to pathogen attack: small RNAs in focus” *International journal of molecular sciences*, vol. 19, p. 515, February 2018.
- [44] W. Islam, M. Qasim, A. Noman, M. Adnan, M. Tayyab, T.H. Farooq, H. Wei, and L. Wang, “Plant microRNAs: Front line players against invading pathogens”, *Microbial pathogenesis*, vol. 118, pp. 9-17, May 2018.
- [45] J. Huang, M. Yang, L. Lu, and X. Zhang, “Diverse functions of small RNAs in different plant-pathogen communications”, *Frontiers in microbiology*, vol. 7, p. 1552, October 2016.
- [46] C. Hua, J.H. Zhao, and H.S. Guo, “Trans-kingdom RNA silencing in plant-fungal pathogen interactions”, *Molecular plant*, vol. 11, pp. 235-244, February 2018.
- [47] G. Huang, R. Allen, E.L. Davis, T.J. Baum, and R.S. Hussey, “Engineering broad root-knot resistance in transgenic plants by RNAi silencing of a conserved and essential root-knot nematode parasitism gene”, *Proceedings of the National Academy of Sciences*, vol. 103, pp.14302-14306, September 2006.
- [48] Q. Cai, L. Qiao, M. Wang, B. He, F.M. Lin, J. Palmquist, S.D. Huang, and H. Jin, “Plants send small RNAs in extracellular vesicles to fungal pathogen to silence virulence genes”, *Science*, vol. 360, pp. 1126-1129, June 2018.

- [49] H. Zhang, J. Guo, R.T. Voegelé, J. Zhang, Y. Duan, H. Luo, and Z. Kang, "Functional characterization of calcineurin homologs PsCNA1/PsCNB1 in *Puccinia striiformis* f. sp. *tritici* using a host-induced RNAi system", *PLoS One*, vol. 7, pe. 49262, November 2012.
- [50] X. Zhu, T. Qi, Q. Yang, F. He, C. Tan, W. Ma, R.T. Voegelé, Z. Kang, and J. Guo, "Host-induced gene silencing of the MAPKK gene PsFUZ7 confers stable resistance to wheat stripe rust", *Plant physiology*, vol. 175, pp. 1853-1863, December 2017.
- [51] L. Miozzi, G. Gambino, J. Burgyan, and V. Pantaleo, "Genome-wide identification of viral and host transcripts targeted by viral siRNAs in *Vitis vinifera*", *Molecular plant pathology*, vol. 14, pp. 30-43, January 2013.
- [52] Y. Song, and S.P. Thomma, "Host-induced gene silencing compromises *Verticillium* wilt in tomato and *Arabidopsis*", *Molecular plant pathology*, vol. 19, pp. 77-89, January 2018.
- [53] A. Niehl, M. Soininen, M.M. Poranen, and M. Heinlein, "Synthetic biology approach for plant protection using dsRNA", *Plant biotechnology journal*, vol. 16, pp.1679-1687, September 2018.
- [54] A. Koch, D. Biedenkopf, A. Furch, L. Weber, O. Rossbach, E. Abdellatef, L. Linicus, J. Johannsmeier, L. Jelonek, A. Goesmann, and V. Cardoza, "An RNAi-based control of *Fusarium graminearum* infections through spraying of long dsRNAs involves a plant passage and is controlled by the fungal silencing machinery", *PLoS pathogens*, vol. 12, pe. 1005901, October 2016.
- [55] A.G. McLoughlin, N. Wytinck, P.L. Walker, I.J. Girard, K.Y. Rashid, T. de Kievit, W.G. Fernando, S. Whyard, and M.F. Belmonte, "Identification and application of exogenous dsRNA confers plant protection against *Sclerotinia sclerotiorum* and *Botrytis cinerea*", *Scientific Reports*, vol. 8, pp.1-14, May 2018.
- [56] L. Nerva, M. Sandrini, G. Gambino, and W. Chitarra, "Double-stranded RNAs (dsRNAs) as a sustainable tool against gray mold (*Botrytis cinerea*) in grapevine: Effectiveness of different application methods in an open-air environment", *Biomolecules*, vol. 10, p. 200, January 2020.
- [57] C.N. Taning, S. Arpaia, O. Christiaens, A. Dietz-Pfeilstetter, H. Jones, B. Mezzetti, S. Sabbadini, H.G. Sorteberg, J. Sweet, V. Ventura, and G. Smagghe, "RNA-based biocontrol compounds: current status and perspectives to reach the market", *Pest management science*, vol. 76, pp. 841-845, March 2020.
- [58] P. Bachman, J. Fischer, Z. Song, E. Urbanczyk-Wochniak, and G. Watson, "Environmental fate and dissipation of applied dsRNA in soil, aquatic systems, and plants", *Frontiers in plant science*, vol. 11, p. 21, February 2020.
- [59] X.S. Song, K.X. Gu, X.X. Duan, X.M. Xiao, Y.P. Hou, Y.B. Duan, J.X. Wang, N. Yu, and M.G. Zhou, "Secondary amplification of siRNA machinery limits the application of spray-induced gene silencing", *Molecular plant pathology*, vol. 19, pp. 2543-2560, December 2018.
- [60] L. Qiao, C. Lan, L. Capriotti, A. Ah-Fong, J. Nino Sanchez, R. Hamby, J. Heller, H. Zhao, N.L. Glass, H.S. Judelson, and B. Mezzetti, "Spray-induced gene silencing for disease control is dependent on the efficiency of pathogen RNA uptake", *Plant biotechnology journal*, vol. 19, pp. 1756-1768, September 2021.