

An Analytical Review on Cognitive training for Behavioral Improvement, Brain Rewiring and Memorization during Adolescence Period Using Neural Plasticity

Prateek Pratyasha¹, Saurabh Gupta¹, Aditya Prasad Padhy²

¹Department of Biomedical Engineering, NIT, Raipur, Chhattisgarh

²Department of Electrical and Electronics Engineering, ARKA Jain University, Jamshedpur, Jharkhand

ABSTRACT: Projection of neural interconnection goes after some critical synaptic alteration during adolescence period. The adolescent brain is yet to be relied on pre-frontal cortex. All emotions, impulsive behaviors affect the brain maturity which ultimately resists self-rewiring of brain. Memories are associated with brain-rewiring; hence needs to be retrieved efficiently for memory preservation. As there is no reliability of diagnosis or medication to improve the potentiality of brain; Neural Plasticity can be implicated to reorganize the neural pattern and boost up brain rewiring. This article is intended towards analyzing the anatomical and psychological attributes of brain during adolescence period and implicating neural plasticity as a cognitive training process to facilitate synaptic activities in neurons. For a holistic neuro-anatomical improvement, neural plasticity is served as the driving force to rewire brain functionality. It brings an immense change in neural pattern, there by triggers the synaptic activities. Although an inflicted fluctuate in plasticity level may impair the growth of rewiring, adolescent brain can regain its plasticity no sooner. Instead of any pharmacy dependent technique, we should follow brain stimulation based cognitive trainings for the overall cognition and memorization.

KEYWORDS Brain rewiring, Neural Plasticity, Adolescence Period, Memorization, Synaptic activities, Brain maturation, Prefrontal Cortex.

1. INTRODUCTION

Adolescence period, the most sensitive phase of life is often ambiguous for its dramatic neural and psychological changes. It encompasses with vigorous puberty hormones that effecting emotional, behavioral and experiential independency between the ages of 12 years to 19 years which influence neural transition. The complex neural transition initiates from the genetic blueprint and synaptic synchronization of neurons where memories reside[1]. Although, development of human neural schema and its functionality begins from pre-natal phase; the neural system tends to stabilize slowly with biological development of infants and it varies from one individual to another[2]. The reorganization of neuron alters utmost psychologically as we tend to reach adolescence period; but doesn't cease the process of stabilization.

During adolescence, brain deals with some sensitive phases such as self-consciousness, social isolation, interpersonal violence, peer pressure and other puberty issues [3]. This may be the prior reason behind such dramatic variation affecting physiological, mental and psychological state of mind. Undoubtedly, the neuro-anatomical alteration occurs due to changes in hormones and social surroundings of that individual. The experience from the internal and external surroundings plays an

important role in brain development (see Table 1.1). The ability of thinking, learning, cognition and memorizing occurs in the anterior and posterior lobes of brain and the mechanism is unstoppable. In the meanwhile, some neural windows are specifically adaptive for new cognitions and experiences. Environmental stimuli also exert profound consequences on neural connectivity. Later no, all the taken experiences are stored as explicit memories and short-term memories in the hippocampus and prefrontal cortex respectively [4]. Reorganization of those memories critically demands for Neural Plasticity (NP) as a subtractive interconnection of neurons.

The outcome of neuro-anatomical alteration in cortical brain influences the frequent behavioral changes in adolescents [5] gives rise to plasticity or neurologically termed as "Neural Plasticity" (NP). Although the concept was discovered 100 years ago by William James, in his book entitled "Principle of psychology" where he recognized that human brain is capable of reorganizing memory and cognition. But, concept of NP was highlighted to discuss

Adolescence Regulatory Parameters	Dominated Hormones		Affected Brain regions	Application	Ref.
Hormones	Testosterone			New cell growth, Increase in grey matter volume	
	Progesterone				
	Estrogen				
	ADH		Hypothalamus		
	TSH				
	Sex steroids			Development of secondary sexual characteristics	
Social Behavior	Dominant factors	Non-dominant factors			
	Education	Life satisfaction			
	Sexual orientation	Gender differentiation			
	Health literacy	Cognitive function starting at late life			
	Stress	Abuse of other substances			
	Negative Mood (Depression, Anxiety)	Sexual Practice			
	Diet Patterns	Risk Taking behaviors			
	Physical activates	Social support			
	Financial Strain	Social and cultural norms			
	Exposure to drugs and alcohol	-			
	Social connection and Isolation	-			
	Exposure to violence	-			

about brain mapping by Hubel and Wiesel in 1967 where they emphasize inherent stimuli and period of experience for the cognitive development of cortical brain. NP is active in specified brain regions such as inferior parietal cortex and prefrontal cortex exposing highest synapses which is responsible for neuro-anatomical reorganization

of synaptic network [6] related to independent aspects. However during brain injuries, the cognitive re-organization of prefrontal cortex is different from other fore-brain regions affecting cognitive flexibility, prospective memory, impulsive inhibition and behavior.

Every teenager experiences physical and psychological changes activated by hormonal systems to cope of with all

kinds of physical, mental and social situations. Physical changes are mediated by synaptic nervous system whereas psychological changes are slower and more responsive towards protracted parameters.

Figure 1(a) and (b) shows a comparative graph of both physical and psychological parameters between adolescents and adults accordingly.




Fig. 1(a)




Fig. 1(b)

Reproductive behaviors of adolescents are initiated at the age of 13 years where genital hormones are responsible for organizational and behavioral changes. However, sex-oriented functional changes can only be observed during adulthood concluding the fact that organizational changes are constant whereas functional changes are inconsistent. Growth hormones are responsible for neurogenesis and synaptogenesis during neural development. Additionally, it is involved in other brain functions such as learning and memory. It has drawn the attention when researches are made on preservation of neural integrity after any brain injury. All these physical and psychological parameters are responsible for regulating neurogenesis, promoting neural proliferation and activate neuro-protection. Among the physical hormones, the puberty is majorly affected by puberty hormones such as testosterone in male, estrogen and progesterone in female.

Apart from the earlier generalized discoveries in neurogenesis, NP underscores predominately in cognitive learning accomplishing the competencies. As learning

process tends to ignite by age, plasticity has implemented through various methods such as auditory-visual training programme, When behavioral consequences and practical challenges of life are outlined with the neural architecture, neuroimaging techniques have associated with NP to describe all shapes of emerging brain [7]. However, many studies have shed light on the role of synaptic activities gleaned as an executive function to refine the prefrontal cortex connectivity, simultaneously highlighting vulnerabilities during adolescent periods [8]. As the brain starts to mature during this phase, neural connectivity depends on sleep spindles for the enhancement of memory [9]. But, this process is a way long to initiate brain maturation in adolescents when it calls for a simple stimulus-response learning task. Reinforcement learning often tends to speed up in adolescents when working memory subtle due to puberty hormones [10]. Observing many such issues, NP should be preferred as a presumptive option for indisputable initiation of brain rewiring as well as memory enhancement.

This review article aims to analyze the schematic framework of NP dependent on internal and external sources implicated for cognitive learning, brain rewiring, memory synthesize, decision making process etc. Many literatures of recent years are referenced manifesting the development of NP to amend synaptic activities through neurons. To extend this, we have discussed briefly about brain rewiring and existence of memory in adolescents. Then, we have reviewed various aspects of NP as a fillip up and evaluate the gap between excessive and deficiency of NP during the intervention. Finally, regaining of plasticity has recuperated by numerous techniques such as neural replacement, synaptic modeling, and metaplasticity along with many pharmacological approaches.

2. THE STATE-OF-ART OF BRAIN REWIRING

The dynamic reorganization of brain is perpetually adaptive for skillful learning and acquiring experience from the circumstances. Brain accommodates to impulsive synaptic changes while activating the neurons from dendrites extensions and alters its mode of functionality. Meanwhile, the alteration in brain functionality provides fruitful scope to rewire the brain. Earlier, it was a myth that brain rewires itself according to the neuro-anatomical development and eventually becomes static for adults. Although, the brain reaches its 3/4th of maturity by the age of 2, the prefrontal cortex (PC) still has to wait up to 25 years for its complete maturation [11]. The immaturity of PC reflects physically, hormonally and intellectually during adolescence period. But the reorganization of brain never stops and rewires itself whenever the neurons tend to reconnect among themselves.

Modern cognitive science observed that we have to put some extra effort to rewire the brain to its full efficiency. Initiation of brain rewiring appears in sub-cortical regions, but it is accustomed with PC involving an effortful learning-experience pathway. However, it is influenced by brain mapping as an experiential outcome. This process of lasts till we die, and not only restricted to the early stages of life. From a report given by Carnegie Mellon's Center for Cognitive Brain Imaging (CCBI), learning new skills influences the brain to rewire itself in younger generation[12]. The age between 5 to 12 years is perfect period for physical and mental training as children in this age group are adaptive to express their interest, capable of receive any new skill and active in memorization. Millions of neurons producing synapses are activated which promotes the structural and functional brain development. However, most of the unused synapses are died off due to synaptic pruning [8].

By the time teenage has arrived, the pressure of self recognition, understanding the intentions and emotions of others unfold neural interaction strategically. The role of neural interaction during adolescence period has put an unconventional impact to rewire the "Social brain". Understanding the social dogma quicken brain rewiring and cultivates its functionality minutely[13]. In addition to that, infestation of immense synapses strengthens the brain which is completely dominated by synaptic pruning. The end process of pruning predominate myelin over grey matter in PC lobe to reconstruct the brain extensively. Existence of numerous emotions such as thoughts, beliefs, experiences, skills and decision-making abilities may be last for lifetime or may be lost forever in teenagers. Synaptic pruning selects the neural interactions for those emotions to rewire the brain efficiently; simultaneously eradicating unused neural connections [14].

A rewired brain unfolds the sections where planning, hypothesis, insights and mood regulation resides. The freedom to explore such sections is exquisitely demands for social flexibility. Although brain rewiring is a legacy of neurogenesis, an akin rivet slants towards different sensory and motor functions, Gonad hormones, age, sex, nutrition, stimulants and social key factors[15]. Despite of recognizing the emotions like cry, laugh, anger, fear and so forth that developing the brain roots biologically; the adolescent brain may also discover the existence of some complex emotions to deal with such as embracement, guilt, aggression, confusion etc. All of these key issues can affect the way of brain rewiring[16, 17]. Placing all cells together and engages the brain in its right way is not an easy job to do. The myth about genes regulating our emotions, behavior and experience has shuttered. Recently, Neuroscience has witnessed that prevision and potentiality of learning based experience change the neural

interaction. To acknowledge this legitimacy, our mind should be stress free, tamed and calm before learning something new. Therefore, teenagers should lay out their brain for an upright training to exaggerate brain rewiring.

3. EXISTENCE OF MEMORY

The unstoppable process of acquiring information, storing it and later retrieving them when needed over time is what we called as "Memory"[18]. It resides in the hippocampus, amygdale, cerebellum and prefrontal cortex cerebrated with neuro-anatomy. This biological process is not eternal; it may be lost to recall its evolution either in the pre-natal phase of life or when the living body is brain dead. In aid to that, subsistence of memories can highly affected by several brain diseases, stress, lack up sleep, indulgent life style and the list goes on. However, formation of memory starts from an unpredictable node and hit the infancy queerly[19].

Due to this phenomenon of infantile amnesia[20, 21], we have never triggered back to our flashback. After the age of 3, the child starts to learn and remember language, people, quotidian automatically and those memories are stores as implicit memories. Contradictorily, children are well acquainted with their mother's voice since their birth which proves the existence of declarative memory developed in toddlers. As the child jump into a tough school life; episodic and semantic memories have already built up[22]. By the time; daily routines such as wearing shoes, organizing books in the bag, riding bicycle etc. are also started to pile up as procedural memories in cerebellum. This memory is continues to germinate as per the child's neuro-anatomical development. At the door step to adolescence period; auto-biographical memory, long-term and short-term memory are bloomed at the hippocampus. To expand the cognitive process, memory serves as an alpha factor which exercises the learning, events, skill and other symbiotic entities from past as well as present. It treats the past events as references, working on present while storing experiences and reusing those experiences in future[23].

Memory and disremembering co-exist in mind since childhood, but there are vigorous ways to overcome the bad habit of forgetting. Visual memory is always in prior to recall the information in an agile mind of a child. However, it fades away as the child tends to learn from the real world [24]. To overcome this problem, different Mnemonics are implicated to increase the recalling ability by storing each memory segments as a cyclorama. Memory palaces have proved to be the best ever discovered mnemonics devices of all the time to keep the memories as files and import them when needed. Since ancient times, adequate nourishment impacts cognition, hence parents put their children on a Mediterranean diet with ample of omega-3, fatty acid, Docosahexaenoic acid

(DHA), vitamin B6 complex and a few amount of epicatechin usually found in tea, coffee and berries. However, bellyful nutrition is not enough during childhood to memorize uncountable numbers of iterative and non-iterative information[25]. Therefore, chunking helps in clubbing our memories in a bunch and leaves it to retrieve later. Cognitive training has turned up to change the most usual way of memory development in triggering working memory for the repetition of multiple tasks as well as by incrementing the information quantity.

All these methodologies mature the memorizing abilities throughout entire childhood, but brainstorming and cognitive performances tends to drop off in the crucial period of adolescence. Besides that adolescent brain is intensified in memorizing plenty number of stuffs, some peculiar teenager attributes can sluggish its performance. As mentioned in [26], disturbances in memory such as unable to recall recent information, often mislay things around, forgetting names, events whenever required not only affect their routine life and mental stability but also lead to depression. Many neurologists and psychologists have failed to go underneath the root cause of memory deficiency. However, slow learning activities, lack of creativity, lack of concentration, excessive drug addiction, peer pressure, stress, anxiety, outdistance from society are some of the possible reasons behind memory loss in teens. Adding to that, some medical history of brain injuries, neuro-degeneration, seizures, and endocrine dysfunctions can cause either sub-accurate or chronic memory loss[27].

Life with such memory deficiency becomes chaotic when brain gets over-stimulated by puberty hormones. Memory is affected the most when the mind deals with thousands of emotions, planning, strategies, ambitions and social stigma. Moreover, sleeping pattern overpowers memory in such a way that a deprived adolescent experiences lack of concentration compared to a rested teenager. To check the memory density and concentration, a preliminary skill test has conducted based on tricky mental math questions, puzzle solving, basic information regarding dates and events. Such delicate issues need a medical practitioner to keep an eye on their sleeping pattern as well as diagnose possibilities of any pre-term or post-term seizures, brain injuries, anxiety issues etc[28]. However, medical science doesn't provide any facilities to diagnose or cure memory deficiency. Hence, we need to train our brain in a way that would reactivate all the dozy neurons automatically.

4. NEURAL PLASTICITY AS AN ADDED FILLIP

Originally, Neuropsychology recounted that development of brain depends on plasticity. During this development, brain is subjected to reorganize neural patterns and synchronize its functionality depending on the

external and internal stimuli. The occurrence of NP begins at pre-natal stage of life where the immature brain prepares itself for functioning which is the most sensitive period for vigorous studies. The memories developed during this stage may be responsible for maternal recognition. During the occurrence of synaptogenesis, most of the events are carried forward if something new is learned and memorized. In case of brain injury, it adapts to repair the damaged neurons and compensate learning process from remaining lobes of brain[29].

The most frequently used technique to measure NP level is functional MRI (fMRI) which figures out the changes of blood flow in cerebral cortex and identifies the functional brain regions to when subjected to a given task in both active and resting state[29]. University of Miami experimented about the induced NP by intermittent theta burst stimulation (ITBS), a non-invasive technique to study the impact of plasticity on aging. Amplitude of highly emerging brain waves is validated initially through transcranial magnetic stimulation (TMS). However, the potentiality of the synapses can be modulated by using long-term potentiation (LTP) and long-term depression (LTD). But all these techniques are prone to clinical trials and only meant for the treatment of neurological and psychological defects in children and hence, such argumentative proposals are failed to alter the synaptic sites in adolescents. The time and frequency domain analysis of NP was carried out by STDP, but was also conk out due to onetime generation of single brain signal stipulated with *in vitro* only. Later, all the neural alteration subjected to paired associative stimulation (PAS) was meticulously shifted towards the study of NP. Some research have used paired associative stimulation (PAS) technique to measure the amount of motor cortex to increase the amplitude of motor potentials in teens and failed due to the raid cortical changes occur in children. Displaying the morphological structure of brain alterations in a three dimensional image are exhibited by enlarging the synapses or by increasing the terminal density. But, such neuroimaging techniques through positron emission tomography (PET), magnetic resonance or cranial ultrasound are spurious to depict the morphologic attributes of brain. Hence, a scrupulous attention is needed when the child reaches to the most sensitive period of life.

NP is segregated in two forms where Structural plasticity alters the neuro-anatomical structure and synaptic plasticity which alters the neuro-chemical composition of body. During the beginning period of neurogenesis, NP was circumscribed as the alteration in brain connectivity depending on external and internal stimuli. No sooner, the definition was modified when a cognitive neural transition occurs due to some critical synaptic alteration. Later on, the theory was ramified when

our aptitude, behavior and strategies to do any task and decision making ability have build up at growing stage of life. A special attention should be given to all the adolescents as it is the shadow of one's personality that reflects from behavior. The more a teen brain learns, it becomes difficult to self-rewire the brain due to synaptic pruning. As pruning can't be avoided at the striking age of puberty, the process of rewiring often faces some difficulty to maintain itself in late 20's. Hence, NP is one the more trial & tested technique to get rid of pruning and stabilizes the synaptic alterations.

By the age of 5, a child's brain is open to all the positive and negative thoughts. Hence, the foundation of NP needs to be authorized at such early stage. NP helps in strengthening brain connectivity and expanding the dendrite forks and there by detecting relationships between diverse sources of information, learning something new or more specifically being curious to learn, reviewing every minute events in wider picture, and understanding the global implications of specific issues, capable of taking right decisions at right time and so on. In special cases of brain injury or disorder, the learning skill helps in recovering the lost neural interconnections more quickly. NP encourages our capacity of learning, cognition and memory in response with external and internal surroundings. It is associated with the structural and functional changes in prefrontal cortex [30].

Alteration of brain connectivity is highly dependent on the capability of learning, memory and adaption. A proficient intervention of plasticity synthesizes an adolescent brain during skillful learning and multi-tasking. As cognition is the pure reflection of learning something new, a child's brain should be trained accordingly to be more flexible regarding accessing and retrieving information. During the intervention, the synapses fluctuates more as we respond to outer stimuli. Specific neural tracts associated with synaptic networks for cognitive intervention. The interventions promise for deep brain stimulation, cognitive training and neuro-pharmacology processing. Dynamic modification of neural networks leads to memory formation and brain rewiring under the framework of synaptogenesis, neurogenesis, synaptic synchronization and fluctuation. As synaptic strengthening is one of the consolidated features of memory, memory retrieve are stringently dependent on type of synaptic alterations.

A.ADEQUACY OF PLASTICITY FOR ADOLESCENTS

How much plasticity is sufficient to rewire the adolescent brain is the prior concern for researchers. When plasticity takes place, number of neurons and synapses raise accordingly. As we proceed towards adulthood, the

synaptic synchronization between neurons has reduced to half of its number, which is technically termed as synaptic pruning. If a cognitive training is alimented to a group of children, then a better results would expected from adolescents as compared to adolescents and adults. While the quality of plasticity is increased in adolescents, major hormonal and neuro-anatomical transition occurs which impacts the cognitive process profoundly[31]. When adolescents practice their cognitive skills, the working memory or short-term memory capacity enhances and simultaneously all the synaptic activities synchronize with the increased neural tissues[32].

Since a few years, psycho-stimulant medications such as Amphetamine, Rivastigmine, Citalopram are clinically tested to induce the potentiality of NP by altering the neural interconnection and strengthening the synaptic sites[33]. But, no pills are guaranteed for a sharp brain to store unlimited information. Besides that, a mild overdose of pills can impair the biological stimuli leading to a maladaptive orientation. Recently, non-pharmacological agents such as music, quality of sleep, meditation, brain mapping, brain stimulation etc are introduced to enhance cognitive connections. Despite of being universally accepted, it is not taken too seriously to implement in real life. They seem to be more efficacious compared to currently available drugs. Adjuvant therapies using flavonids are encountered as a standard treatment to enhance neuro stimulations and repair the impaired brain functions. The common hallmark of nerve tissue damage is increased by oxidative stress and inflammation. Thus, the studies on flavonoids with strong antioxidant and anti-inflammatory properties as a potential application in neuro intervention have been carried out for a long time. However, recent results have revealed another important property of these compounds in CNS therapy. Flavonoids are inhibited with antioxidants and anti-inflammatory properties to actuate neuro protective activity, and promote synaptogenesis and neurogenesis[34].

On the other hand, neural deterioration is a common symptom occurs according to the grown up age. Hence, the level of plasticity reduces gradually with teenager's naïve towards life. Traumatic brain injuries can severely interrupt the cognitive and emotional behavior followed by uncontrolled sensory-perception, impaired neuro-modulatory function. Such distorted plasticity or "negative plasticity" often restricts the teenagers participating in academic activities, socio-cultural functions and cognition grooming programs[35]. In aid to that, aging factor is the most unavoidable parameter which naturally declines NP despite of vigorous NP enhancer implications. Therefore, possibilities of gaining memory and rewiring the unstoppable brain can be balanced by enhancing the cognitive training. Many challenge based

tasks are essential elements to trigger plasticity which merely expecting some relevancy and importance in return. Although, several studies have given hypothesis on enlarged plasticity either enhanced or impaired the cognitive learning for same intervention; but the crystallized cognition remains invariant irrespective to age or even may mutate during adulthood. As it takes time to bring a stringent structural and functional changes in plasticity, inventive brush up of task schedule must be refocused.

B. REGAINING PLASTICITY

An adolescent brain has the ability to regain plasticity by repairing the suppressed neurons and drive the synaptic arc efficiently. Brain extends the heterogeneity of neuronal subtypes to drive stem cells and specifies only a few essential subtypes to regain the dysfunctional neural functionality. However, the new neurons are integrated with the pre-exist ones near the hippocampus and too difficult for the hierarchical reorganization. An alternative approach to enhance plasticity is to replace the non-functional neurons with the activated ones under the process of neurogenesis. But this neuron replacement approach must be implemented in newborn for the effective synaptic synchronization. Indeed, the cell replacement technique may be unsafe because of its unknown potential. Hence, both the strategies demands for a successful clinical trial.

But, modern contextual learning always demands for a quick and efficient approach. Hence, an approachable pharmaceutical glutamate receptor, namely “AMPA” is meant to be delivered to the center of learning and memory that’s the hippocampus region. For that, a cellular model of synaptic chaotic activities has been depicted by LTP associated with associative learning. Although neural stimulations are transmitted in synaptic activities, the AMPA receptor acts as an interceder for faster synaptic transmission which is used widely in synaptic potentiation and cognitive learning during experience-dependent NP [36]. In another aspect, remodeling of NP demands for stimulus transduction to maintain the learning and memory related neural function. Hence, an epigenetic enzyme lysine-specific demethylase-1 (LSD1) or neuro LSD1 have been implicated in the neuro specific splicing variant process through an antagonistic mechanism. This transcriptional receptor co-activates the androgen and estrogen receptors which modulates emotional behavior and enhances neural maturation during memory formation. However, the single neuro LSD1 genomic structure has both positive and negative modulation depending on the transcription activities[37].

The balance between dominated and suppressed plasticity exists by entailing the threshold of NP in

metaplasticity where synapses improve their ability to regain NP within a stipulated period. Metaplasticity encountered the change which outlast for a longer bout compared to common plasticity[38]. It can line up the learning intervention strategically despite of any reward uncertainties or explicit optimization of sources. A metaplasticity model is more precise to predict uncertainty than any optimal models considering the neural activities occur in prefrontal cortex[39]. Moreover, meta-plastic models bring synaptic changes in the brain, but doesn’t encroach the synaptic efficacy. Hence, metaplasticity acts as an extenuation between adaptability and precision via simulation whose results can substantially enhance synaptic efficiency. These models strengthen the adaptability-precision palliation by two meta-state frameworks (1) Reservoirs and (2) Buffers[40]. It has observed that synaptic efficiency is robust for reservoirs creating a strong synaptic activity without any evident plasticity[41]. In belie, synaptic efficiency is restricted under the circumstances of buffer meta-state without any major change in plasticity. Hence, meta-plastic models are quite adaptive for plasticity transition and capable of understating any uncertainties.

5. CONCLUSION

Pubertal hormones, impulsive behavior, external and internal environmental stimuli and other factors may restrict synaptic potential for self-rewiring which alternatively retarding memorizing ability. Although a tiny frequent change in neural connectivity provides a scope for self-rewiring, however it has proved to be a myth during the grown-up age. Of course, the inter-relation between brain rewiring and memory enhancement becomes chaotic as the adolescent brain deals with thousands of mixed emotions, stress, hormonal alterations and social hindrances. And, as we tend to adulthood, synaptic pruning starts decaying our long term memory and makes the brain a static organ to respond to obvious stimuli only. Although, basic training sessions has been conducted since decade to enlarge memory density such as meditation, audio-visualization therapy, healthy nutrients and exceptionally psychopathic drugs; still are uncertain in their full-efficiency. However, development of memory is a never-ending process and it explores the synaptic activities with exuberance of external and internal stimuli. This article suggests a cognitive training session for adolescents in which neural plasticity has implicated as an additive fill-up to boost the synaptic activities. However, measurement of NP is seemed to be necessary as we are dealing with a sensitive phase of life. Therefore, adequacy of brain stimulation, brain mapping and Adjuvant therapies can be imposed at the beginning to conciliate the brain from past neuro-anatomical injuries. Sometimes, care should be taken to avoid the negative plasticity as it may cause neural deterioration during brain maturity. Moreover, plasticity

can be refined and crystallized with an amendable way by regaining the threshold of plasticity. For this, metaplasticity is to be suggested which has the ability to bridge the adaptability-precision conflict and cope of with plasticity transition efficiently. But, this is not the termination of neurogenesis. Likewise the unstoppable memory and brain rewiring; the cognitive training process should be modified to deal with synaptic transitions uninterruptedly.

Conflict of Interest: None

REFERENCES

- [1] S.-J. Blakemore and K. L. Mills, "Is adolescence a sensitive period for sociocultural processing?," *Annual review of psychology*, vol. 65, pp. 187-207, 2014.
- [2] J. Stiles, "Neural plasticity and cognitive development," *Developmental neuropsychology*, vol. 18, pp. 237-272, 2000.
- [3] E. C. Dunn, Y. Wang, J. Tse, K. A. McLaughlin, G. Fitzmaurice, S. E. Gilman, *et al.*, "Sensitive periods for the effect of childhood interpersonal violence on psychiatric disorder onset among adolescents," *The British Journal of Psychiatry*, vol. 211, pp. 365-372, 2017.
- [4] J. M. Cooper, F. Vargha-Khadem, D. G. Gadian, and E. A. Maguire, "The effect of hippocampal damage in children on recalling the past and imagining new experiences," *Neuropsychologia*, vol. 49, pp. 1843-1850, 2011.
- [5] F. Crews, J. He, and C. Hodge, "Adolescent cortical development: a critical period of vulnerability for addiction," *Pharmacology Biochemistry and Behavior*, vol. 86, pp. 189-199, 2007.
- [6] S. Chokron, C. Perez, and C. Peyrin, "Behavioral consequences and cortical reorganization in homonymous hemianopia," *Frontiers in systems neuroscience*, vol. 10, p. 57, 2016.
- [7] A. Galván, "Neural plasticity of development and learning," *Human brain mapping*, vol. 31, pp. 879-890, 2010.
- [8] L. D. Selemon, "A role for synaptic plasticity in the adolescent development of executive function," *Translational psychiatry*, vol. 3, pp. e238-e238, 2013.
- [9] M. A. Hahn, D. Heib, M. Schabus, K. Hoedlmoser, and R. F. Helfrich, "Slow oscillation-spindle coupling predicts enhanced memory formation from childhood to adolescence," *Elife*, vol. 9, p. e53730, 2020.
- [10] S. L. Master, M. K. Eckstein, N. Gotlieb, R. Dahl, L. Wilbrecht, and A. G. Collins, "Distangling the systems contributing to changes in learning during adolescence," *Developmental cognitive neuroscience*, vol. 41, p. 100732, 2020.
- [11] N. Ramnani and A. M. Owen, "Anterior prefrontal cortex: insights into function from anatomy and neuroimaging," *Nature reviews neuroscience*, vol. 5, pp. 184-194, 2004.
- [12] M. A. Just, "Brain Imaging and rTMS Studies of Individual Differences in Cognitive Processing," CARNEGIE-MELLON UNIV PITTSBURGH PA DEPT OF PSYCHOLOGY 2013.
- [13] J. B. Arden, *Rewire your brain: Think your way to a better life*: John Wiley & Sons, 2010.
- [14] O. J. Lieberman, A. F. McQuirt, G. Tang, and D. Sulzer, "Roles for neuronal and glial autophagy in synaptic pruning during development," *Neurobiology of disease*, vol. 122, pp. 49-63, 2019.
- [15] R. M. O'Connor and J. F. Cryan, "Adolescent brain vulnerability and psychopathology through the generations: role of diet and dopamine," *Biological psychiatry*, vol. 75, pp. 4-6, 2014.
- [16] K. L. Mills, A.-L. Goddings, L. S. Clasen, J. N. Giedd, and S.-J. Blakemore, "The developmental mismatch in structural brain maturation during adolescence," *Developmental neuroscience*, vol. 36, pp. 147-160, 2014.
- [17] S. J. Ordaz, W. Foran, K. Velanova, and B. Luna, "Longitudinal growth curves of brain function underlying inhibitory control through adolescence," *Journal of Neuroscience*, vol. 33, pp. 18109-18124, 2013.
- [18] H. Ebbinghaus, "Memory: A contribution to experimental psychology," *Annals of neurosciences*, vol. 20, p. 155, 2013.
- [19] A. Manthiou, J. Kang, S. S. Hyun, and X. X. Fu, "The impact of brand authenticity on building brand love: An investigation of impression in memory and lifestyle-congruence," *International Journal of Hospitality Management*, vol. 75, pp. 38-47, 2018.
- [20] S. A. Josselyn and P. W. Frankland, "Infantile amnesia: a neurogenic hypothesis," *Learning & Memory*, vol. 19, pp. 423-433, 2012.
- [21] G. Kartikeswar, T. B. Parikh, D. Pandya, and A. Pandit, "Ionizing Radiation Exposure in NICU," *The Indian Journal of Pediatrics*, vol. 87, pp. 158-160, 2020.
- [22] L. J. Otten and M. D. Rugg, "The birth of a memory," *Trends in neurosciences*, vol. 25, pp. 279-281, 2002.
- [23] C. Hitchcock, J. Newby, E. Timm, R. M. Howard, A.-M. Golden, W. Kuyken, *et al.*, "Memory category fluency, memory specificity, and the fading affect bias for positive and negative autobiographical events: Performance on a good day-bad day task in healthy and depressed individuals," *Journal of Experimental Psychology: General*, vol. 149, p. 198, 2020.
- [24] P. J. Olesen, J. Macoveanu, J. Tegnér, and T. Klingberg, "Brain activity related to working memory and distraction in children and adults," *Cerebral cortex*, vol. 17, pp. 1047-1054, 2007.
- [25] G. Lynch, "Memory enhancement: the search for mechanism-based drugs," *nature neuroscience*, vol. 5, pp. 1035-1038, 2002.
- [26] P. B. Paulus, V. L. Putman, K. L. Dugosh, M. T. Dzindolet, and H. Coskun, "Social and cognitive influences in group brainstorming: Predicting production gains and losses," *European review of social psychology*, vol. 12, pp. 299-325, 2002.
- [27] D. F. Stramaccia, A.-K. Meyer, K. M. Rischer, J. M. Fawcett, and R. G. Benoit, "Memory suppression and its deficiency in psychological disorders: A focused meta-analysis," *Journal of Experimental Psychology: General*, 2020.
- [28] Z. Shipstead, T. S. Redick, and R. W. Engle, "Is working memory training effective?," *Psychological bulletin*, vol. 138, p. 628, 2012.
- [29] M. V. Johnston, A. Ishida, W. N. Ishida, H. B. Matsushita, A. Nishimura, and M. Tsuji, "Plasticity and injury in the developing brain," *Brain and Development*, vol. 31, pp. 1-10, 2009.
- [30] P. R. Huttenlocher, *Neural plasticity*: Harvard University Press, 2009.
- [31] D. L. Dunning, K. Griffiths, W. Kuyken, C. Crane, L. Foulkes, J. Parker, *et al.*, "Research Review: The effects of mindfulness-based interventions on cognition and mental health in children and adolescents—a meta-analysis of randomized controlled trials," *Journal of Child Psychology and Psychiatry*, vol. 60, pp. 244-258, 2019.
- [32] M. Buschkuhl, S. M. Jaeggi, and J. Jonides, "Neuronal effects following working memory training," *Developmental cognitive neuroscience*, vol. 2, pp. S167-S179, 2012.
- [33] C.-s. R. Li and R. Sinha, "Inhibitory control and emotional stress regulation: Neuroimaging evidence for frontal-limbic dysfunction in psycho-stimulant addiction," *Neuroscience & Biobehavioral Reviews*, vol. 32, pp. 581-597, 2008.
- [34] J. B. Harborne, H. Marby, and T. Marby, *The flavonoids*: Springer, 2013.
- [35] V. Dietz, "Neuronal plasticity after a human spinal cord injury: positive and negative effects," *Experimental neurology*, vol. 235, pp. 110-115, 2012.
- [36] R. Malinow and R. C. Malenka, "AMPA receptor trafficking and synaptic plasticity," *Annual review of neuroscience*, vol. 25, pp. 103-126, 2002.

- [37] M. T. Borrello, B. Schinor, K. Bartels, H. Benelkebir, S. Pereira, W. T. Al-Jamal, *et al.*, "Fluorinated tranlycypromine analogues as inhibitors of lysine-specific demethylase 1 (LSD1, KDM1A)," *Bioorganic & Medicinal Chemistry Letters*, vol. 27, pp. 2099-2101, 2017.
- [38] S. Farashahi, C. H. Donahue, P. Khorsand, H. Seo, D. Lee, and A. Soltani, "Metaplasticity as a neural substrate for adaptive learning and choice under uncertainty," *Neuron*, vol. 94, pp. 401-414. e6, 2017.
- [39] K. Iigaya, "Adaptive learning and decision-making under uncertainty by metaplastic synapses guided by a surprise detection system," *Elife*, vol. 5, p. e18073, 2016.
- [40] P. Khorsand and A. Soltani, "Optimal structure of metaplasticity for adaptive learning," *PLoS computational biology*, vol. 13, p. e1005630, 2017.
- [41] Q. Wu, H. Wang, Q. Luo, W. Banerjee, J. Cao, X. Zhang, *et al.*, "Full imitation of synaptic metaplasticity based on memristor devices," *Nanoscale*, vol. 10, pp. 5875-5881, 2018.