

A Survey Study of Some Graph Labeling Techniques.

V. T. Dhokrat¹, P. G. Jadhav², and S.B.Gaikwad³

¹Department of Mathematics, New Art's, Commerce and Science College, Ahmednagar, Maharashtra, India.

²Department of Mathematics, Dnyaneshwar Gramonnati Mandal's Hon. B.J.Art's, Comm. and Sci. College, Ale Phata, Ta: Junnar Dist: Pune, Maharashtra, India

³Department of Mathematics, New Art's, Comm. and Sci. College, Ahmednagar, Maharashtra, India.

¹Email: vibhadhokrat@gmail.com, pgjmaths@gmail.com, sbgmathsnagar@gmail.com

Abstract

Graph theory is a well-known branch of Mathematical sciences (Arithmetic). Mathematics gives the same name to different things. But graph theory creates a graph labeling techniques in which different labeling are given to same graphical structures. Graph is a mathematical structure describing dots, curves, bars, or traces. In Graph labeling natural numbers including zero are used to label the nodes or edges, or all, which fulfill some mathematical constraints. Graphs are used to describe various mathematical models in Operations Research, control, and Engineering, in studies domains of lap-top technological know-how which include statistics mining, photograph segmentation, clustering, photograph capturing, networking that's used in structural fashions, and so forth. Graph labeling has many applications in the social community, verbal exchange (communication) community, circuit design, Database management, coding principle, radar, astronomy, and X-ray crystallography. Depending on the trouble scenario a type of graph is used for representing the hassle (problem) and by way of applying appropriate graph labeling techniques the hassle may be solved. Graph labeling is a flourishing as well as application-oriented area of research in Mathematics.

In this chapter, we discuss the different graph labeling techniques related to different graphs.

Keywords- Graph, Graph labeling, Cordial, Magic, Mean, Radio, Power mean, Permutation, combination.

2010 Mathematics Subject Classification: Primary 05C22, 05C78, 05C38, 05C76.

1 Introduction

Graph concept has been invented in 1735 with the Koinsberg Seven Bridge problem. The phrase graph is derived from the Greek word 'graphein'. A "graph" in this paper consists of "vertices" or "nodes" and lines called edges that connect with vertices. Graph labeling have been introduced within the mid 1960's by Alexander Rosa. Over the past six decades, the research in graph labeling developed very fast. Graph labeling is a flourishing as well as an application-oriented area of research in the theory of Graphs. To date, more than 200 types of graph labeling techniques had been introduced. In this chapter we try to collect some important graph labeling techniques with suitable graphs. Here we study mainly graph without loops and parallel edges, non-directed having countable number of edges and vertices. Here we consider the terminology and symbols of graph from Harary's 'Graph Theory'.

1.1 Graph :

Definition: 1. A linear graph (simple graph) $G = (N(G), L(G), F(G))$, consisting of two non empty sets viz: $N(G) = \{v_1, v_2, \dots\}$ whose elements are known as nodes or vertices, and a set of elements which are known as lines or edges $L(G) = \{e_1, e_2, \dots\}$. The incidence function $F(G)$ which maps every element of E i.e. $e = xy$ to two

elements x and y of $N(G)$.

Order of a graph: The total count of elements i.e. vertex set $N(G)$ is called order of a graph.


Graph Size: The total count of an elements in edge set $L(G)$ is called graph size.

Vertex degree:The total number of lines occurring on a vertex is called a vertex degree .

Graph degree:The sum of all degrees of each node(vertex) in a given graph G is known as graph degree.

1.2 Graph Labeling :

Definition: 2. In a Graph, an allotment of positive integers including zero to the nodes or positive integers to the lines(edges), or both, having some mathematical criterion is known as Graph labeling.




1.3 Different Graph Labeling Techniques :

1.3.1 Graceful Labeling:

Definition: 3. A graceful labeling of a simple connected graph $G = (N(G), L(G))$ having p nodes and q lines is a particular graph labeling in which the nodes or vertices are labeled with a subset of distinct integers from 0 to q and the graph lines or edges are labeled with the positive differences between vertex label values. The labeling for edges used are from 1 to q both must be included. The resultant graph is known a graceful.


Exa. Banana tree ,trees with vertices ≤ 35 ,Caterpillars,bananas etc. are always graceful.



Examples of graph having Graceful Labeling.

- All symmetrical trees
- Trees with vertices ≤ 35 .
- Caterpillars, banana trees etc.

Non-graceful graph: A simple graph G in which we can not make graceful labeling is said to be non-graceful.



Result

In graceful graph with q (q is positive integer) edges and $q + 1$ vertices the following holds:

- We can draw distinct graceful labeling for the same graph i.e. it is not unique.
- The vertices labeled with 0 and q are always in neighbouring.
- If the graph has q edges then each graceful labeling must contains vertex label as q .
- Graceful labeling graph may contains a triangle graph.
- The complementarity property for graceful labeling is satisfied. That is for a given graph with graceful labeling if we swap every vertex label q with $q - k$, the resulting labeling is also graceful since the edge labels will not have changed the extreme vertices of an edge. The new labeling for a and b now are $q - a$ and $q - b$.


1.3.2 Harmonious labeling:

Graham and Sloane [33]

Definition: 4. In a graph $G = (N(G), L(G))$ with q lines, a function g is called as harmonious labeling of G if there exists an injective function f from vertex/node set $N(G)$ to $L(G)$ and the induced function g^* from $N(G)$ to $L(G)$ defined by $g^*(e = xy) = (g(x) + g(y)) \pmod{q}$ must be bijective.

A graph is called a harmonious graph if it has harmonious labeling .

Exa. $K_{1,7}$



Examples of graph having Harmonious Labeling.

- Cycle $C_n (n \geq 3)$ has harmonious labeling if n is an odd number and vice versa.
- Ladder graph $L_n, n \neq 2$.
- F_n , Friendship graph excluding n is congruent to $2 \pmod{4}$.
- The fan graph f_n .

- The graph g_n for $(n \geq 2)$.


Result

- Harmonious labeling is non distinctive.
- h is harmonious labeling then $xh(a) + y$ also a harmonious labeling , here the inverse of an element x in q exist and y is any arbitrary element in q . (i.e. Z_q).
- We can label 0 to any vertex in a graph.
- All trees are harmonious. non-distinctive
- Complete graph C_n is harmonious iff $n \leq 4$.
- The Peterson graph is Harmonious.
- Wheel graph $W_n = C_n + K_1$ is harmonious.
- $K_p^{(2)}$ has harmonious labeling for $p=4$. $K_p^{(2)}$ has not harmonious labeling for $n = 6$ or an odd number. If $p = 4$, $K_p^{(2)}$ is harmonious, if $p = 6$ it is not harmonious..

1.3.3 Cordial Labeling :

Cahit [9]

Definition: 5. Let $G = (N(G),L(G))$ be a connected, simple graph with p nodes/vertices q edges/lines.If in this graph , if it is possible to label its vertices with 0's and 1's such that the number of vertices labeled with 0 and the number of vertices labeled with 1 differ by at most 1 and similarly the number of edges labeled with 0 and the number of edges labeled 1 differ at most by 1.Then such a labeling is called cordial labeling Exa. Cordial labeling graph of Jelly Fish $J_{5,5}$



Examples of standard graph having Cordial Labeling.

- All trees.
- Complete graph K_n iff n is less than or equal to 3.
- Complete bipartite graph $K_{p,q}$ where p and q are natural numbers.
- Friendship graph $C_3^{(s)}$ iff s is not congruent to 2 modulo 4.
- All fan graph f_n .
- mK_n for particular values of m and n .

- Skolem-graceful graph .

Types of Cordial Labeling.

- Difference cordial graph.
- Edge product cordial graph.
- Prime cordial graph.
- Planar grid cordial graph.
- Context of Duplication cordial graphs.
- Second order cordial labeling graph.
- SET cordial graph.
- Integer cordial graph.
- Signed product cordial graph.
- Mean cordial graph.
- Harmonic Mean cordial graph.


1.3.4 Magic labeling of a graph:

There is a well-known concept of magic squares in number theory , Sedlacek in 1963 use this concept to discover a new labeling techniques known as magic labeling. [41]. **a) Node magic labeling of a graph(VML):**

Definition: 6. Let $G = (N(G), L(G))$ be a simple, connected graph having p nodes q lines. A graph G is said to be node magic if there exists a bijective function $M: N(G) \cup L(G) \rightarrow \{1, 2, \dots, p + q\}$, such that for all nodes of G , the sum of label on a node and the labels of its incident lines is constant such a bijection is called node magic labeling of G .

A graph that admits node magic labeling is called a node magic graph.

Exa.cycle C_8 having VML with magic constant $K=22$.




b) Line magic labeling of a graph

Definition: 7. Let $G = (N(G), L(G))$ be a simple, connected graph having p nodes q lines. A graph G is said to be line magic if there exists a bijective function $M: N(G) \cup L(G) \rightarrow \{1, 2, \dots, p + q\}$, such that $M(u) + M(v) + M(uv)$ is constant for all edges uv .

A graph is a line magic graph if it has an line magic labeling .

If the sum of all labels of lines incident on a node v is the same for all nodes v then such a connected graph is said to be semi-magic .

cycle graph C_6 with magic constant $K=20$.



Cycle C_6

Examples of a graph having Magic Labeling.

- For $p = 2$ and p greater than or equal to 5, K_p .
- $K_{p,p}$ for all p greater than or equal to 3.
- Fan graph f_n iff $n=$ odd and n greater than or equal to 3.
- The wheel graph on p vertices i.e. W_p for $p \geq 4$.
- A connected (m,n) graph iff $5m/4 < n \leq m(m - 1)/2$.

Types of Magic Labeling.

- Semi-Magic
- Super-Magic
- Anti-magic
- Prime-magic
- H-magic
- Sigma labeling/ Distance-magic labeling

1.3.5 Radio labeling:


Chartrand, Erwin, Zhang, and Harary [15]

Radio labeling of graph is used for synchronization of channel allotments to FM radio stations.

Definition: 8. Let $G = (N(G), L(G))$ be a simple, connected graph having p nodes q lines. A radio labeling of a connected graph G is an injective function c from the node set $N(G)$ to the set of natural numbers such that $d(x,y) + |c(x) - c(y)| \geq 1 + \text{diam}(G)$ for every two distinct nodes x and y of G .

The radio number denoted by $rn(c)$ is the greatest value allotted by radio labeling c to any node of G by radio labeling c .

The smallest value of radio numbers $rn(c)$ taken from radio labeling c of G is the radio number of G and is denoted by $rn(G)$.



Cycle C_8


Examples of Radio Labeling graph.

- The $rn(G)$ of a sunflower graph SF_n is equal to its order.
- The $rn(G)$ of a Helm graph H_n is an odd no. i.e. $2n + 1$.
- The $rn(G)$ of a gear graph G_n is also an odd no. i.e. $2n + 1$.

1.3.6 Mean labeling of graphs:

Somasundaram and Ponraj [40].

Definition: 9. Let $G = (N(G), L(G))$ be a simple, connected graph having p nodes q lines. A graph G is said to be a mean graph if there is a one-onemapping M from the nodes of G to $\{0, 1, 2, \dots, q\}$ such that when each edge (r, s) is labeled with $(M(r) + M(s))/2$ if $M(r) + M(s)$ is even and $(M(r) + M(s) + 1)/2$ if $M(r) + M(s)$ is odd, the resultant line labels are easily distinguishable.



$F_{2,4}$

Examples of Mean Labeling graph.

- The graph P_n, C_n , the bipartite graph $K_{2,n}$, triangular snakes, quadrilateral snakes etc. all are mean graphs..
- K_p iff p is less than 3.
- $K_{1,r}$ iff r is less than 3.
- The friendship graph $C_3^{(t)}$ iff t less than 2.
- Bistars $B_{p,q} (p > q)$ iff $p < q + 2$

Types of Mean Labeling graph.


- Vertex even and odd mean graph.
- Super graph.

1.3.7 Irregular labeling of graph:

Ruiz, Chartrand, Saba, Lehel, Oellermann and Jacobson. [14]

Definition: 10. Let $G = (N(G), L(G))$ be a simple, connected graph having p nodes q lines. An irregular labeling of a graph G without zero degree nodes is an allotment of natural numbers as weights to the lines of G in such a way that the sums of all the weights of the lines incident at every node are disjoint.

The lowest of the greatest weight of lines over all irregular labeling of a particular graph is known as the irregularity strength of G and is denoted by $s(G)$. If no such weight exists for an irregular labeling then $s(G) = \infty$.




An irregular 3-labeling of the wheel W_3

1.3.8 Prime labeling of graph:

Tout, Dabboucy, and Howalla. [20]

Definition: 11. Let $G = (N(G), L(G))$ be a simple, connected graph having p nodes q lines. A graph G is said to possess a prime labeling if there is an injective and surjective mapping $P: (N(G) \rightarrow \{1, 2, \dots, q\})$ such that for each edge (r, s) , $\gcd(P(r), P(s)) = 1$ i.e. $(P(r)$ and $P(s)$ are relatively prime.



R_8


Examples of Prime Labeling of graph.

- Caterpillar graph with maximum degree 5.
- Path graph, star graph, complete binary trees, spider graph.

1.3.9 Power Mean Labeling of graph:

Mercy P.[37]

Definition: 12. Let $G = (N(G), L(G))$ be a simple, connected graph having p nodes q lines. A graph G is said to be a Power Mean Graph if it is feasible to label the vertices $r \in N(G)$ with distinct labels P from $\{1, 2, 3, \dots, m+1\}$ i.e. there is an one-one function $P: N(G) \rightarrow \{1, 2, \dots, q\}$ in such a way that when each edge $e = rs$ is labeled with $P(e = rs) = \lfloor (r)^{P(s)} * P(s)^{P(r)} \frac{1}{P(r)+P(s)} \rfloor$ then the line labels are distinct and are from $\{1, 2, 3, \dots, m\}$. F is called Power mean labeling of G .



Examples of Power Mean Labeling of a graph.

- Tadpoles $T(n, k)$.

Result:

- When two cycles C_p and C_q are joined by a path P_q , the resultant graph is a Power mean graph
- $T(P_m)$ for $m \geq 2$ is a Power mean graph.
- Subdivision of any P_n for $n \geq 2$ is a Power mean graph.
- Subdivision of any C_n for n greater than or equal to 3 is a Power mean graph.


1.3.10 Combination and Permutation labeling of graph:

Hegde and Shetty [20]

a) Combination labeling of graph:

Definition: 13. Let $G = (N(G), L(G))$ be a simple, connected graph having p nodes q lines. A function $C: N(G) \rightarrow \{1, 2, \dots, q\}$ is called as combination labeling of graph G if every line (r, s) is labeled with $(C(r))! / [(C(r) - C(s))!(C(s))!]$ where $C(r) > C(s)$.

The resulting line labels are all well-defined.




Examples of Combination Labeling of a graph.

- K_n if and only if n is less than or equal to 5.
- C_n , n greater than 3.
- $K_{n,n}$ iff n is less equals to 2. .

b) Permutation Labeling of graph:

Definition: 14. Let $G = (N(G), L(G))$ be a simple, connected graph having p nodes q lines. A function $P: N(G) \rightarrow \{1, 2, \dots, m\}$ is called as permutation labeling of graph G if every edge (r, s) is labeled with $(P(r))!/[P(r) - P(s)]!$ where $P(r) > P(s)$. The resulting line labels all are well-defined.



Examples of Permutation Labeling of a graph.

- K_p iff p is less equals 5.


Result:

- In a permutation graph G if we remove any edge from this graph then the resultant graph is again a permutation graph.
- The graph k -wheel, for all integer $n \geq 2$.
- The graph k -fan $F_{n,k}$.
- The gear graph G_{2n} is also a permutation graph.

1.3.11 Triangular sum labeling of graph:

Hegade and Shankaran[36]

Definition: 15. Let $G = (N(G), L(G))$ be a simple, connected graph having p nodes and q lines. A labeling of the graph with q lines is called a triangular sum labeling if the nodes can be determined with distinct not-negative integers in such a way that, when an line whose end vertices are labeled with r and s is labeled with the value $r + s$. All vertex(node) and edge(line) labels are different.



Examples of Triangular sum Labeling of graph.

- Star graph, Path graph, n -ary complete trees.
- K_p iff $p = 1$ or $p = 2$

References

- [1] M. E. Abdel-Aal, *New classes of odd graceful graphs*, *Internat. J. Appl. Graph Theory in Wireless ad hoc Networks and Sensor Networks*, 5(2) June 2013.
- [2] J. Abrham and A. Kotzig, *Extensions of graceful valuations of 2-regular graphs consisting of 4-gons*, *Ars Combin.*, 32 (1991) 257-262.
- [3] B. D. Acharya, K. A. Germina, K. Abhishek, and P. J. Slater, *Some new results on set-graceful and set-sequential graphs*, *Combin. System Sci.*, 37(2-4) (2012) 229-239
- [4] B. D. Acharya, S. B. Rao, T. Singh, and V. Parameswaran, *Neighborhood magic graphs*, 2004, unpublished.
- [5] B. D. Acharya and S. M. Hegde, *Strongly indexable graphs*, *Discrete Math.*, 93 (1991) 123-129.
- [6] V. Ajitha, S.Arumugum and K.A. Gremina, *On square sum graphs*, *AKCE International Journal of graphs and Combinatorics*, Volume 6(1) (2009),1-10.
- [7] B. Beavers, *Golomb rulers and graceful graphs*, http://webcourse.cs.technion.ac.il/236801/Spring2009/ho/WCFiles/Golomb_Ruler_Gracefu_Graphs.pdf
- [8] L. W. Beineke and S. M. Hegde, *Strongly multiplicative graphs*, *Discuss. Math. Graph Theory*, 21 (2001) 63-75.
- [9] I. Cahit, *Cordial graphs: a weaker version of graceful and harmonious graphs*, *Ars Combin.*, 23 (1987) 201-207.
- [10] ahit, *On harmonious tree labellings*, *Ars Combin.*, 41 (1995) 311-317.
- [11] . Cattell, *Graceful labellings of paths*, *Discrete Math.*, 307 (2007) 3161-3176.
- [12] . J. Chang, D. F. Hsu, and D. G. Rogers, *Additive variations on a graceful theme: some results on harmonious and other related graphs*, *Congr. Numer.*, 32 (1981) 181-197.
- [13] G. Chartrand, S. M. Lee, and P. Zhang, *Uniformly cordial graphs*, *Discrete Math.*, 306 (2006) 726-737.
- [14] G. Chartrand, M. Jacobson, J. Lehel, O. Oellermann, S. Ruiz, and F. Saba, *Irregular networks*, *Congr. Numer.*, 64 (1988) 187-192.
- [15] G. Chartrand, D. Erwin, and P. Zhang, *A graph labeling problem suggested by FM channel restrictions*, *Bull. Inst. Combin. Appl.*, 43 (2005) 43-57.
- [16] G. Chartrand, D. Erwin, P. Zhang, and F. Harary, *Radio labelings of graphs*, *Bull. Inst. Combin. Appl.*, 33 (2001) 77-85.
- [17] A. Durai Baskar, S. Arockiaraj, and B. Rajendran, *F-geometric mean labeling of some chain graphs and thorn graphs*, *Kragujevac J. Math.* 37(1) (2013) 163-186.
- [18] H. L. Fu and K. C. Huang, *On prime labelling*, *Discrete Math.*, 127 (1994) 181-186.
- [19] Y. Fukuchi, *Edge-magic labelings of wheel graphs*, *Tokyo J. Math.*, 24 (2001) 153-167.
- [20] Gallian, JA 2010, 'A dynamic survey of graph labeling', *The Electronic Journal of Combinatorics*, vol.17 (DS6).
- [21] J. A. Gallian, *Labeling prisms and prism related graphs*, *Congr. Numer.*, 59 (1989) 89-100.
- [22] J. A. Gallian, *A survey: recent results, conjectures and open problems on labelinggraphs*, *J. Graph Theory*, 13 (1989) 491-504.
- [23] J. A. Gallian, *Open problems in grid labeling*, *Amer. Math. Monthly*, 97 (1990)133-135.
- [24] J. A. Gallian, *A guide to the graph labeling zoo*, *Discrete Appl. Math.*, 49 (1994)213-229.

- [25] J. A. Gallian and D. S. Jungreis, *Labeling books*, *Scientia*, 1 (1988) 53-57.
- [26] J. A. Gallian, J. Prout, and S. Winters, *Graceful and harmonious labelings of prisms and related graphs*, *Ars Combin.*, 34 (1992) 213-222.
- [27] J. A. Gallian and L. A. Schoenhard, *Even harmonious graphs*, *AKCE J. Graphs Combin.*, 11(1) (2014) 27-49.
- [28] J. A. Gallian and D. Stewart, *Properly even harmonious labelings of disconnected graphs*, *AKCE J. Graphs Combin.*, 12(2-3) (2015) 193-203.
- [29] J. A. Gallian and D. Stewart, *Even harmonious labelings of disjoint graphs with a small component*, *AKCE J. Graphs Combin.*, 12(2-3) (2015) 204-215.
- [30] J. A. Gallian and D. Stewart, *Even harmonious labelings of disjoint unions with even sequential graphs*, *J. Graph Labeling*, 1(1) (2015) 1-10.
- [31] T. Grace, *Graceful, Harmonious, and Sequential Graphs*, Ph. D. Thesis, University Illinois at Chicago Circle, 1982.
- [32] T. Grace, *On sequential labelings of graphs*, *J. Graph Theory*, 7 (1983) 195-201
- [33] R. L. Graham and N. J. A. Sloane, *On additive bases and harmonious graphs*, *SIAM J. Alg. Discrete Methods*, 1 (1980) 382-404.
- [34] Harary, F 1988, 'Graph Theory', Narosa publishing House Reading, New Delhi.
- [35] N. Hartsfield and G. Ringel, *Pearls in Graph Theory*, Academic Press, San Diego, 1990.
- [36] S. M. Hegde and P. Shankaran, *On triangular sum labelings of graphs*, in *Labeling of Discrete Structures and Applications*, Narosa Publishing House, New Delhi, 2008, 109-115.
- [37] Mercy P., *Power mean series*, Ph.D. Thesis.
- [38] M. Mollard, C. Payan, and S. Shixin, *Graceful problems*, *Seventh Hungarian Colloquium on Finite and Infinite*.
- [39] M. A. Razaq, S. T. Rizvi, and K. Ali, *H-groupmagic total labelings of families of fan graphs*, *Util. Math.*, 108 (2018) 213219
- [40] Rosa, A 1967, 'On certain valuations of the vertices of a graphs', *Theory of graphs*, (Internet, symposium, Rome, July 1966), Gordon and Breach, N.Y. and Dunod Paris, pp.349-355.
- [41] S. S. Sandhya, S. Somasundaram, R. Ponraj, *Some results on harmonic mean graphs*, *Int. J. Contemp. Math. Sci.*, 7 (2012) 197-208.
- [42] J. Sedlacek, *Problem 27*, in *Theory of Graphs and its Applications*, Proc. Symposium Smolenice, June, (1963) 163.
- [43] S. K. Vaidya and C. M. Barasara, *Product cordial labeling of line graph of some graphs*, *Kragujevac J. Math.*, 40(2) (2016) 290-297.
- [44] S. Somasundaram and R. Ponraj, *Mean labelings of graphs*, *Natl. Acad. Sci. Let.*, 26 (2003) 210-213.
- [45] S. Somasundaram, P. Vidhyarani and R. Ponraj, *Geometric mean labelings of graphs*, *Bull. Pure Appl. Sci.*, 30E(2) (2011) 153-160.