

AN OVERVIEW OF *MORINGA OLEIFERA* PODS, SEEDS, BARK AND ROOT

Muthu Reka.S¹ and S.S. Vijayanchali²

¹ Ph. D Full Time Research Scholar, Department of Home Science, The Gandhigram Rural Institute (Deemed to be University), Gandhigram, Dindigul Dist., Tamil Nadu, India.

E-mail Id - muthurekasankar@gmail.com

²Professor, Department of Home Science, The Gandhigram Rural Institute (Deemed to be University), Gandhigram, Dindigul Dist., Tamil Nadu, India.

*Corresponding Author: muthurekasankar@gmail.com

Abstract

Moringa oleifera is also called as moringa, drumstick leaves, miracle tree. *Moringa oleifera* belongs to *Moringaceae* family. *Moringa oleifera* contains numerous health benefits. *Moringa oleifera* is a rich source of specific essential nutrients, bioactive components and phytochemicals like sterols, tannins, flavonoids, terpenoids, anthraquinones, saponins, alkaloids, isothiocyanates and glycoside compounds. *Moringa oleifera* is also beneficial for skin, hair, liver, eye, blood pressure, treating anemia, kidney disease, and diabetes. The main focus of this review is about the health benefits of *Moringa oleifera* seeds, bark, pod and root.

Keywords: *Moringa oleifera*, drumstick leaves, pods, seeds, bark, root, antioxidant

Introduction

Plants are rich in secondary metabolites that can be used to treat a wide range of illnesses and problems. These phytoconstituents are often obtained from various plant components such as the leaf, flower, bark, twig, stem, tuber and they act directly as medicinal agents (Cragg & Newman, 2001).

Moringa oleifera is a member of the *Moringaceae* family, is one of the most effective medicinal herbs. Native to India, Southern America, Africa, Europe, Sub-Himalayan regions, Asia, and the Middle East, it grows in tropical and subtropical climates (Pakade et al., 2013), (Alegbeleye, 2018), (Raman et al., 2018), (Vickers, 2017). The *moringa oleifera* tree, which grow to a height of 10 to 12 m, is a small, quickly-growing evergreen tree (Athira et al., 2021).

Moringa oleifera is nutritionally dense due to the presence of several important compounds in its leaves, pods, and seeds (Mahmood et al., 2010), (Rockwood et al., 2013). *Moringa oleifera* is a well-known plant that is abundant in essential minerals and

phytochemicals(Gupta & Soin, 2020). Numerous bioactive phytochemicals such as flavonoids, saponin, vanillin, omega fatty acids, carotenoids, ascorbates, tocopherols, β -sitosterol, kaempferol, and quercetin can be found in *Moringa oleifera's* flowers, roots, fruits, and seeds and have a wide range of medical applications(Ma et al., 2020).

Moringa oleifera is known to have hypoglycemic, antibacterial, antifungal, antioxidant and anti-inflammatory properties (Choudhary et al., 2013; Sutorialangka et al., 2013),(Oladeji et al., 2020). Due to its significant medicinal properties, it might be substituted for coriander leaves to increase flavour (Gandji et al., 2018).

Moringa oleifera has also been used to modify the immune system in pathological situations including anaemia, anxiety, asthma, blood impurities and cholera. It has also been used to treat malnutrition, edema, bacterial contaminations, viral infections, hyperglycemia, and cancer(Hamza, 2010),(Saini et al., n.d.). *Moringa oleifera* is also good for the skin, hair, eyes, liver(Meireles et al., 2020).

***Moringa oleifera* Pods**

Moringa oleifera pods are brown, pendulous, triangular in shape and divided into three sections according to length. 30-120cm long and 1.8cm broad when completely dried. Fruit production takes place mostly in March and April. Each fruit has 26 seeds at the beginning of its growth. Pods are green while juvenile and become brown when they are grown(Islam et al., 2021).

Moringa oleifera pods are incredibly nutrient-dense and include phytochemicals, vitamins, minerals, and amino acids. A good dose of protein, ash, non-structural carbohydrates, lipids, and fiber is found in *Moringa oleifera* pods. Fatty acids such as oleic, linoleic, palmitic, and linolenic acids are also present. Vitamins C and A are very abundant in the pods. The complex chemical components found in *Moringa oleifera* pods include antibacterial and anti-inflammatory capabilities, which help strengthen the immune system's innate defense(Prentki & Nolan, 2006).

Moringa oleifera immature pods have 19.34% protein, 1.28% fats, 46.78% fibre,30% amino acids and 24.98% nonstructural carbs(Sánchez-Machado et al., 2010). According to one research, one cup of fresh, sliced pods (100 g) contains 157% of an adult's daily need of ascorbic acid. It is also used to extend the shelf life of food and to store it. Because pods

contain anti-helminthic effects, they are used to treat spleen and liver infections. It is also used to alleviate articular discomfort(Masih et al., 2019).

The pods of *Moringa oleifera* also guard against bacterial activity, cancer, oxidative stress, antidiabetic, inflammation, hepatic fibrosis, liver damage, hypercholesterolemia, and other harmful conditions(SWATIq et al., 2018),(Shah et al., 2022).

***Moringa oleifera* Seeds**

Moringa oleifera seed was harmless and suggested using it as a coagulant in developing nations (Okereke & Akaninwor, 2013). *Moringa oleifera* seeds are spherical shape semi-permeable seed shell. Three wings on the hull itself are spaced 120 degrees apart from top to bottom. Each tree may yield 15,000 to 25,000 seeds annually(Shah et al., 2022).

Seeds feature a mid-brown seed core with three light wings. The seed core can be dark brown or black in colour, although it can also be milky. *Moringa oleifera* has a variety of bioactive categories, including alkaloidal steroidal, phenolic compounds, flavonoids, terpenes, zeatin, quercetin and different carbohydrates such as L-arabinose, L-galactose, mannose, and xylose, among other phytonutrients(Stadtlander & Becker, 2017).

The high levels of protein, minerals and unsaturated fatty acids, primarily oleic acid is an excellent source in *Moringa oleifera* seeds make them valuable raw materials for culinary applications(Milla et al., 2021), (Özcan, 2020). According to recent studies, *Moringa oleifera* seed powder increases the nutritional content of a variety of foods, including bread, cookies, cakes, snacks, and cereal gruels(Rabie et al., 2020).

The seeds of the *Moringa oleifera* seed are a good source of antioxidant activity, lipids, proteins, and minerals like calcium, magnesium, zinc(Ijarotimi et al., 2013), (Liang et al., 2019). *Moringa oleifera* seeds' nutritional analysis revealed that they included a range of protein concentrations (9.98–51.80 g/100 g), ash concentrations (3.60–5.0 g/100 g), fibre concentrations (17.26–20.0 g/100 g), and carbohydrate concentrations (3.6–18.0 g/100 g)(Busani et al., 2011).

Moringa oleifera seeds consist of 28.02% crude protein, 33.78% crude fat, 28.77% carbs, 94.74 mg/100 g vitamin C, 2.84 mg/100 g calcium, and 129.03 mg/100 g sodium(Igwilo et al., 2017). The amount of cystine and methionine in *Moringa oleifera* seeds is comparable to that of eggs and milk. *Moringa oleifera* seed protein is highly digestible (93%) due to urease activity and the absence of trypsin inhibitors. As a natural flocculent, *Moringa oleifera* seed

powder is used to clear murky water(Masih et al., 2019). *Moringa oleifera* seed extract lowers lipid peroxides in the liver and contains antihypertensive substances(Ali et al., 2022).

A clinical trial is being conducted to assess the effectiveness of *Moringa oleifera* seed kernels in the treatment of bronchial asthma. For three weeks, 20 patients with mild to severe asthma were given 3 kg of dried seed kernels. The results demonstrated that *Moringa oleifera* extracts significantly reduced the intensity of asthma symptoms while simultaneously improving lung function metrics(Agrawal & Mehta, 2008).

Patients with bronchial asthma can benefit from using *Moringa oleifera* seeds as a treatment. 3g daily for three weeks is the recommended dose. It produced positive effects such a decrease in the erythrocyte sedimentation rate, an increase in the heartbeat value symptoms score, and a decrease in the intensity of asthma episodes. Enforced expiratory volume, vital capacity, and expiratory flow rate values all improved after 3 weeks of therapy. *Moringa oleifera* seed alcohol extract has spasmolytic effects on histamine- and acetylcholine-induced bronchospasm(Paikra & Gidwani, 2017).

Polyunsaturated fatty acids abound in *Moringa oleifera* seeds. Aside from that, the existence of phytochemical substances is responsible for medicinal plants' therapeutic effectiveness(Mahfuz & Piao, 2019). Due to the presence of benzyl isothiocyanate and moringine the seeds of this plant have antibacterial potential(Kamran et al., 2020).

In addition to being used to season food, *Moringa oleifera* seeds can also be used to purify water. In some regions, they are also consumed as roasted nuts(James & Zikankuba, 2017). The seeds of the *Moringa oleifera* plant are used as a purgative and to treat hysteria, tumors and scurvy (Posmontier, 2011). In addition to acting as antibacterial and anti-inflammatory agents, seeds of *Moringa oleifera* are used in the treatment of hyperthyroidism, antiheroes-simplex virus arthritis, rheumatism, gout cramp, epilepsy, and sexually transmitted infections(Ram & Rastogi, 2004).

***Moringa oleifera* Bark**

Bark extract in aqueous form lowers the weight of stones formed using 1% ethylene glycol. *Moringa oleifera* bark has curative and preventative potential (Sekhar et al., 2018). The bark is used as an aphrodisiac and to treat upper respiratory tract infections(Posmontier, 2011).

It was demonstrated in a clinical trial involving 30 patients that 40 mL of a *Moringa oleifera* stem bark decoction taken twice daily was beneficial in treating the signs and symptoms of a

urinary tract infection(Maurya & Singh, 2014). With the use of several *Moringa oleifera* bark extracts, anticholesteremic and antilipidemic efficacy in rats was also confirmed(Senecha et al., 2012).

The bark of the *Moringa oleifera* plant is used to treat eye problems and stop the growth of tuberculous glands in the neck and spleen, eliminate tumors, and treat ulcers. The root bark juice is applied to the ears to relieve earaches and is also used as a pain reliever in dental cavities(Anwar et al., 2007), (Qi et al., 2019).

***Moringa oleifera* Root**

As a spice *Moringa oleifera* root is eaten. The antibacterial, anticancer, antifungal, antiulcer, and anti-inflammatory actions of roots are also present (Shank et al., 2013; Steve & Babatunde, 2013), (Abubakar & Usman, 2016), (Abd-Rabou et al., 2017).

The effects of *Moringa oleifera* root extract on foot edema formation in rats were equivalent to those of the analgesic and antipyretic nonsteroidal anti-inflammatory drug phenylbutazone(Sandeep et al., 2019). *Moringa oleifera* root has anti-ulcer, anti-inflammatory, and cardiac-stimulating properties(Aderinola et al., 2020).

Indians utilize the roots of the *Moringa oleifera* plant to cure ailments including edema, wounds, ulcers, hiccups, asthma, rheumatism, dyspepsia, gout, eye problem, kidney stones, and illnesses of the liver and spleen(Fuglie, 2002).

Conclusion

From the review it is concluded that the *Moringa oleifera* has numerous health benefits. *Moringa oleifera* has high amount of antioxidant, phytochemicals, bioactive components. *Moringa oleifera* is a rich source of carbohydrate, protein, fat, vitamins like A and C, and minerals like calcium, sodium and potassium. *Moringa oleifera* is also beneficial for skin, hair, liver, eye, blood pressure, treating anemia, kidney disease, and diabetes.

References

- Abd-Rabou, A. A., Abdalla, A. M., Ali, N. A., & Zoheir, K. M. A. (2017). *Moringa oleifera* root induces cancer apoptosis more effectively than leave nanocomposites and its free counterpart. *Asian Pacific Journal of Cancer Prevention: APJCP*, 18(8), 2141.
- Abubakar, I., & Usman, A. (2016). *Phytochemical and antibacterial investigations of moringa (Moringa oleifera) leaf extract on selected bacterial pathogens.*

- Aderinola, T. A., Alashi, A. M., Nwachukwu, I. D., Fagbemi, T. N., Enujiugha, V. N., & Aluko, R. E. (2020). In vitro digestibility, structural and functional properties of Moringa oleifera seed proteins. *Food Hydrocolloids*, *101*, 105574.
- Agrawal, B., & Mehta, A. (2008). Antiasthmatic activity of Moringa oleifera Lam: A clinical study. *Indian Journal of Pharmacology*, *40*(1), 28.
- Alegbeleye, O. O. (2018). How functional is Moringa oleifera? A review of its nutritive, medicinal, and socioeconomic potential. *Food and Nutrition Bulletin*, *39*(1), 149–170.
- Ali, M. Y., Khalil, M. I., Jahan, F. N., Hossain, M. B., & Samanta, A. K. (2022). Moringa oleifera: A review on nutritional attributes, therapeutic applications and value-added product generation: A review on nutritional attributes of moringa oleifera. *SAARC Journal of Agriculture*, *20*(2), 1–15.
- Anwar, F., Latif, S., Ashraf, M., & Gilani, A. H. (2007). Moringa oleifera: a food plant with multiple medicinal uses. *Phytotherapy Research: An International Journal Devoted to Pharmacological and Toxicological Evaluation of Natural Product Derivatives*, *21*(1), 17–25.
- Athira, K. A., Panjikaran, S. T., Aneena, E. R., Sharon, C. L., Pathrose, B., Gopal, K. S., & Lakshmi, P. S. (2021). Moringa Oleifera-Proximate and Anti-nutritional Composition. *The Indian Journal of Nutrition and Dietetics*, 390–397.
- Busani, M., Patrick, J. M., Arnold, H., & Voster, M. (2011). Nutritional characterization of Moringa (Moringa oleifera Lam.) leaves. *African Journal of Biotechnology*, *10*(60), 12925–12933.
- Choudhary, M. K., Bodakhe, S. H., & Gupta, S. K. (2013). Assessment of the antiulcer potential of Moringa oleifera root-bark extract in rats. *Journal of Acupuncture and Meridian Studies*, *6*(4), 214–220.
- Cragg, G. M., & Newman, D. J. (2001). Natural product drug discovery in the next millennium. *Pharmaceutical Biology*, *39*(sup1), 8–17.
- Fuglie, L. J. (2002). The Tree of Life. *Dakar, Senegal, Western Africa: Church World Services*, 40–60.
- Gandji, K., Chadare, F. J., Idohou, R., Salako, V. K., Assogbadjo, A. E., & Kakai, R. L. G. (2018). Status and utilisation of Moringa oleifera Lam: A review. *African Crop Science Journal*, *26*(1), 137–156.
- Gupta, D., & Soin, D. (2020). Recent Advances in Health Benefits of Moringa Oleifera. *International Journal of Pharmaceutical Sciences and Nanotechnology (IJPSN)*, *13*(3), 4870–4875.
- Hamza, A. A. (2010). Ameliorative effects of Moringa oleifera Lam seed extract on liver fibrosis in rats. *Food and Chemical Toxicology*, *48*(1), 345–355.
- Igwilo, I. O., Ugochukwu, G. C., Ezekwesili, C. N., & Nwenyi, V. (2017). COMPARATIVE STUDIES ON THE NUTRIENT COMPOSITION AND ANTINUTRITIONAL

FACTORS IN DIFFERENT PARTS OF MORINGA OLEIFERA PLANT FOUND IN AWKA, NIGERIA. *The Bioscientist Journal*, 5(1), 1–12.

- Ijarotimi, O. S., Adeoti, O. A., & Ariyo, O. (2013). Comparative study on nutrient composition, phytochemical, and functional characteristics of raw, germinated, and fermented Moringa oleifera seed flour. *Food Science & Nutrition*, 1(6), 452–463.
- Islam, Z., Islam, S. M., Hossen, F., Mahtab-ul-Islam, K., Hasan, M. R., & Karim, R. (2021). Moringa oleifera is a prominent source of nutrients with potential health benefits. *International Journal of Food Science*, 2021.
- James, A., & Zikankuba, V. (2017). Moringa oleifera a potential tree for nutrition security in sub-Saharan Africa. *American Journal of Research Communication*, 5(4), 1–14.
- Kamran, M., Hussain, S., Abid, M. A., Syed, S. K., Suleman, M., Riaz, M., Iqbal, M., Mahmood, S., Saba, I., & Qadir, R. (2020). Phytochemical composition of moringa oleifera its nutritional and pharmacological importance. *Postepy Biologii Komorki*, 47(3), 321–334.
- Liang, L., Wang, C., Li, S., Chu, X., & Sun, K. (2019). Nutritional compositions of Indian Moringa oleifera seed and antioxidant activity of its polypeptides. *Food Science & Nutrition*, 7(5), 1754–1760.
- Ma, Z. F., Ahmad, J., Zhang, H., Khan, I., & Muhammad, S. (2020). Evaluation of phytochemical and medicinal properties of Moringa (Moringa oleifera) as a potential functional food. *South African Journal of Botany*, 129, 40–46.
- Mahfuz, S., & Piao, X. S. (2019). Application of Moringa (Moringa oleifera) as natural feed supplement in poultry diets. *Animals*, 9(7), 431.
- Mahmood, K. T., Mugal, T., & Haq, I. U. (2010). Moringa oleifera: a natural gift-A review. *Journal of Pharmaceutical Sciences and Research*, 2(11), 775.
- Masih, L. P., Singh, S., Elamathi, S., Anandhi, P., & Abraham, T. (2019). Moringa: A multipurpose potential crop-A review. *Proc. Indian Natl. Sci. Acad*, 85, 589–601.
- Maurya, S. K., & Singh, A. K. (2014). Clinical efficacy of Moringa oleifera Lam. stems bark in urinary tract infections. *International Scholarly Research Notices*, 2014.
- Meireles, D., Gomes, J., Lopes, L., Hinzmann, M., & Machado, J. (2020). A review of properties, nutritional and pharmaceutical applications of Moringa oleifera: integrative approach on conventional and traditional Asian medicine. *Advances in Traditional Medicine*, 20(4), 495–515.
- Milla, P. G., Peñalver, R., & Nieto, G. (2021). Health benefits of uses and applications of Moringa oleifera in bakery products. *Plants*, 10(2), 318.
- Okereke, C. J., & Akaninwor, J. O. (2013). The protein quality of raw leaf, seed and root of Moringa oleifera grown in Rivers State, Nigeria. *Annals of Biological Research*, 4(11), 34–38.

- Oladeji, O. S., Odelade, K. A., & Oloke, J. K. (2020). Phytochemical screening and antimicrobial investigation of Moringa oleifera leaf extracts. *African Journal of Science, Technology, Innovation and Development*, 12(1), 79–84.
- Özcan, M. M. (2020). Moringa spp: Composition and bioactive properties. *South African Journal of Botany*, 129, 25–31.
- Paikra, B. K., & Gidwani, B. (2017). Phytochemistry and pharmacology of Moringa oleifera Lam. *Journal of Pharmacopuncture*, 20(3), 194.
- Pakade, V., Cukrowska, E., & Chimuka, L. (2013). Metal and flavonol contents of Moringa oleifera grown in South Africa. *South African Journal of Science*, 109(3), 1–7.
- Posmontier, B. (2011). The medicinal qualities of Moringa oleifera. *Holistic Nursing Practice*, 25(2), 80–87.
- Prentki, M., & Nolan, C. J. (2006). Islet β cell failure in type 2 diabetes. *The Journal of Clinical Investigation*, 116(7), 1802–1812.
- Qi, L., Zhou, Y., Li, W., Zheng, M., Zhong, R., Jin, X., & Lin, Y. (2019). Effect of Moringa oleifera stem extract on hydrogen peroxide-induced opacity of cultured mouse lens. *BMC Complementary and Alternative Medicine*, 19, 1–9.
- Rabie, M. M., Ibrahim, F. Y., Youssif, M. R. G., & Ezz El-Ragal, N. M. (2020). Effect of Moringa oleifera leaves and seeds powder supplementation on quality characteristics of cookies. *Journal of Food and Dairy Sciences*, 11(2), 65–73.
- Ram, P., & Rastogi, B. N. (2004). Mehrotra. *Compendium of Indian Medicinal Plants. Central Drug Research Institute, Lucknow and National Institute of Science Communication and Information Resources, New Delhi*, 4, 484.
- Raman, J. K., Alves, C. M., & Gnansounou, E. (2018). A review on moringa tree and vetiver grass–Potential biorefinery feedstocks. *Bioresource Technology*, 249, 1044–1051.
- Rockwood, J. L., Anderson, B. G., & Casamatta, D. A. (2013). Potential uses of Moringa oleifera and an examination of antibiotic efficacy conferred by M. oleifera seed and leaf extracts using crude extraction techniques available to underserved indigenous populations. *International Journal of Phytotherapy Research*, 3(2), 61–71.
- Saini, R. K., Sivanesan, I., & Keum, Y. S. (n.d.). Phytochemicals of Moringa oleifera: A Review of Their Nutritional. *Therapeutic and Industrial Significance*, 3.
- Sánchez-Machado, D. I., Núñez-Gastélum, J. A., Reyes-Moreno, C., Ramírez-Wong, B., & López-Cervantes, J. (2010). Nutritional quality of edible parts of Moringa oleifera. *Food Analytical Methods*, 3, 175–180.
- Sandeep, G., Anitha, T., Vijayalatha, K. R., & Sadasakthi, A. (2019). Moringa for nutritional security (Moringa oleifera Lam.). *International Journal of Botany Studies*, 4(1), 21–24.
- Sekhar, C., Venkatesan, N., Muruganathi, D., & Vidhyavathi, A. (2018). Status of value addition and export of moringa produce in Tamil Nadu-a case study. *International Journal of Horticulture*, 8.

- Senecha, C., Shama, P. K., D'Souza, U. P., & Shastry, C. S. (2012). Anticholesteremic and antilipidemic activity of stem bark extracts of *Moringa oleifera* in diet induced hyperlipidemia model in rats. *International Journal of Pharmaceutical and Chemical Sciences*, 1(3), 567–574.
- Shah, L., Baxi, N., Shah, A., & Akhiani, T. K. (2022). *An Overview of the Health Benefits of Drumstick Pods*.
- Shank, L. P., Riyathong, T., Lee, V. S., & Dheeranupattana, S. (2013). Peroxidase activity in native and callus culture of *Moringa oleifera* Lam. *J Med Bio Eng*, 2(3).
- Steve, I. O., & Babatunde, O. I. (2013). Chemical compositions and nutritional properties of popcorn-based complementary foods supplemented with *Moringa oleifera* leaves flour. *Journal of Food Research*, 2(6), 117.
- Sutalangka, C., Wattanathorn, J., Muchimapura, S., & Thukham-mee, W. (2013). *Moringa oleifera* mitigates memory impairment and neurodegeneration in animal model of age-related dementia. *Oxidative Medicine and Cellular Longevity*, 2013.
- SWATIġ, A. K., Kumari, C., Ali, A., Garg, P., Thakur, P., Attri, C., & Kulshrestha, S. (2018). *Moringa oleifera*—A never die tree: An overview. *Asian J Pharm Clin Res*, 11(12), 57–65.
- Vickers, N. J. (2017). Animal communication: when i'm calling you, will you answer too? *Current Biology*, 27(14), R713–R715.