

Ecological Consequences of Harmful Algal Blooms

Jayashri Mahadev Swamy ^a, Preeti ^b Somnath R. Yadav^c

^a Directorate of Research Karnataka Veterinary Animal and Fisheries Science University
Bidar 585226

^c Maharashtra Animal and Fishery Sciences University, Nagpur, Maharashtra 4400001, India.

^b Veterinary Gynaecology and Obstetrics. Karnataka Veterinary Animal and Fisheries
Sciences University Bidar

Abstract

This book chapter delves into the complex dynamics of harmful algal blooms (HABs) and their far-reaching implications in marine ecosystems. Highlighting the critical role of microscopic planktonic algae as a primary food source for crustaceans, fish, and shellfish, the narrative explores the dual nature of algal blooms, oscillating between advantageous and detrimental impacts on aquaculture and wild fisheries. The chapter underscores the global prevalence of harmful algal blooms, defined by the prolific expansion of specific microalgae, macroalgae, or cyanobacteria. Recognizing the escalation of HABs as intricately linked to climate change, nutrient enrichment, and habitat disturbance, the authors elucidate the diverse causes behind their occurrence. Focusing on categories of bloom-forming algae, the chapter details their distinctive characteristics, impacts on aquatic ecosystems, and associated health risks, including various toxins such as paralytic shellfish poisons and neurotoxic shellfish poisons. The discussion further delves into the environmental forces driving HABs, including eutrophication, climate change, and ocean acidification. The consequences of HABs span phycotoxin accumulation, toxin transfer, high biomass blooms, ecosystem disturbance, aerosolized toxins, and localized hazardous blooms. In advocating for a nuanced understanding of the multifaceted contributors to HABs, the chapter emphasizes the need for effective mitigation and management strategies. A holistic approach that considers both anthropogenic influences and broader ecological contexts is crucial to curtail the expanding impact of HABs and restore balance to coastal ecosystems globally.

Key Words: Harmful algal blooms, Categories, Consequences, Mitigation

Introduction

In the expansive realm of the world's oceans, microscopic planktonic algae stand as a vital sustenance source for the early stages of economically consequential crustaceans and fish, along with serving the nutritional needs of filter-feeding bivalve shellfish, including oysters, mussels, scallops, and clams. The growth dynamics of plankton algae, colloquially referred to as "algal blooms" when reaching densities of millions of cells per liter, play a pivotal role, oscillating between being advantageous and detrimental for both aquaculture and wild fisheries operations.

Numerous scholarly sources (Hallegraeff 1993; Anderson 2012; Bresnan et al. 2013; GlobalHAB, 2017) collectively define algal blooms as the prolific expansion of specific microalgae, macroalgae, or cyanobacteria (blue/green algae). In favorable environmental conditions, these blooms can attain levels that impact aquatic life and, in turn, pose consequences for human well-being. Harmful algal blooms (HABs) have been documented globally, exerting a detrimental influence on tourism, recreation, aquaculture, and fisheries (Berdalet *et al.*, 2016; Hallegraeff, 2003). Recognizing the rich diversity of marine phytoplankton, Sournia *et al.* (1991) identified 5,000 species. However, a mere 80 species possess the capacity to produce toxic substances, affecting humans through fish and shellfish consumption. Intriguingly, non-toxic species experiencing high-biomass blooms can also serve as sources of HABs.

The aftermath of algal blooms extends beyond the proliferation phase, as the shaded benthic environment witnesses a drop in oxygen levels following the bloom's decay. The resultant accumulation of algal biomass may induce water discoloration, imparting negative impacts that reverberate across the ecosystem, ultimately leading to a reduction in biodiversity.

The escalating frequency of HABs is intricately linked to climate change, nutrient enrichment, and habitat disturbance (Hallegraeff 1993; GESAMP, 2001; Smayda 2004; Anderson 2012; Berdalet et al. 2016). This increase in HAB occurrences carries profound negative effects, ranging from the poisoning or asphyxiation of finfish, shellfish, and human consumers to severe consequences such as fish, bird, and mammal (including human) deaths. Afflictions of the respiratory or digestive tract, memory loss, seizures, lesions, and skin irritation are among the most pronounced and memorable impacts of HABs. Additionally, coastal resources, including submerged aquatic vegetation and benthic epi- and in-fauna, undergo significant losses. As we delve into the scientific exploration of these phenomena, this

chapter aims to unravel the intricate dynamics of harmful algal blooms and their far-reaching implications.

Causes of Harmful Algal Blooms:

Harmful algal blooms (HABs) represent a pervasive ecological challenge, encompassing the unchecked proliferation of autotrophic algae and heterotrophic protists, a phenomenon increasingly observed in coastal waterways across the globe. The alarming surge in HAB occurrences serves as a poignant indicator of an ecological imbalance, a consequence of the intricate interplay between the expanding global human footprint and the pervasive influence of climate change.

At the heart of this burgeoning issue lie two overarching causes: eutrophication, a byproduct of human activities, and a suite of natural processes, including circulation, upwelling relaxation, and river movement. Eutrophication, driven by nutrient loading from agricultural runoff, sewage discharge, and industrial activities, provides an enriched substrate for algal growth, fueling the proliferation of harmful species. Simultaneously, natural phenomena such as oceanic currents and river dynamics contribute to the spatial and temporal distribution of these blooms, creating a complex tapestry of environmental factors influencing their prevalence.


Historically, the escalation of HABs has coincided with episodes wherein environmental conditions approach those conducive to maximum algal growth. This dynamic relationship underscores the intricate interplay between climate variations and the intensification of algal blooms (Moore et al., 2009; Gobler et al., 2017; Anderson et al., 2012). The intricate web of interacting physical, chemical, and biological factors further complicates this scenario, influencing the blooming of specific species across diverse habitats (Zingone and Enevoldsen, 2000).

Additionally, the geographic expansion of certain algal species reflects a complex interplay of factors, including anthropogenic introductions, natural occurrences such as ocean currents, and the historical presence of species at undetectable concentrations or in the distant past. This dynamic landscape of causes underscores the imperative for a nuanced understanding of the multifaceted contributors to HABs.

In navigating this intricate web of causation, there arises a pressing need for effective mitigation and management strategies. A holistic and scientifically informed approach is vital,

acknowledging the diverse factors at play and charting a course towards sustainable solutions that address both anthropogenic influences and the broader ecological context. Only through such nuanced comprehension can we hope to curtail the expanding impact of HABs and restore balance to coastal ecosystems worldwide.

Factors Affecting Harmful Algae Gains and Losses



Harmful Algal Groups:

In the realm of harmful algal blooms (HABs), five primary categories of bloom-forming algae take center stage, each with distinctive characteristics and impacts on aquatic ecosystems.

Dinoflagellates: Dinoflagellates, unicellular microalgae, are key contributors to the primary production of planktonic algae in water bodies and are frequently implicated in harmful algal bloom (HAB) occurrences globally (Burkholder et al., 2006). Exhibiting a range of nutritional strategies, dinoflagellates can be heterotrophic, autotrophic, or mixotrophic. Reproducing primarily through asexual fission, some species engage in sexual reproduction, leading to the formation of dormant cysts capable of withstanding environmental stresses (Smayda, 1997).

Notably, dinoflagellates produce toxins that pose health risks to humans through the consumption of contaminated seafood or direct contact with HABs. Responsible for several types of seafood poisoning, including paralytic shellfish poisoning (PSP), neurotoxic shellfish

poisoning (NPS), ciguatera fish poisoning (CFP), and diarrhetic shellfish poisoning (DSP), dinoflagellates exert a considerable impact on public health (Smayda, 1997; Van Dolah, 2000).

Other Flagellates: Flagellates, specifically Prymnesiophyceae and Raphidophyceae classes, contribute to harmful algal blooms with significant human impact. Prymnesiophytes, like *Prymnesium parvum* and *Chrysochromulina polylepis*, cause fish-kills. *Phaeocystis* spp., another prymnesiophyte, produces mucilaginous material, causing anoxic events that killed 10 million kg of mussels in the Oosterschelde estuary in 2001. Raphidophytes, including *Pseudochattonella verruculosa*, have triggered large-scale fish kills in northern Europe since 1998, highlighting the complex dynamics and widespread consequences of harmful flagellate blooms in marine ecosystems (Edwardsen and Imai, 2006; Smayda, 2006; Peperzak and Poelman, 2008; de Boer, 2006).

Diatoms: Diatoms, unicellular photosynthetic eukaryotes, represent approximately 100 species thriving in freshwater and marine environments globally. Playing a vital role in carbon fixation and marine ecosystem productivity, marine diatoms contribute 20% of global carbon fixation and 40% of primary productivity in marine ecosystems. Known for their siliceous cell wall, or frustule, and spines that can harm fish gills, diatoms are essential yet can be lethal to aquatic life.

Some diatoms, such as *Pseudo-nitzschia* species, produce domoic acid, a neurotoxin responsible for Amnesic Shellfish Poisoning (ASP) in humans. Symptoms include confusion, headaches, short-term memory loss, and, in severe cases, death (Muhseen et al., 2015).

Cyanobacteria: Cyanobacteria, often referred to as "blue-green algae," constitute a diverse group of oxygenic photosynthetic bacteria with the ability to synthesize chlorophyll-a and auxiliary pigments like phycobilins (Francis, 1878). Inhabiting various water types, cyanobacteria exhibit resilience to extreme temperatures and demonstrate a higher affinity for nitrogen and phosphorus than other photosynthetic species, capitalizing on anthropogenic changes in aquatic environments (Chorus and Bartram, 1999).

Unlike other phytoplankton types, cyanobacteria, including species like *Nodularia*, *Aphanizomenon*, *Anabaena*, and *Trichodesmium*, thrive in warm, stratified conditions in both coastal and open ocean environments (Dadheech et al., 2013; Kleinteich et al., 2012). Their rapid growth in response to warming conditions sets them apart from other eukaryotic phytoplankton.

Filamentous Algae: Filamentous algae, encompassing green, red, and brown species like *Cladophora*, *Ectocarpus*, *Pilayella*, *Ulothrix*, *Ulva*, and *Urospora*, along with colonial diatoms like *Melosira*, trigger near-shore green tides. While non-toxic, they form dense overgrowth and drifting mats, outcompeting macrophytes, causing anoxia, and posing threats to fishing gear, boats, and tourism. Events occur notably in the Baltic Sea, Kattegat, Skagerrak, and Atlantic coast, impacting local ecosystems and tourism (Eriksson et al., 2009; Eriksson et al., 2011; Bonsdorff et al., 1997; Pihl et al., 1999; Vahteri et al., 2000; Baden et al., 2003; World Health Organisation, 2002).

These five harmful algal groups exemplify the diverse nature of HABs, emphasizing the intricate ecological dynamics and potential health risks associated with their presence in aquatic ecosystems. Understanding the characteristics and impacts of these algal groups is crucial for developing effective management and mitigation strategies.

Harmful Algal blooms and toxins

In addition to the dinoflagellates, other groups such as raphid ophytes, diatoms, cyanobacteria, and several others also have species that produce toxins.

HAB toxins are mostly categorised by symptoms into the following groups:

- Paralytic shellfish poisons (PSP)
- Neurotoxic shellfish poisons (NSP)
- Amnesic shellfish poisons (ASP),
- Diarrheic shellfish poisons (DSP),
- Azaspiracid shellfish poisoning (AZP),
- Ciguatera fish poisoning (CFP) and
- Cyanobacteria toxin poisoning (CTP)

Poison	Toxin-producing organism	Toxin(s)	Foods likely to be contaminated	Symptoms
Ciguatera fish poisoning (CFP)	Dinoflagellates: <i>Gambierdiscus toxicus</i>	Ciguatoxins, Maitotoxin, Scaritoxin	Reef fish such as barracuda, grouper, red	Nausea, Vomiting, Diarrhea, Stomach pain

			snapper, and amberjack	
Neurotoxic shellfish poisoning (NSP)	Dinoflagellates: <i>Karenia brevis</i> and other <i>Karenia</i>	Brevetoxins	Shellfish, primarily mussels, oysters, scallop	Nausea, Vomiting, Diarrhea, Stomach pain, Numbness of lips, tongue, and throat, Dizziness
Paralytic shellfish poisoning (PSP)	Dinoflagellates: <i>Gyrodinium aureolum</i> , <i>Pyrodinium bahamense</i> , <i>Alexandrium</i>	Saxitoxins	Shellfish, primarily scallop, mussels, clams, oysters, and cockles, Some fish and crabs	Nausea, Vomiting, Diarrhea, Shortness of breath, Irregular heartbeat, Numbness of mouth and lips, Weakness
Amnesiac shellfish poisoning (ASP)	Diatoms: <i>Pseudo-nitzschia</i>	Domoic acid	Shellfish, primarily scallop, mussels, clams, oysters, Possibly some fish species	Nausea, Vomiting, Diarrhea, Stomach pain, Shortness of breath, Memory loss, Disorientation, Possibly coma
Diarrhetic shellfish poisoning (DSP)	Dinoflagellates: <i>Dinophysis</i> species, <i>Prorocentrum lima</i>	Okadaic acid	Shellfish, primarily scallop,	Nausea, Vomiting, Diarrhea, Stomach pain,

			mussels, clams, oysters	Possibly chills, Headache, Fever
Azaspiracid shellfish poisoning (AZP)	Dinoflagellates: <i>Proroperidinium</i>	Azaspiracid	Shellfish	Nausea, Vomiting, Diarrhea, Stomach pain

[\(https://www.cdc.gov/habs/\)](https://www.cdc.gov/habs/)

Environmental Forcing of Harmful Algal Blooms:

The evolution of forecasting tools and mitigation strategies has emerged from extensive research on the ecological processes governing Harmful Algal Blooms (HABs) and the identification of key factors contributing to their global proliferation, as outlined by the Ecological Risks of Harmful Algal Blooms (GEOHAB, 2003, 2006).

Eutrophication:

One of the foremost environmental drivers of HABs is eutrophication, characterized by increased biomass resulting from nutrient over-enrichment in coastal waters. The ecological response to eutrophication is intricate, contingent upon macro- and micronutrient concentrations, the chemical form of nutrients (organic vs. inorganic), and nutrient supply ratios (Anderson et al., 2002, 2008; Heisler et al., 2008; Glibert and Burkholder, 2011; Kudela et al., 2010). Eutrophication not only poses threats to human health but also triggers fish and shellfish die-offs, sporadic deaths of marine mammals and birds, and substantial economic losses for coastal communities. These losses arise from expenses related to beach cleanup, closure of commercially significant areas, and reduced tourism and fisheries (Paerl 1988; Shumway 1990; Villac et al. 1993; Horner et al. 1997; Turner and Tester 1997). The assessment of ecological risks is crucial for understanding the impact of harmful and toxic algae on coastal ecosystems and predicting HAB occurrences.

Climate Change: Another significant environmental force driving HABs is climate change. Changes in temperature, precipitation patterns, and ocean currents influence the distribution and intensity of algal blooms (GEOHAB, 2003, 2006; Smayda, 2008). The warming of coastal waters can create favorable conditions for the rapid growth of harmful algae, exacerbating their impact on marine ecosystems.

Ocean Acidification: Ocean acidification, resulting from increased carbon dioxide absorption by seawater, is an emerging factor influencing HAB dynamics. Altered acidity levels may affect the physiology and competitive advantage of certain algal species, potentially favoring the proliferation of harmful varieties (GEOHAB, 2003, 2006; Gobler and Sunda, 2012).

In evaluating the environmental impacts of HABs, six primary categories emerge, each associated with distinctive ecological consequences and threats to human health:

1. **Phycotoxin Accumulation:** Leading to the accumulation of toxins in suspension-feeding organisms like bivalve shellfish.
2. **Toxin Transfer and Respiratory Damage:** Causing harm to fish and other marine fauna through toxin transfer and damage to respiratory processes, particularly fish gills.
3. **High Biomass Blooms:** Depleting oxygen levels or inducing other detrimental effects.
4. **Ecosystem Disturbance:** Triggering disturbances with cascading impacts on species relationships within ecosystems.
5. **Aerosolized Toxins:** Producing toxins that can affect human respiratory health when aerosolized.
6. **Localized Hazardous Blooms:** Involving benthic or epiphytic microalgae with unique habitation, processes, and severity of adverse effects distinct from planktonic HABs.

Understanding these environmental forces and their implications is fundamental for the development of effective mitigation and management strategies in the face of the growing challenges posed by HABs.

Conclusion:

This comprehensive exploration of harmful algal blooms (HABs) emphasizes the intricate interplay of ecological factors driving their occurrence. The escalating frequency of HABs, linked to climate change, nutrient enrichment, and habitat disturbance, poses significant threats to marine ecosystems and human well-being. The chapter underscores the urgent need for nuanced mitigation and management strategies, acknowledging both anthropogenic influences and broader ecological contexts. By understanding the diverse causes and impacts of HABs, the scientific community is better equipped to develop sustainable solutions, mitigate adverse effects, and restore balance to coastal ecosystems on a global scale.

Reference

Hallegraeff, G. M. (2003). Harmful algal blooms: a global overview. *Manual on harmful marine microalgae*, 33, 1-22.

Sellner, K.G., Doucette, G.J. & Kirkpatrick, G.J. (2003), Harmful algal blooms: causes, impacts and detection. *J IND MICROBIOL BIOTECHNOL*, 30, 383–406.

Brown, A. R., Lilley, M., Shutler, J., Lowe, C., Artioli, Y., Torres, R. ... Tyler, C. R. (2019). Assessing risks and mitigating impacts of harmful algal blooms on mariculture and marine fisheries. *Reviews in Aquaculture*. doi:10.1111/raq.12403

Anderson DM (2012) HABs in a changing world: a perspective on harmful algal blooms, their impacts, and research and management in a dynamic era of climactic and environmental change. *Harmful Algae* 2012: 3–17

Hallegraeff GM (1993) A review of harmful algal blooms and their apparent global increase. *Phycologia* 32(2): 79–99.

Berdalet, E., Fleming, L.E., Gowen, R., Davidson, K., Hess, P., Backer, L.C., Moore, S.K., Hoagland, P., Enevoldsen, H., 2016. Marine harmful algal blooms, human health and wellbeing: challenges and opportunities in the 21st century. *J. Mar. Biol. Assoc. United Kingdom* 96 (1), 61–91.

Bresnan, E., Ar´evalo, F., Belin, C., Branco, M.A.C., Cembella, A.D., Clarke, D., Correa, J., Davidson, K., Dhanji-Rapkova, M., Fern´andez-Lozano, R., Fernandez-Tejedor, ´ M., Guðfinnsson, H., Ja´en-Carbonell, D.J., Laza-Martinez, A., Lemoine, M., Lewis, A.M., Mam´an-Men´endez, L.M., Maskrey, B.H., McKinney, A., Pazos, Y., Revilla, M., Siano, R., Silva, A., Swan, S., Turner, A.D., Schweibold, L., Provoost, P., Enevoldsen, H., accepted for publication. Harmful algae - diversity and regional distribution of harmful algal events along the Atlantic margin of Europe. *Harmful Algae*.

Karlson, B., Andersen, P., Arneborg, L., Cembella, A., Eikrem, W., John, U., ... Suikkanen, S. (2021). Harmful algal blooms and their effects in coastal seas of Northern Europe. *Harmful Algae*, 102, 101989.

SOURNIA, A.; CHRETIENNOT-DINET, M. J.; RICARD, M. 1991. Marine phytoplankton: how many species in the world ocean? *J. Plankton Res.*, No. 13, pp. 1093–9.

Hallegraeff GM (1993) A review of harmful algal blooms and their apparent global increase. *Phycologia* 32(2): 79–99.

GESAMP (2001) Group of Experts on the Scientific Aspects of Marine Pollution (GESAMP) Reports and Studies. United Nations Environment Programme, Nairobi: 2001. Global Environment Outlook 3, EARTH SCAN, Past, present and future perspectives. Protecting the oceans from land-based activities. Land-based sources and activities affecting the quality and uses of the marine, coastal and associated freshwater environment. No. 71.11

Smayda TJ (2004) Harmful algal bloom communities in Scottish coastal waters: relationship to fish farming and regional comparisons – a review. Scottish Executive Environment Group Paper 2006/3.

Anderson DM (2012) HABs in a changing world: a perspective on harmful algal blooms, their impacts, and research and management in a dynamic era of climactic and environmental change. *Harmful Algae* 2012: 3–17.

Berdalet E, Fleming LE, Gowen R, Davidson K, Hess P, Backer LC et al. (2016) Marine Harmful Algal Blooms, human health and wellbeing: challenges and opportunities in the 21st century. *Journal of the Marine Biology Association of the United Kingdom* 96 (Special Issue 01): 61.

Sellner, K. G., Doucette, G. J., & Kirkpatrick, G. J. (2003). Harmful algal blooms: causes, impacts and detection. *Journal of Industrial Microbiology and Biotechnology*, 30(7), 383–406.

Ismael, A.A., Succession of heterotrophic and mixotrophic dinoflagellates as well as autotrophic microplankton in the harbour of Alexandria, Egypt. *Journal of Plankton Research*, 2003. 25(2): p. 193-202.

Mohamed, Z.A. and A.M. Al-Shehri, Occurrence and germination of dinoflagellate cysts in surface sediments from the Red Sea off the coasts of Saudi Arabia. *Oceanologia*, 2011. 53(1): p. 121-136.

Smayda T.J., 1997. Harmful algal blooms: Their ecophysiology and general relevance to phytoplankton blooms in the sea. *Limnology and Oceanography* 42, 1137-1153.

Van Dolah F.M., 2000. Marine algal toxins: Origins, health effects, and their increased occurrence. *Environmental Health Perspectives* 108, 133-141.

Edwardsen B., Imai I., 2006. The ecology of harmful flagellates. In: Graneli E., Turner J.T. (editors). *Ecology of harmful algal blooms*. Springer.

Jiang, L., et al., Diatoms: a novel source for the neurotoxin BMAA in aquatic environments. *PLoS One*, 2014. 9(1): p. e84578.

Muhseen, Z.T., et al., Proteomics studies on stress responses in diatoms. *Proteomics*, 2015. 15(23-24): p. 3943-3953.

Pulido, O.M., Domoic acid toxicologic pathology: a review. *Mar Drugs*, 2008. 6(2): p. 180-219.

Lefebvre, K.A. and A. Robertson, Domoic acid and human exposure risks: a review. *Toxicon*, 2010. 56(2): p. 218-30.

Francis, G., Poisonous Australian Lake. *Nature*, 1878. 18(444): p. 11-12.

Dadheech, P.K., et al., Cyanobacterial diversity in the hot spring, pelagic and benthic habitats of a tropical soda lake. *FEMS MicrobiolEcol*, 2013. 85(2): p. 389-401. 39.

Kleinteich, J., et al., Temperature-related changes in polar cyanobacterial mat diversity and toxin production. *Nature Climate Change*, 2012. 2(5): p. 356-360.

Chorus, I. and J. Bartram, *Toxic Cyanobacteria in Water: A guide to their public health consequences, monitoring and management*. 1999.

Paerl, H.W., Mitigating harmful cyanobacterial blooms in a human- and climatically-impacted world. *Life (Basel)*, 2014. 4(4): p. 988-1012.

Elliott, J.A., The seasonal sensitivity of Cyanobacteria and other phytoplankton to changes in flushing rate and water temperature. *Global Change Biology*, 2010. 16(2): p. 864-876.

Conley, D.J., et al., Ecology. Controlling eutrophication: nitrogen and phosphorus. *Science*, 2009. 323(5917): p. 1014-5.

Kevin G Sellner and others, Harmful algal blooms: causes, impacts and detection, *Journal of Industrial Microbiology and Biotechnology*, Volume 30, Issue 7, 1 July 2003, Pages 383–406, <https://doi.org/10.1007/s10295-003-0074-9>

Zingone, A., &Enevoldsen, H.O., 2000. The diversity of harmful algal blooms: a challenge for science and management. *Ocean and Coastal Management* 43, 725–748.

Sellner, K. G., Doucette, G. J., & Kirkpatrick, G. J. (2003). Harmful algal blooms: causes, impacts and detection. *Journal of Industrial Microbiology and Biotechnology*, 30(7), 383–406.

Perl TM, Bedard L, Kosatsky T, et al. 1990. An outbreak of toxic encephalopathy caused by eating mussels contaminated with domoic acid. *New Engl J Med* 322:775-1780.

Shumway SE. 1990. A review of the effects of algal blooms on shellfish and aquaculture. *J World Aquaculture Soc* 21:65-104.

Villac MC, Roelke DL, Villareal TA, et al. 1993. Comparison of two domoic acid producing diatoms: A review. *Hydrobiologia* 269/270:213-24.

Horner RA, Garrison DL, and Plumley FG. 1997. Harmful algal blooms and red tide problems on the U.S. west coast. *LimnolOceanogr* 42:1076-88.

Tester PA and Steidinger KA. 1997. *Gymnodiniumbreve* red tide blooms: Initiation, transport, and consequences of surface circulation. *LimnolOceanogr* 42:1039-51.

Turner JT and Tester PA. 1997. Toxic marine phytoplankton, zooplankton grazers, and pelagic food webs. *LimnolOceanogr* 42:1203-14.

Gentien. P., Pitcher. G., Cembella. A.D., Glibert. P.M. (Eds.), 2003. GEOHAB: Global Ecology and Oceanography of harmful algal Blooms. SCOR and IOC , Baltimore and Paris.

Glibert. P.M., 2006. Global Ecology and Oceanography of harmful algal Blooms: Eutrophication and HABs. IOC and SCOR, Paris and Baltimore.

Anderson, D. M., Glibert, P. M., & Burkholder, J. M. (2002). Harmful algal blooms and eutrophication: Nutrient sources, composition, and consequences. *Estuaries*, 25(4), 704–726.

Anderson, D. M., Burkholder, J. M., Cochlan, W. P., Glibert, P. M., Gobler, C. J., Heil, C. A., Vargo, G. A. (2008). Harmful algal blooms and eutrophication: Examining linkages from selected coastal regions of the United States. *Harmful Algae*, 8(1), 39–53.

Heisler, J., Glibert, P. M., Burkholder, J. M., Anderson, D. M., Cochlan, W., Dennison, W. C., Suddleson, M. (2008). Eutrophication and harmful algal blooms: A scientific consensus. *Harmful Algae*, 8(1), 3–13.

Glibert, P.M., Burkholder, J.M. Harmful algal blooms and eutrophication: “strategies” for nutrient uptake and growth outside the Redfield comfort zone. *Chin. J. Ocean. Limnol.* 29, 724–738 (2011). <https://doi.org/10.1007/s00343-011-0502-z>

Kudela, R. M., Seeyave, S., & Cochlan, W. P. (2010). The role of nutrients in regulation and promotion of harmful algal blooms in upwelling systems. *Progress in Oceanography*, 85(1-2), 122–135. doi:10.1016/j.pocean.2010.02.008